

Original Research

Mitigation of Organ Specific Toxicity Following Acute Dinotefuran Exposure through Vitamin C Supplementation

H. Muhammad Ahmad¹, Sadia Nawaz¹, Muhammad Shahbaz Yousaf¹,
Taha Hassan², Muhammad Osama³, Rahat Naseer^{1*}

¹Institute of Biochemistry and Biotechnology, University of Veterinary and Animal Sciences, Lahore, Pakistan

²Hameed Latif teaching Hospital, Lahore, Pakistan

³Institute of Pharmaceutical Sciences, University of Veterinary and Animal Sciences, Lahore, Pakistan

Received: 11 September 2023

Accepted: 07 February 2024

Abstract

The objective of this research was to assess the impact of 2-methyl-2-nitro-3-guanidine on mammalian species, otherwise not the intended targets, followed by one-time exposure and its adverse effects. Thirty-five Sprague Dawley rats of either male or female sex, aged four weeks, and with an average weight of 80±20g, were divided into three groups randomly: control E exposed and exposed and supplemented with vitamin C, and assigned the titles C,E, and V, respectively. Each group comprised fifteen rats. Both E and V groups were further divided into subgroups: E1, E2, and E3, exposed to LD10, LD25, and LD50 doses of Dinotefuran respectively, and V1, V2, and V3, exposed in the same manner but supplemented with a fixed dose of ascorbic acid (vitamin C).

Ascorbic acid was administered in aqueous form 35mg/100mL of water and provided *ad libitum*. Eight hours after exposure to Dinotefuran, 5ml of blood was collected under sedation through the cardiac vein. After 48 hours, two rats randomly selected from each subgroup, including the control group, were anesthetized, euthanized, and dissected. Different body tissues, and the kidney, liver, bones, and heart, were isolated and preserved in formalin solution for subsequent analysis. CBC, liver, renal, and cardiac biomarkers were evaluated. In addition, histopathology and bone characteristics of soft tissues were also conducted. Mortality and morbidity were recorded.

The result showed significant disruptions in CBC and other biomarkers related to kidney, liver, and heart in a dose-dependent pattern. Although vitamin supplementation improved the overall outcome, the improvement was not statistically significant. Histopathological examination displayed changes in the exposed (E) group, with no observable improvements with vitamin supplementation. Moreover, the bone-related parameters exhibited similar trends.

Keywords: histopathology, insecticides, ascorbic acid, cellular toxicity

Introduction

Neonicotinoids are systemic insecticides with broad-spectrum efficacy and relatively low toxicity to mammals, fish, and birds. However, they have been linked to harmful effects on pollinating insects, including honey bees. Dinotefuran, a third-generation insecticide, is widely acceptable due to its efficacy and low toxicity to non-target species. Neonicotinoids mostly target the nervous system of insects by binding to specific receptors known as nicotinic acetylcholine receptors (*nAChRs*). These receptors are blocked upon binding with Dinotefuran, leading to paralysis and ultimately death in insects. It is reported that neonicotinoids may induce oxidative stress in cells, resulting in immunomodulatory effects and affecting the central nervous system of the target species [1, 2]. It's important to note that the impact of neonicotinoids is not limited to insects. Some studies have also suggested potential effects on non-target organisms, including birds, fish, and mammals [3]. However, the exact cellular mechanisms and long-term effects on these organisms are still being studied. Regulatory authorities and researchers continue to evaluate the risks and benefits of neonicotinoids to determine appropriate usage and minimize potential harm to pollinators and other organisms in the environment. Despite their widespread use, there is limited experimental data on the impact and mechanism of action of neonicotinoids on animals. This study aims to evaluate the effect of acute exposure to dinotefuran on different mammalian tissues.

Materials and Methods

Chemical Reagents

Dinotefuran [1-methyl-2-nitro-3-(tetrahydro-3-furyl methyl) guanidine] was provided by "Four Brothers" pesticide suppliers L. Ascorbic Acid (Merck), Formalin Buffer (Reidel-de Haen), Ethanol (VWR), distilled water, and Diethyl Ether (Merck Germany).

Methodology

Thirty-five Sprague Dawley rats, of male or female sex, weighing approximately 80±20 g, and aged around 4±0.5 weeks, were acclimatized in accordance with European Union guidelines for animal research at the Laboratory animal housing facility at UVAS, Lahore. The rats were kept in steel cages, and the temperature was maintained at 21±2°C with 50% humidity. They were provided with a basic diet and water *ad libitum* for a 12-hour adjustment period. Subsequently, the rats were distributed randomly into three groups, designated as the control, exposed, and vitamin supplemented groups. There were fifteen rats in each group and five in each subgroup (E1, E2, E3 & V1, V2, V3). Control

(Group C) also had five rats. The first subgroup of group E, E1, received LD₁₀ (400 mg/kg), the second subgroup received LD₂₅ (1000 mg/kg), and the third, E3, received LD₅₀ (2000 mg/kg). All the subgroups of "V" were exposed in the same manner as those of "E" and received a vitamin C supplement (100 mg/ 350 mL of water) additionally.

Twelve hours after the exposure, 5 mL of blood was collected directly from the cardiac puncture of randomly selected rats from each group. The blood sample was divided into two halves for further analysis of serum and whole blood.

Zero mortality was observed after 48 hours, but four rats belonging to different groups showed symptoms of anorexia and lethargy. All the rats were anesthetized and euthanized as per ethical guidelines for organ extraction. The bones and vital organs, except the brain, were preserved in formalin for subsequent analysis [4]. All procedures were conducted in accordance with the institution's code of animal research, ensuring humane practices.

Histopathological analysis of the tissues from the liver, kidney, and heart was performed using the method devised by [3] Cui et al. in 2009. The left tibia was removed, cleansed, and kept in boiling water for 10 minutes [4]. The weight/length index and robusticity index of the removed bone were determined using the following formulas:

$$\text{Length Index} = \frac{gm(\text{thigwe})}{\text{Length}(mm)}$$

Robusticity Index = $\frac{\text{Bone length}(mm)}{\sqrt[3]{\text{Bone Weight}}}$ [5]. All observations were collected in triplicates, and then the obtained data were analyzed using ANOVA in SPSS software. The p-value of each experimental parameter was also determined with $\{\alpha = 0.05\}$.

Results and Discussion

Zero mortality or weight loss symptoms were recorded after 48 hours of exposure. Similar results were also reported after the chronic exposure of Imidacloprid (10 and 20 mg/kg/60 days) [6] alone and also in combination with Fipronil [7].

Blood Toxicity

CBC was conducted on fifteen sampled rats after a duration of 12 hours. The results showed a significant decrease ($p < 0.05$) in the mean values of white blood cells (WBCs), MID cells, and granulocytes (GRA), while lymphocytosis (LYM) significantly increased ($p < 0.05$) in the experimental group, whereas the control group exhibited normal values of blood parameters. Vitamin C supplementation also demonstrated a significantly positive effect, as shown in Table 1.

Leukocytes and their types decreased in number, which can be attributed to the cellular environment

under oxidative stress, leading to individual cell lysis and depressed hematopoiesis [8]. Moreover, the transient rise in lymphocyte count may be the result of the body's defense mechanism [9, 10]. It should be noted that these findings are based on short-term data from an acute trial [11]. Another study on the sub-lethal toxicity of orally applied cypermethrin reported a significant reduction in leukocyte count, except for lymphocytes, and a promotion in lymphocyte count (lymphocytosis) in adult Sprague Dawley rats [12].

A significant decrease in the mean values of the following parameters, red blood cells (RBCs), hemoglobin (HGB), hematocrit (HCT), and red cell distribution width (RDWc) in the experimental group in comparison to the control group has been observed (Table 2). Conversely, values of the following parameters,

mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), and mean corpuscular hemoglobin concentration (MCHC) were also mounted up in the intoxicated rat group. Vitamin C supplementation demeaned the toxicity effect of dinotefuran on RBCs and related parameters. Factors such as internal hemorrhage, depletion in hemoglobin synthesis, and elevation in hemoglobin destruction have been reported to contribute to the reduction in the erythrocyte count and hemoglobin concentration [13- 16] Toxic substances like insecticides can produce effects in blood cells and lead to hemolysis [17]. Similar results were observed with cypermethrin and imidacloprid [18].

A slight downward trend was also shown by the mean values of platelets (PLT) and platelet percentage (PCT%) in the experimental groups as compared to

Table 1. Mean (\pm SD) values of WBCs and its types in different rat groups. E1 = Rat group fed with LD₁₀, E2 = Rat group fed with LD₂₅, E3 = Rat group fed with LD₅₀, V1 = Rat group fed with LD₁₀ + Vit C, V2 = Rat group fed with LD₂₅ + Vit C, V3 = Rat group fed with LD₅₀ + Vit C (Mean values with different subscripts differ significantly ($p \leq 0.05$)).

Parameter	Control Group	E1	E2	E3	V1	V2	V3
WBC (10^9 / L)	8.62 \pm 0.67 ^a	7.28 \pm 1.06 ^b	6.91 \pm 0.71 ^b	4.62 \pm 0.94 ^c	7.82 \pm 1.01 ^b	7.07 \pm 0.77 ^b	5.08 \pm 1.04 ^c
LYM (10^9 / L)	6.93 \pm 0.01 ^a	7.31 \pm 0.95 ^a	8.08 \pm 1.24 ^a	8.97 \pm 0.59 ^b	7.02 \pm 1.46 ^a	7.84 \pm 0.80 ^a	8.33 \pm 0.21 ^a
MID (10^9 / L)	0.68 \pm 0.18 ^a	0.61 \pm 0.15 ^a	0.50 \pm 0.13 ^b	0.26 \pm 0.02 ^b	0.64 \pm 0.14 ^a	0.55 \pm 0.08 ^b	0.36 \pm 0.06 ^c
GRA (10^9 / L)	0.55 \pm 0.03 ^a	0.43 \pm 0.06 ^b	0.36 \pm 0.04 ^b	0.29 \pm 0.05 ^c	0.45 \pm 0.05 ^b	0.40 \pm 0.03 ^b	0.31 \pm 0.04 ^c

Table 2. Mean (\pm SD) values of RBCs and related parameters in different rat groups. E1 = Rat group fed with LD₁₀, E2 = Rat group fed with LD₂₅, E3 = Rat group fed with LD₅₀, V1 = Rat group fed with LD₁₀ + Vit C, V2 = Rat group fed with LD₂₅ + Vit C, V3 = Rat group fed with LD₅₀ + Vit C.

Parameter	Control Group	E1	E2	E3	V1	V2	V3
RBC (10^{12} / L)	7.63 \pm 0.74	7.02 \pm 0.69	6.34 \pm 0.64	5.18 \pm 0.64	7.40 \pm 0.38	6.87 \pm 0.33	6.13 \pm 0.41
HGB (g/dl)	14.70 \pm 0.68	13.20 \pm 1.07	12.50 \pm 0.72	10.70 \pm 0.45	13.40 \pm 0.38	12.87 \pm 1.24	11.21 \pm 1.09
HCT %	40.84 \pm 2.3	37.63 \pm 1.2	35.95 \pm 0.98	30.29 \pm 1.52	37.95 \pm 1.18	36.31 \pm 0.92	32.71 \pm 1.05
MCV (fl)	52.14 \pm 2.62	54.20 \pm 1.79	57.00 \pm 1.8	58.00 \pm 2.28	53.08 \pm 1.79	56.00 \pm 1.47	57.54 \pm 1.94
MCH (pg)	17.30 \pm 1.46	18.90 \pm 0.32	19.70 \pm 0.48	20.60 \pm 0.8	18.47 \pm 2.07	19.30 \pm 0.93	19.83 \pm 1.17
MCHC (g/dL)	34.13 \pm 1.93	34.90 \pm 3.60	34.98 \pm 0.58	35.20 \pm 0.71	34.77 \pm 2.63	34.92 \pm 2.17	35.13 \pm 0.7
RDWc %	21.55 \pm 2.06	20.50 \pm 1.06	17.27 \pm 0.57	15.30 \pm 1.22	20.64 \pm 1.73	17.80 \pm 1.11	16.67 \pm 2.77

Table 3. Mean (\pm SD) values of Platelets and related parameters in different rat groups. E1 = Rat group fed with LD₁₀, E2 = Rat group fed with LD₂₅, E3 = Rat group fed with LD₅₀, V1 = Rat group fed with LD₁₀ + Vit C, V2 = Rat group fed with LD₂₅ + Vit C, V3 = Rat group fed with LD₅₀ + Vit C.

Parameter	Control Group	E1	E2	E3	V1	V2	V3
PLT (10^9 / L)	554 \pm 5.7	543 \pm 4.24	525 \pm 4.74	459 \pm 3.16	549 \pm 3.74	534 \pm 4.74	507 \pm 5.96
PCT %	0.52 \pm 0.03	0.41 \pm 0.02	0.31 \pm 0.03	0.27 \pm 0.04	0.46 \pm 0.04	0.38 \pm 0.03	0.30 \pm 0.03
MPV (fl)	5.80 \pm 0.68	6.10 \pm 0.33	6.30 \pm 0.51	6.60 \pm 1.09	6.00 \pm 0.27	6.20 \pm 0.88	6.40 \pm 0.11
PDWc (10^9 / L)	33.20 \pm 1.29	33.80 \pm 0.8	34.30 \pm 1.06	35.20 \pm 0.79	33.53 \pm 1.93	34.00 \pm 2.41	34.50 \pm 0.84
PLCR %	9.03 \pm 0.44	10.56 \pm 0.84	17.31 \pm 1.13	21.17 \pm 0.7	9.66 \pm 0.81	12.87 \pm 1.24	20.58 \pm 0.96

Table 4. Mean (\pm SD) values of Renal Function Test (RFT) in different rat groups. E1 = Rat group fed with LD₁₀, E2 = Rat group fed with LD₂₅, E3 = Rat group fed with LD₅₀, V1 = Rat group fed with LD₁₀ + Vit C, V2 = Rat group fed with LD₂₅ + Vit C, V3 = Rat group fed with LD₅₀ + Vit C.

Parameter	Control Group	E1	E2	E3	V1	V2	V3
Serum Creatinine (mg/dL)	0.34 \pm 0.03	0.47 \pm 0.05	0.84 \pm 0.04	1.11 \pm 0.03	0.41 \pm 0.03	0.77 \pm 0.04	0.98 \pm 0.04
BUN (mg/dL)	15.4 \pm 1.14	20.0 \pm 1.86	22.6 \pm 2.70	28 \pm 2.74	17.2 \pm 1.48	21 \pm 2.25	26 \pm 3.74

Table 5. Weight / Length Index (mg/mm) of left tibia of different rat groups. E1 = Rat group fed with LD₁₀, E2 = Rat group fed with LD₂₅, E3 = Rat group fed with LD₅₀, V1 = Rat group fed with LD₁₀ + Vit C, V2 = Rat group fed with LD₂₅ + Vit C, V3 = Rat group fed with LD₅₀ + Vit C.

Parameter	Control Group	E1	E2	E3	V1	V2	V3
W/L Index (mg/mm)	20.63	18.75	17.36	16.83	19.14	17.93	17.13

Table 6. Robusticity Index (mm/mg^{1/3}) of left tibia of different rat groups. E1 = Rat group fed with LD₁₀, E2 = Rat group fed with LD₂₅, E3 = Rat group fed with LD₅₀, V1 = Rat group fed with LD₁₀ + Vit C, V2 = Rat group fed with LD₂₅ + Vit C, V3 = Rat group fed with LD₅₀ + Vit C.

Parameter	Control Group	E1	E2	E3	V1	V2	V3
Robusticity Index (mm/mg ^{1/3})	2.229	2.272	2.279	2.295	2.263	2.283	2.285

the control group. However, mean values of platelet volume (MPV), platelet distribution width (PDWc), and platelet large cell ratio percentage (P-LCR%) were not significantly increased in the experimental groups (Table 3). Vitamin C exhibited a non-significant impact. Toxic substances such as insecticides can have a negative impact on bone marrow cells, leading to impaired cell function, aplasia, and dysplasia [19]. The demeaning platelets may be attributed to the effect of the insecticide on bone marrow cells, affecting their formation or development and subsequently slowing down or halting hematopoiesis [8]. An elevated mean platelet volume (MPV) indicates an enhanced tendency

for blood clotting, which may increase the risk of thrombosis, stroke, or other cardiovascular anomalies [11]. Previous studies have reported a decreased platelet number due to the toxicity of diazinon, imidacloprid, and other insecticides [17, 20].

Liver Toxicity

Liver tissue histopathology, initially preserved, was carried out. The serum biomarkers for the Liver Function Test (LFT) were also performed. These two procedures helped us determine the liver's health after acute exposure of varying degrees. It was found that

White blood cell & related




Fig. 1. Comparative analysis of WBCs and its types in different rat groups. E1 = Rat group fed with LD₁₀, E2 = Rat group fed with LD₂₅, E3 = Rat group fed with LD₅₀, V1 = Rat group fed with LD₁₀ + Vit C, V2 = Rat group fed with LD₂₅ + Vit C, V3 = Rat group fed with LD₅₀ + Vit C.

RBC & related parameters




Fig. 2. Comparative analysis of RBCs and related parameters in different rat groups. E1 = Rat group fed with LD₁₀, E2 = Rat group fed with LD₂₅, E3 = Rat group fed with LD₅₀, V1 = Rat group fed with LD₁₀ + Vit C, V2 = Rat group fed with LD₂₅ + Vit C, V3 = Rat group fed with LD₅₀ + Vit C.

Platelets & related parameters




Fig. 3. Comparative analysis of Platelets and related parameters in different rat groups. E1 = Rat group fed with LD₁₀, E2 = Rat group fed with LD₂₅, E3 = Rat group fed with LD₅₀, V1 = Rat group fed with LD₁₀ + Vit C, V2 = Rat group fed with LD₂₅ + Vit C, V3 = Rat group fed with LD₅₀ + Vit C.

Biomarkers of LFT & KFT




Fig. 4. Comparative analysis of Liver Function Test (LFT) and Renal Function Test (RFT) in different rat groups. E1 = Rat group fed with LD₁₀, E2 = Rat group fed with LD₂₅, E3 = Rat group fed with LD₅₀, V1 = Rat group fed with LD₁₀ + Vit C, V2 = Rat group fed with LD₂₅ + Vit C, V3 = Rat group fed with LD₅₀ + Vit C.

total bilirubin, alkaline phosphatase (ALP), alanine transaminase (ALT), and aspartate transaminase (AST) increased significantly in the experimental group in comparison to the control group. Vitamin C exhibited a significant ameliorating effect on the mentioned liver biomarkers (Table 4).

There is a fair argument that these mounted up levels of liver enzymes could be the result of the insecticide's interference in the biosynthesis pathways of these enzymes, or any alterations in membrane permeability could be the potential cause of hepatic tissue degeneration. These abnormal findings demonstrate tissue damage and probable narcosis of hepatic tissues. These findings were supported by various [21, 22] studies of hepatic toxicity with different molecules like chlorpyrifos, imidacloprid, and fipronil. It is worth mentioning that most of the studies were carried out with chronic exposure rather than acute exposure, hence

further studies are required to examine the effects of acute exposure [6, 23-25].

Histopathological examination of liver tissues revealed severe hydropic degeneration and coagulative necrosis in Experiment Group E3 (LD₅₀), as well as mild congestion and coagulative necrosis in Group V (Fig. 5).


Renal and Cardiac Toxicity

The Renal Function Test (RFT) and kidney histopathology test were performed to assess the extent of renal damage. Table 5 shows that serum creatinine and Blood Urea Nitrogen (BUN) values were significantly increased in intoxicated rat groups compared to the control group. Vitamin C also demonstrated antioxidant and protective effects on these kidney-related serum parameters. Histopathological examination of renal tissues revealed congestion peri-tubular in nature, pyknosis, karyolysis, and coagulative necrosis in exposed groups, while mild congestion and coagulative necrosis have also been observed in the ascorbic acid supplemented groups (Fig. 6).

Histopathology of cardiac tissues showed loss of striated muscles and the fragmentation of cardiac cells in the exposed groups in a dose-dependent manner, whereas ascorbic acid supplementation exhibited mild congestion and cellular degeneration (Fig. 7). These results indicate the nephrotoxic potential of Dinotefuran. Hence, exposure-related alterations in different biomarkers associated with histopathology can be attributed to oxidative damage and the subsequent demeaning of the glomerular filtration rate, as reported in other studies by [26].


The tibia of the rat was used to assess the damage. The mean weight and length decreased in varyingly exposed groups as compared to the control group. Consequently, it led to a low Weight/Length Index and Robusticity Index, as indicated in Table No. 6. These findings highlight the inhibitory and protective effects of Vitamin C against the toxic effects of Dinotefuran. Overall, these results elucidate the reduced bone density [27]. It is important to note that there is currently no available data in regard to the weight/length index and robusticity index of any rat bone in relation to insecticide or pesticide toxicity.

Based on the aforementioned discussion, it is to be concluded that acute exposure to dinotefuran has an impact on the structure and functioning of vital organs in mammals, including the kidney, liver, heart, and bone [28]. It leads to alterations in hematological, biochemical, and only histopathological and bone-related parameters. Even acute exposure to Dinotefuran can be hazardous in a dose-dependent manner. Vitamin C supplementation shows potential in mitigating the toxicity of the insecticide, although the effects are dose-dependent.




A = Normal Liver of Control (40X) B = Liver of Experiment Group (40X) C = Liver of Vitamin Group (40X)

Fig. 5. Liver Histopathology of different rat groups.



D = Normal Kidney of Control (10X) E = Kidney of Experiment Group (40X) F = Kidney of Vitamin Group (40X)

Fig 6. Renal Histopathology of different rat groups.



F = Normal Heart of Control (10X) G = Heart of Experiment Group (40X) H = Heart of Vitamin Group (40X)

Fig. 7. Histopathology of heart of different rat groups.

Conclusions

The above mentioned study concluded that one-time exposure to Dinotefuran can damage mammalian cells in a dose-dependent manner. A strict regulatory policy must be in place when dispensing these insecticides. A regular intake of vitamin C, both by humans and livestock, can minimize damage. There is a need to explore acute exposure studies further.

Conflict of Interest

On the behalf of all the authors, I declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data Availability

Raw data is available on request.

Consent to participate

All the authors gave full consent to participate.

Consent to Publish

All the authors gave consent to publish this data.

Funding

Only departmental sources used.

Ethical Clearance

The research got approval from independent ethical committee of university.

References

- HOUCCHAT J.N., CARTEREAU A., LE MAUFF A., TAILLEBOIS E., THANY S.H. An Overview on the Effect of Neonicotinoid Insecticides on Mammalian Cholinergic Functions through the Activation of Neuronal Nicotinic Acetylcholine Receptors. *International journal of environmental research and public health*, **17** (9), 3222, **2020**.
- XU X., WANG X., YANG Y., ARES I., MARTÍNEZ M., LOPEZ-TORRES B., MARTÍNEZ-LARRAÑAGA M.R., WANG X., ANADÓN A., MARTINEZ M.A. Neonicotinoids: mechanisms of systemic toxicity based on oxidative stress-mitochondrial damage. *Archives of toxicology*, **96** (6), 1493, **2022**.
- ARSLAN P., GÜNAL A.Ç. Does Fipronil Effect on Aquatic Organisms? Physiological, Biochemical, and Histopathological Alterations of Non-Target Freshwater Mussel Species. *Water*, **15** (2), 334, **2023**.
- DINCA V., DOCEA A.O., DROCAS A.I., NIKOLOUZAKIS T.K., STIVAKTAKIS P.D., NIKITOVIC D., GOLOKHAVAST K.S., HERNANDEZ A.F., CALINA D., TSATSAKIS A. A mixture of 13 pesticides, contaminants, and food additives below individual NOAELs produces histopathological and organ weight changes in rats. *Archives of toxicology*, **97** (5), 1285, **2023**.
- KHAN I., ZANEB H., MASOOD S., ASHRAF S., REHMAN H.F., TAHIR S.K., REHMAN H.U., KHAN A., TAJ R., RAHMAN S.U., SHAH M. Supplementation of Selenium Nanoparticles-Loaded Chitosan Improves Production Performance, Intestinal Morphology, and Gut Microflora in Broiler Chickens. *The journal of poultry science*, **59** (3), 272, **2022**.
- VOHRA P., KHERA K.S., SANGHA G.K. Physiological, biochemical and histological alterations induced by administration of imidacloprid in female albino rats. *Pesticide biochemistry and physiology*, **110**, 50, **2014**.
- SEEDOR J.G., QUARTUCCIO H.A., THOMPSON D.D. The bisphosphonate alendronate (MK-217) inhibits bone loss due to ovariectomy in rats. *Journal of Bone and Mineral Research*, **6** (4), 339, **1991**.
- NOREEN S., KHAN I.M., KHAN M.S., ZARNAAB B., GUL I., KHAN M.Z., JADOON W.A., GHAYYUR S., LIU Y. Comparative valuation of the chlorpyrifos, acetamiprid, and lambda-cyhalothrin toxicity and their hematological and histopathological consequences in pigeons. *Environmental science and pollution research international*, **30** (40), 92817, **2023**.
- ALIZADEH S.A., ANANI-SARAB G., AMIRI H., HASHEMI M. Paraquat induced oxidative stress, DNA damage, and cytotoxicity in lymphocytes. *Heliyon*, **8** (7), e09895, **2022**.
- GHAZI S., HABIBIAN M., MOEINI M., ABDOLMOHAMMADI A. Effects of different levels of organic and inorganic chromium on growth performance and immunocompetence of broilers under heat stress. *Biological trace element research*, **146**, 309, **2012**.
- MUNDY-HEISZ K.A., PROSSER R.S., RAINE N.E. Acute oral toxicity and risks of four classes of systemic insecticide to the Common Eastern Bumblebee (*Bombus impatiens*). *Chemosphere*, **295**, 133771, **2022**.
- HATHOUT H., M.SOBHY H., ABOU-GHANIMA S., EL-GARAWANI I.M. Ameliorative role of ascorbic acid on the oxidative stress and genotoxicity induced by acetamiprid in Nile tilapia (*Oreochromis niloticus*). *Environmental Science and Pollution Research*, **28**, 55089, **2021**.
- CHATTERJEE S., BASAK P., CHAKLADER M., DAS P., PEREIRA J.A., CHAUDHURI S., LAW S. Pesticide induced alterations in marrow physiology and depletion of stem and stromal progenitor population: an experimental model to study the toxic effects of pesticide. *Environmental toxicology*, **29** (1), 84, **2014**.
- KATARIA S.K., CHHILLAR A.K., KUMAR A., TOMAR M., MALIK V. Cytogenetic and hematological alterations induced by acute oral exposure of imidacloprid in female mice. *Drug and chemical toxicology*, **39** (1), 59, **2016**.
- JASPER R., LOCATELLI G.O., PILATI C., LOCATELLI C. Evaluation of biochemical, hematological and oxidative parameters in mice exposed to the herbicide glyphosate-Roundup®. *Interdisciplinary toxicology*, **5** (3), 133, **2012**.
- SUBRAMANIAN M. *Toxicology: Principles and Methods*. MJM Publisher, **2019**.
- AHMAD L., GUL T.S., DU X., HUSSAIN R., KHANANI M.R., IRAM S., REHMAN A., KHAN A. Dose- and dosage-dependent spectrum of respiratory toxicity of cypermethrin in rabbits. *Toxin Reviews*, **42** (3), 573, **2023**.
- MUNDY-HEISZ K.A., PROSSER R.S., RAINE N.E. Acute oral toxicity and risks of four classes of systemic insecticide to the Common Eastern Bumblebee (*Bombus impatiens*). *Chemosphere*, **295**, 133771, **2022**.
- ZEINALI M., MEYBODI N.T., REZAEI S.A., RAFATPANAH H., HOSSEINZADEH H. Protective effects of chrysin on sub-acute diazinon-induced biochemical, hematological, histopathological alterations, and genotoxicity indices in male BALB/c mice. *Drug and chemical toxicology*, **41** (3), 270, **2018**.
- KATARIA S.K., CHHILLAR A.K., KUMAR A., TOMAR M., MALIK V. Cytogenetic and hematological alterations induced by acute oral exposure of imidacloprid in female mice. *Drug and chemical toxicology*, **39** (1), 59, **2016**.
- FENG X., LIANG R., SHI D., WANG D., XU T., CHEN W. Urinary acrolein metabolites, systemic inflammation, and blood lipids: Results from the National Health and Nutrition Examination Survey. *Chemosphere*, **286** (Pt 2), 131791, **2022**.
- HOSSAIN M.A., SUTRADHAR L., SARKER T.R., SAHA S., IQBAL M.M. Toxic effects of chlorpyrifos on the growth, hematology, and different organs histopathology of Nile tilapia, *Oreochromis niloticus*. *Saudi journal of biological sciences*, **29** (7), 103316, **2022**.
- CHAKROUN S., EZZI L., GRISSA I., KERKENI E., NEFFATI F., BHOURI R., SALLEM A., NAJJAR M.F., HASSINE M., MEHDI M. Hematological, biochemical, and toxicopathic effects of subchronic acetamiprid toxicity in Wistar rats. *Environmental Science and Pollution Research*, **23**, 25191, **2016**.
- SUN Y.J., LIANG Y.J., YANG L., LONG D.X., WANG H.P., WU Y.J. Long-term low-dose exposure of permethrin induces liver and kidney damage in rats. *BMC pharmacology & toxicology*, **23** (1), 46, **2022**.

25. MARQUES L.P., JOVIANO-SANTOS J.V., SOUZA D.S., SANTOS-MIRANDA A., ROMAN-CAMPOS D. Cardiotoxicity of pyrethroids: molecular mechanisms and therapeutic options for acute and long-term toxicity. *Biochemical Society transactions*, **50** (6), 1737, **2022**.
26. BDEL-DAIM M.M., ABDEEN A. Protective effects of rosuvastatin and vitamin E against fipronil-mediated oxidative damage and apoptosis in rat liver and kidney. *Food and chemical toxicology*, **114**, 69, **2018**.
27. BABU E., GOPALAKRISHNAN V., SRIGANTH I.N. P., GOPALAKRISHNAN R., SAKTHISEKARAN D. Cisplatin induced nephrotoxicity and the modulating effect of glutathione ester. *Molecular and cellular biochemistry*, **144**, 7, **1995**.
28. MONTEAGUDO M., HERNANDEZ E., SECO C., GONZALEZ-RIOLA J., REVILLA M., VILLA L., RICO H. Comparison of the bone robusticity index and bone weight/bone length index with the results of bone densitometry and bone histomorphometry in experimental studies. *Cells Tissues Organs*, **160** (3), 195, **1997**.