

Original Research

Effects of Nano-Potassium Molybdate Preventing the Heavy Metal Poisoning in the Przewalski's Gazelle

Wanyi Zeng^{1#}, Hong Ren^{2#}, Xiaoyun Shen^{1*}¹School of Life Science and Engineering, Southwest University of Science and Technology, Mianyang, China²North Sichuan Medical College, Nanchong 637100, China*Received: 14 November 2023**Accepted: 25 March 2024*

Abstract

In the past 10 years, the level of heavy metals has sharply increased in the Jiangxigou farm in the Qinghai Lake basin, which has seriously affected the health of Przewalski's gazelle. Nano-potassium molybdate (nano-K₂MoO₄), as a new nano-fertilizer, has been applied in agriculture. To study the effect of applying different levels of nano-K₂MoO₄ fertilization (8 kg/hm², group I; 9 kg/hm², group II; 10 kg/hm², group III) on preventing heavy metal toxicity in Przewalski's gazelle. The samples of soil, forage, and animal tissues were collected for testing heavy metals and mineral contents, oxidative stress, and blood indexes. The empirical findings showed that the contents of selenium (Se) and copper (Cu) in the tested forage were significantly higher than those in the control group (p<0.01). The levels of Cu and lead (Pb) in the blood and liver of the tested Przewalski's gazelles were remarkably higher than those in the control animals (p < 0.01), but the levels of molybdenum (Mo) and Pb were remarkably lower than those in the control animals (p < 0.01). The levels of hemoglobin (Hb), platelets (PLT), and red blood cells (RBC) in the tested Przewalski's gazelles were remarkably higher than those in the control animals (p<0.01). The enzymatic activities of glutathione peroxidase (GSH-Px), superoxide dismutase (SOD), catalase (CAT), and total antioxidant capacity (T-AOC) in the serum of the tested Przewalski's gazelles exhibited a significant increase in comparison to the control animals (p<0.01), but the content of malondialdehyde (MDA) was lower than that in the control animals (p<0.01). In summary, the application of nano-K₂MoO₄ not only significantly improved the antioxidant capacity and alleviated effectively the anemia symptoms of Przewalski's gazelle, but also reduced the heavy-metal toxicity of Przewalski's gazelle in the fertilization areas.

Keywords: Przewalski's gazelle, nano-potassium molybdate, antioxidant capacity, toxicity of heavy metal, anaemia

[#]These authors contributed equally to this work and should be considered co-first authors.

*e-mail: xyshen@swust.edu.cn

Table 2. The physical and chemical properties in the soil from tested and healthy districts.

Items	Polluted district	Healthy district
OM (g/kg)	36.93±2.62	37.96±3.31
TS (mg/kg)	5577.77±43.67	5317.00±48.83

OM: organic matter; TS: total salt.

Sample Analysis

The samples were tested for mineral contents on 25 June 2020. Hematological and biochemical analyses were performed on 29 June 2020 (Table 4).

Statistical Analyses

These data were shown as mean±standard deviation. The data were performed using the Statistical Package for the Social Sciences analyzed (SPSS, version 23.0, Inc., IL, USA). The differences between the two groups were analyzed by applying the T- test. The very extreme difference was indicated by ** $p < 0.01$.

Results and Discussion

Detection of Nutritional Value in Forages

The values of CP and EE in the forage from fertilized areas were remarkably higher than those from the control group ($p < 0.01$, Table 5). In contrast, the value of N in the forage from fertilized areas was remarkably higher ($p < 0.01$) than that from the control areas. The value of S in the forage from fertilized areas was remarkably lower ($p < 0.01$) than that from the control group. No significant difference in the other indicators was found.

Detection of Heavy Metal Content in Forages

Compared to the control group, the levels of Se and Cu in the forage from the fertilized areas were remarkably higher ($p < 0.01$, Table 6). On the other hand, the levels of Mo and Pb in the fertilized forages were significantly lower than those in the control forages. There were no major differences observed in the levels of other elements.

Table 3. Samples of collection method.

Samples	Collection method
Soil samples	The samples of soil were gleaned from surface layer in randomly distributed locations in each farm. The soils were dried at 20-25°C until analysis [20].
Forage samples	The samples of forages were gleaned by using a mower, dried in a forced-air oven at 80°C, and ground to pass a 0.5-mm screen [4].
Tissue samples	The blood samples were collected from the jugular vein by vacuum blood collection tubes with EDTA-K ₂ [21]. The samples of blood were stored at 4°C until analysis. The serum samples were separated by centrifuge of 3 000 g for 15 min, and were stored at -20°C until analysis. Liver samples collections were performed by a trained technician, and stored at -20°C for analysis.

Table 4. The analysis of methods in the samples.

Indicators	Determination methods
Heavy metals	The analysis of heavy metals, including Cu, Zn, Cd, Pb, Hg, Se, and chromium, using an AA-7000 absorption spectrophotometer (Shimadzu Corporation, Japan) [3]. Mo was analyzed by using atomic absorption spectrophotometer (Perkin-Elmer 3030 graphite furnace with a Zeeman background correction).
Soil properties	The organic matter (OM) in the soil was analyzed by potassium dichromate sulfuric acid oxidation titration, and the TS in the soil was analyzed by drying residue mass method, and the water-soil ratio was 5:1 [5]. The pH value in the soil solution (water-soil ratio, 5:1) was analyzed with potentiometric method (PHS-3C, Shanghai Precision Scientific Instrument Co., Ltd) [22].
Physiology index	The blood indexes, including hemoglobin (Hb), red blood cell count (RBC), packed cell volume (PCV), mean corpuscular hemoglobin (MCH), mean corpuscular volume (MCV), mean corpuscular hemoglobin concentration (MCHC), white blood cell count (WBC) and platelet count (PLT), were analyzed by automatic blood cell analyzer (SF-3000, Sysmex-Toa Medical Electronics, Kobe, Japan) [23].
Nutrition values	Crude protein (CP) and crude fat (EE) of the forage were analyzed by kjeldahl method and Soxhlet extractor method, respectively. Crude fiber (CF) in the forage was analyzed by crude fiber analyzed apparatus (CXC-06, Wuhan Glemo Testing Equipment Co., Ltd).
Digestibility	The digestibility of the forage was analyzed with in vitro gas production technique [3]. Organic matter digestibility (OMD) and metabolic energy (ME) in the forage were calculated by gas production.

- atresia. *In Vitro Cellular and Developmental Biology – Animal*. **54** (6), 401, **2018**.
44. HUO B., WU T., CHI Y.K., MIN X.Y., SHEN X.Y. Effect of molybdenum fertilizer treatment to copper pollution meadow on copper metabolism in Wumeng semi-fine wool sheep. *Journal of Domestic Animal Ecology*. **40** (07), 44, **2019**.
 45. Song C.J., Shen X.Y. Effects of environmental zinc deficiency on antioxidant system function in Wumeng semi-fine wool sheep. *Biological Trace Element Research*. **195** (1), 110, **2020**.
 46. WU T., SONG M., SHEN X.Y. Seasonal dynamics of copper deficiency in Wumeng semi-fine wool sheep. *Biological Trace Element Research*. **197** (2), 487, **2020**.
 47. CHI Y.K., XIONG K.N., CHEN H., MIN X.Y., XIAO H., LIAO J.J., SHEN X.Y. Effect of Grazing to Copper Pollution Meadow on Copper Metabolism in Wumeng Semi-fine Wool Sheep. *Polish Journal Environmental Studies*. **28** (3), 1083, **2019**.
 48. SONG C.J., GAN S.Q., HE J., SHEN X.Y. Effects of nano-zinc on immune function in Qianbei-Pockmarked goats. *Biological Trace Element Research*. **199** (2), 578, **2020**.
 49. LOPEZ-ALONSO M., MIRANDA M. Copper Supplementation, A Challenge in Cattle. *Animals (Basel)*. **10** (10), 1890, **2020**.
 50. EDWARDS J.R., PROZIALECK W.C. Cadmium, diabetes and chronic kidney disease. *Toxicology and Applied Pharmacology*. **238** (3), 289, **2009**.
 51. CHUNG Y., DESIRAJU S., NAMACHIVAYAM K., GUZMAN P., HE L., MOHANKUMAR K. Hematological changes in neonatal mice with phlebotomy-induced anemia. *Pediatric Research*. **92** (6), 1575, **2022**.
 52. ASHOUR M., MABROUK M.M., AYOUB H.F., EL-FEKY M.M.M.M., ZAKI S.Z., HOSEINIFAR S.H., ROSSI W. Jr, VAN DOAN H., EL-HAROUN E., GODA A.M.A-S. Effect of dietary seaweed extract supplementation on growth, feed utilization, hematological indices, and non-specific immunity of Nile Tilapia, *Oreochromis niloticus* challenged with *Aeromonas hydrophila*. *Journal of Applied Phycology*. **32** (5), 3467, **2020**.
 53. EL BOUJI S., KAMIL N., BEIDOURI Z. Experimental study of an oscillating water column wave energy converter based on regular waves. *Marine Systems and Ocean Technology*. **17**, 147, **2023**.
 54. DASS S., SIDDIQUI J.A., MULAVEESALA R. Applicability of nanoparticle coating in bone density evaluation using gaussian-weighted linear frequency-modulated thermal wave imaging. *Russian Journal of Nondestructive Testing*. **59**, 228, **2023**.
 55. KUMAR B.S., SRIKANTH K. A study on properties of pervious concrete with high-volume usage of supplementary cementitious materials as substitutes for cement. *Asian Journal of Civil Engineering*. **24**, 1997, **2023**.
 56. MA Y.X., CHEN Q., ZHANG Y., XUE J.H., LIU Q.W., ZHAO Y.Y., YANG Y.P., HUANG Y., FANG W.F., HOU Z.G, LI S.R., WANG J., ZHANG L., ZHAO H.Y. Pharmacokinetics, safety, tolerability, and feasibility of apatinib in combination with gefitinib in stage III/IV EGFR-mutated non-squamous NSCLC: a drug-drug interaction study. *Cancer Chemotherapy and Pharmacology*. **92** (5), 411, **2023**.