Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy

Pol. J. Environ. Stud. Vol. 33, No. X (2024), 1-16
DOI: *10.15244/pjoes/186936* ONLINE PUBLICATION DATE:

Original Research

Species Diversity and Phytosociological Study of Herbaceous Layer of Rangeland Ecosystem

Muhammad Saleem[1], Faisal Hussain[1*], Muhammad Amjad Bashir[2], Yasser S. Mostafa[3], Saad A. Alamri[3], Suliman Mohammed Suliman Alghanem[4], Ibtisam Mohammed Alsudays[5], Fahad Mohammed Alzuaibr[6], Usama Ahmad Khan[2], Abdul Basit[2]

[1]Department of Botany, Faculty of Sciences, Ghazi University, Dera Ghazi Khan 32200-Pakistan
[2]Department of Plant Protection, Faculty of Agricultural Sciences, Ghazi University Dera Ghazi Khan 32200-Punjab, Pakistan
[3]Department of Biology, College of Science, King Khalid University, Abha P.O. Box 9004, Saudi Arabia
[4]Department of Biology, College of Science, Qassim University, Burydah 52571, Saudi Arabia
[5]Department of Biology, College of Science and Arts, Qassim University, Unaizah, Saudi Arabia
[6]Department of Biology, Faculty of Science, University of Tabuk, Tabuk-71491, Saudi Arabia

Received: 11 December 2023
Accepted: 8 April 2024

Abstract

The present study was proposed to work on species diversity and phytosociological study of the herbaceous layer at eight different sites in the surrounding area of Dera Ghazi Khan, particularly Choti Zareen, to determine a comprehensive phytosociological analysis, assessment of species diversity, and distribution of herbaceous species. The sampling data was obtained to analyze density, relative density, frequency, relative frequency, the IVI index, and to conduct the analysis of species diversity in herbaceous and dynamic communities. For this purpose, a total of 400 quadrats were selected for the evaluation of 8 different sites by identifying and analyzing various indices (Shannon Diversity-Index (H), species evenness, their variance, and Simpson index D). During the period of field surveys, 120 herbaceous species were analyzed, belonging to 92 genera, and distributed in 35 different families. Our phytosociological analysis revealed that *Chloris flagellifera* in site no. 2 with IVI 22.9, followed by *Aerva javanica* with IVI 13.24, were found to be the most dominant species as compared to others, respectively. The highest density was 38 plant/m^2 recorded in sampling site no. 4. The highest species richness was 30 at site no. 2. The highest values of the Shannon Diversity Index (H) and Simp Index (D) are also shown at site 2. The phytosociological analysis of the studied sites provides valuable insight

*e-mail: faisal.botanist2011@gmail.com
Tel.: +92-344-2743495.

Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy

to ecologists for the management and conservation of plant diversity, which is influenced by many climatic factors such as water, air, and rain, as well as anthropogenic stress.

Keywords: Species Diversity, herbaceous layer, Index, Density

Introduction

Phytosociology is an arch to describe the population dynamics, community structure, and diversity indices of plant communities [1]. A phytosociological study of the area gives a detailed explanation of the distribution pattern of species and their relationships between different plant communities [2]. The study of plant communities is the best way to learn about the habitat and vegetation structure of these communities [3]. Phytosociological methods often involve the quantitative understanding of various parameters like abundance, density, frequency, etc. [4]. The ecological properties of vegetation composition, like productivity, competition, niche structure, stability, and the relationship of plants in their community, are all determined by species diversity [5]. Such properties are often of critical ecological significance as they shed light on many interesting processes operating at the community level and provide an explanation of not only within-community patterns but also vegetation processes such as dynamics, stability, structuring, and functioning of communities and species diversity [6-9]. The species diversity of these communities is related to their structural habitat, which is supported by the internal and external function and potential of their habitat [10, 11].

Naturally, herbs serve as shelter and habitat for many animals and form the complex food chain and food webs [12-15]. Grasses provide shelter for a large number of microbial communities [16]. They also protect topsoil from soil erosion, improve water penetration into soils, and increase soil holding capacity [17-19]. Anthropogenic and multi-environmental factors affect species diversity, such as air, water, and other atmospheric factors [20-23]. Environmental factors, disturbance, dispersal ability and rate of plant species, and various ecological interactions determined the structure of the vegetation community in a complex ecosystem [24-26].

Humans, through overpopulation, exerted pressure on habitat destruction and reduced this influx [27, 28]. It was observed that agricultural land use in the study area affects plant composition and pattern [29, 30]. Making decisions for the conservation and management of floristic composition and structures of vegetation and biodiversity provides basic information to evaluate these results [10].

The aim of the present investigation was to analyze the phytosociological characteristics and diversity patterns of the herbaceous plants of the area, to measure the species diversity and its components for the phytosociological analysis, and to examine the environmental factors and anthropogenic pressure that were exerted on the herb layer of the study area as one of the main medicinal, economic, and fodder resources in Choti Zareen, Dera Ghazi Khan, Pakistan.

Material and Methods

Study Area

The study was conducted in Choti Zareen locality, 30 km south of the district of Dera Ghazi Khan, Pakistan, and 26 km east of the Koh-e-Suleiman range, extending from latitude 29°41`N to 29°52`N and from longitude 70°20` to 70°34`E (Fig. 1). The study area



Fig. 1. The map of the study area.

Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy

Table 1. Coordinates, altitude, and soil type of the study sites.

Site #	Location	Latitude (N)	Longitude (E)	Altitude (M)	Soil Type
1.	Basti Talpur	29˚43` N	70˚ 29` E	93 m	Sandy loam
2.	Mithawan	29˚46` N	70˚ 20` E	190 m	Sandy loam
3.	Chak Talpur	29˚47` N	70˚22` E	157 m	Loam
4.	Qaim Wala	29˚52` N	70˚ 29` E	116 m	Sandy loam
5.	Khanpur	29˚45` N	70˚ 34` E	86 m	Sandy loam
6.	Bakherwah	29˚41` N	70˚ 24` E	136 m	Silt loam
7.	Rohi Talpur	29˚ 45` N	70˚ 26` E	134 m	Silt loam
8.	Chak Jalloharr	29˚50` N	70˚ 24` E	145 m	Sandy loam

has a tropical monsoon and temperate climate, with high temperatures during the summer of about 50ºC, while during the winter the average temperature is 4ºC. The rainy season extends from July to August, with annual monsoon rainfall ranging from 87 to 165.83 mm. The field study was conducted from 2022 to 2023, covering both dry and rainy seasons. Eight vegetation sites were selected based on their physiognomy, topography, and altitude, representing vegetation variation (Table 1). The Choti Zareen supported plain, desert, fluvial, and wetland habitats, causing a large amount of plant diversity.

Vegetation Sampling and Data Collection

At each of the 8 sites, random quadrate sampling was used to measure the herbaceous diversity of the study area. 50 quadrates (1 × 1 m) in size were randomly placed for sampling in each site, and a total of 400 quadrates were taken from 8 different sites. For the valuation of the dominance of plant species in the vegetation community, the density, frequency, and abundance, and were converted into relative values and summed to obtain statistical data. The importance value index (IVI) of the study area was calculated by the methods of Dangoli & Shivakoti [31] and Chaudhry et al. [32]. Species richness and evenness were determined in the studied sites, and species diversity was measured using the Shannon diversity index (H) [33]. The Pielou index was used for the estimation of species evenness (E) [34].

The vegetation analysis of the studied communities was analyzed for the Importance Value Index, which was obtained by Density (D), Frequency, and their relative values.

$$\text{Density(D)} = \frac{\text{Number of individuals of species in all the sample plots}}{\text{Total number of sample plots studied}}$$

$$\text{Relative Density (R.D)} = \frac{\text{Number of individuals of species}}{\text{Total number of sample plots studied}} \times 100$$

$$\text{Frequency(F)} = \frac{\text{Number of points at which species occour}}{\text{Total number of points sampled}} \times 100$$

$$\text{Relative Frequency (R.F)} = \frac{\text{frequency of a specie}}{\text{Total frequencies of all species}} \times 100$$

$$\text{Importance Value Index} = \frac{\textbf{Relatve Density+Relative Frequency}}{\textbf{2}}$$

Diversity Indices

For the measurement of species diversity in herbaceous communities, commonly indices are usually used in ecology. The purpose of diversity indices was to measure species diversity and vegetation composition in plant communities [35]. For quantifying community diversity, indices provided an authentic and fruitful source, and they were useful in the analysis of the community structure of the plain and desert areas. Different types of diversity indices had been applied to contrast the plant species sampled from different sites. The general species diversity of the vegetation communities was measured by the popular Shannon–Wiener information theory function:

$$H = -\sum Pi \ln Pi \quad \text{I=1}\ldots\text{. S}$$

Where H is showing the Shannon diversity-index, which is general species diversity, Pi is the proportion of individuals in a community, ln is the natural log, taking the base e and expressed heterogeneity, and S denotes the total number present in the community [33, 36]. The variance of general diversity, Var (H), was calculated by Magurran [35], as follows:

$$\text{Var}(H) = \frac{\sum Pi\,(\ln Pi)\,2 - (\sum \text{Pi} \ln Pi)2}{N} + \frac{(S-1)}{2N\,2} \quad \text{i=1, S}$$

General diversity consists of two diversity components: species richness and evenness. Species richness is expressed as S; the function ratio of the total number of individuals (N); and equitability (J) measures the evenness of the allocation of individuals among the species [35]. The evenness and its variance, Var (J), were measured in accordance with Pielou [37].

Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy



Fig. 2. Top families according to the number of species present in the study area.

$$J = \frac{H}{Hmax} = \frac{H}{ln\ S}$$

The equitability (J) is the ratio between observed H and maximal diversity H max. The variance of equitability was estimated as:

$$Var\ (J) = \frac{(H)}{(ln\ S)2}$$

The Simpson Index determines the dominance of species. The formula for the Simpson index is:

$$D = \frac{\Sigma ni(ni-1)}{N(N-1)}$$

Where D = Simpson's Index, ni = the number of individuals of species I, and N = the total number of individuals.

Results and Discussion

A total of 120 species, representing 92 genera from 35 families, were recorded from the studied area. Poaceae was the dominant family with 28 species, followed by Asteraceae (14), Amaranthaceae (8), Euphorbiaceae (7), Boraginaceae, Cyperaceae, and Polygonaceae (5 species each), Brassicaceae, Convolvulaceae, Fabaceae, and Solanaceae (4 species each), Cucurbitaceae and Plantaginaceae (3 species

Table 2. Vegetation composition of the herbaceous layer of various sites (Frequency (F), Relative Density (R.D.), and Relative Frequency (R.F.).

Species	Density/m^2	F	R.D.	R.F.
Site 1: Basti Talpur				
Cynodon dactylon L.	2.8	24	6.024	7.5
Dactyloctenium aegyptium L.	2.24	20	4.819	6.25
Desmotachya bipinnata L.	2.64	20	5.680	6.25
Dichanthium annulatum Forssk.	2.88	16	6.196	5
Typha angustifolia L.	2	12	4.303	3.75
Site 2: Mithawan				
Chloris flagellifera Nees.	3.84	60	15.190	15.464
Aerva javanica Burm. f.	2.24	34	8.861	8.763
Eragrostis minor Host.	1.84	32	7.278	8.247
Fagonia cretica L.	1.2	20	4.747	5.155
Solanum virginianum L.	1.6	20	6.329	5.155

Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy

Table 2. Continued.

Site 3: Chak Talpur				
Aerva javanica Burm. f.	2.08	30	5.977	6.466
Chloris flagellifera Nees.	1.92	24	5.517	5.172
Fagonia cretica L.	1.76	24	5.057	5.172
Solanum virginianum L.	1.2	20	3.448	4.310
Glinus lotoides L.	1.44	20	4.138	4.310
Site 4: Qaim Wala				
Cenchrus ciliaris L.	2.32	32	6.713	7.442
Eragrostis minor Host.	1.68	18	4.861	4.186
Aerva javanica Burm. f.	0.96	18	2.778	4.186
Chloris flagellifera Nees.	1.12	18	3.241	4.186
Alternanthera sessilis L.	1.52	16	4.398	3.721
Site 5: Khanpur				
Chenopodiastrum murale L.	3.04	24	7.95	5.941
Typha angustifolia L.	1.44	22	3.766	5.446
Melilotus indicus L.	1.28	18	3.347	4.455
Fagonia cretica L.	1.84	18	4.812	4.455
Vicia sativa L.	1.76	16	4.603	3.960
Site 6: Bakherwah				
Fagonia cretica L.	2.32	24	5.906	5.505
Citrullus colocynthis L.	1.68	24	4.277	5.505
Eragrostis minor Host.	1.44	16	3.666	3.670
Euphorbia serpens Kunth.	0.96	16	2.444	3.670
Aerva javanica Burm. f.	1.12	16	2.851	3.670
Site 7: Rohi Talpur				
Chenopodiastrum murale L.	3.76	24	9.457	6.154
Cenchrus ciliaris L.	1.52	20	3.823	5.128
Fagonia cretica L.	1.84	18	4.628	4.615
Cyperus rotundus (L.)	1.2	14	3.018	3.590
Chrozophora plicata Vahl.	1.04	14	2.616	3.590
Site 8: Chak Jalloharr				
Aerva javanica Burm. f.	1.52	24	4.158	5.455
Convolvulus arvenses L.	2.8	24	7.659	5.455
Glinus lotoides L.	2	24	5.470	5.455
Convolvulus prostratus Forssk.	1.04	20	2.845	4.545
Solanum virginianum L.	0.88	18	2.407	4.091

each), Aizoaceae, Caryophyllaceae, Oxalidaceae, and Zygophyllaceae (2 species each), while the other 18 families were represented by only one species (Fig. 2).

For the vegetation composition, a total of 5 species have been selected from each site with the highest density and frequency. In the present study, density (density/m^2), relative density (%), frequency (%), and relative frequency (%) were calculated for all 120 species (Table 2). The result highlighted that the species *Chloris flagellifera* showed the highest density and relative









Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy

Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy

Fig. 3. Vegetation composition with species diversity=H, species evenness=J, variance of evenness=Var(J), variance of diversity=Var(H), species richness=d_1 and Dominance=D values of different sites (1= *Achyranthes aspera,* 2= *Aerva javanica,* 3= *Aeluropus lagopoides,* 4= *Alternanthera sessilis,* 5= *Amaranthus deflexus,* 6= *Anagallis arvensis,* 7= *Asphodelus tenuifolius,* 8= *Boerhavia diffusa,* 9= *Carthamus lanatus,* 10= *Cenchrus ciliaris,* 11= *Chenopodiastrum murale,* 12= *Cleome brachycarpa,* 13= *Chloris flagellifera,* 14= *Chrozophora plicata,* 15= *Citrullus colocynthis,* 16= *Convolvulus arvenses,* 17= *Convolvulus prostrates,* 18= *Cressa cretica,* 19= *Crisium arvense,* 20= *Cynodon dactylon,* 21= *Cyperus rotundus,* 22= *Dichanthium annulatum,* 23= *Echinops echinatus,* 24= *Eclipta alba,* 25= *Eragrostis minor,* 26= *Euphorbia helioscopia,* 27= *Euphorbia hirta,* 28= *Euphorbia serpens,* 29= *Fagonia cretica,* 30= *Fumaria parviflora,* 31= *Galium aparine,* 32= *Glinus lotoides,* 33= *Heliotropium europoeum,* 34= *Heliotropium ovalifolium,* 35= *Imperata cylindrica,* 36= *Launaea nudicaulis,* 37= *Malcolmia Africana,* 38= *Medicago polymorpha,* 39= *Melilotus indicus,* 40= *Oxalis dilleniid,* 41= *Parthenium hysterophorus,* 42= *Persicaria glabra,* 43= *Phalaris minor,* 44= *Phragmites australis,* 45= *Phyla nodiflora,* 46= *Polypogon monspeliensis,* 47= *Ranunculus sceleratus,* 48= *Reseda villossa,* 49= *Solanum americanum,* 50= *Solanum virginianum,* 51= *Sonchus oleraceus,* 52= *Sorghum halepense,* 53= *Spergularia marina,* 54= *Taraxacum officinale,* 55= *Tribulus terrestris,* 56= *Typha angustifolia,* 57= *Vicia sativa,* 58= *Zaleya pentandra*).

density, followed by *Chenopodiastrum murale* in sites 2, 7, and 5, respectively. Whereas frequency and relative frequency were found highest from *Chloris flagellifera* in site 2, followed by *Aerva javanica*, *Cenchrus ciliaris,* and *Eragrostis minor* as compared to other sites (Table 2).

There are 10 important species selected with diversity indices from each site concerning their diversity pattern and conservation status at different sites. The result showed that the highest Simpson Index (D) was observed in Khanpur site no. 5 by *Dichanthium annulatum* because the study was conducted from December to April, which is the best season for its growth and yield production (Fig. 3).

The lowest Simpson index (D) was shown by *Euphorbia hirta* in site no. 2, Mithawan. Unusually, the overall highest Simpson index (D) is shown by Basti Talpur site no. 1 and the lowest by Mithawan site no. 2, which is the fluvial site. Mithawan is facing large grazing pressure, and agricultural activity is increasing day by day. *Chloris flagellifera* in site 2 had the highest general diversity (H) among all the sites, followed by *Cenchrus ciliaris* in site 4 and *Aerva javanica* in site 8 with Shannon diversity-index 2.61 and 2.41, respectively. The highest values of the Shannon Diversity-Index (H) in these sites indicate greater species richness and diverse community structure. Conversely, *Sonchus oleraceus* had the lowest 0.41 Shannon Diversity index in site 3 as compared to other sites. Across all sites, we observed that *Aerva javanica* had consistent Shannon Diversity-Index (H) values, which indicated its suitable habitat. In comparison to all sites, 4 and 5 had an even distribution of plant species. This means these are least disturbed by human impact, and these sites can maintain their diversity pattern through many ecological processes (Fig. 3).

The clear superiority in terms of species richness and evenness had been taken by *Chloris flagellifera* in site 2, followed by *Cenchrus ciliaris* and *Aerva javanica* in sites 4 and 3, respectively. The highest values of species richness and evenness of these species are an indication of dominance and adaptation in their respective ecosystems. In terms of even distribution and stability, the communities of site 4, site 5, and site



Fig. 4. Shows the highest diversity indices of different sites in the study area.

Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy

6 had the highest stability in their species richness and evenness in comparison to the other 5 sites, which had fluctuations between high and low in their values of species richness and evenness. Our observations of the diversity indices of each site indicated that a large number of dominant species in the study area are invasive, and the fluctuation of values between high and low for some species is proof of this phenomenon. The invasive species pose a major threat to the diversity of the area. The other reason is that the pressure exerted on many plant species, mainly fodder plants that belong to the Poaceae family, is because people of the study area are farmers in professions that have grazing animals like goats, sheep, cows, etc. and we know these animals consume more fodder and exert pressure on the natural vegetation of the study area (Fig. 4).

Table 3. Importance Value Index (IVI) and random distribution pattern of recorded plant species from various sites (Ann.=Annual, Per.= Perennial, Fd. = Fodder, Med.= Medicinal).

Families	Life cycle	Economic importance	Site 1	Site 2	Site 3	Site 4	Site 5	Site 6	Site 7	Site 8
Aizoaceae										
Trianthema portulacastrum (L.)	Ann.	Fd.	2.28	_	_	_	_	_	_	_
Zaleya pentandra (L.)	Per.	Fd.	_	1.15	2.30	3.48	_	_	4.30	1.78
Amaranthaceae										
Achyranthes aspera (L.)	Per.	Med.	4.24	_	_	5.34	3.24	_	_	_
Aerva javanica (Burm.f.) Juss. & Schult.	Per.	Med.	3.53	13.24	9.45	5.58	3.16	4.69	4.21	6.89
Alternanthera sessilis (L.)	Ann.	Med.	2.20	_	_	5.92	_	2.14	_	2.88
Amaranthus albus (L.)	Ann.	Med.	_	_	_	_	_	_	2.78	_
Amaranthus deflexus (L.)	Ann.	Med.	3.71	_	_	_	_	_	3.18	_
Amaranthus viridis (L.)	Ann.	Med.	1.59	_	_	_	_	_	_	_
Chenopodiastrum murale (L.)	Ann.	Med.	1.51	1.78	4.20	_	10.92	7.31	12.53	4.42
Chenopodium album (L.)	Ann.	Med.	_	1.84	_	_	_	6.49	2.18	3.99
Asphodelaceae										
Asphodelus tenuifolius (Cav.)	Ann.	Med.	_	_	2.64	_	_	2.29	4.45	_
Asteraceae										
Carthamus lanatus (L.)	Ann.	Med.	_	_	_	_	_	_	3.18	_
Crisium arvense (L.)	Ann.	Med.	2.82	_	_	4.53	4.66	5.65	_	_
Echinops echinatus (Roxb.)	Ann.	Med.	_	2.36	_	4.53	2.84	1.07	2.58	3.33
Eclipta alba (L.)	Per.	Med.	2.20	_	_	4.53	_	_	_	2.43
Erigeron bonariensis (L.)	Ann.	Med.	_	1.15	_	_	_	_	_	2.00
Gnaphalium uliginosum (L.)	Ann.	Med.	_	_	_	_	_	2.78	_	_
Launaea capitata (Spreng.) Dandy	Ann.	Med.	2.71	_	_	_	_	_	_	_
Launaea nudicaulis (L.) Hook.f.	Ann.	Med.	_	_	_	_	2.84	1.48	3.50	1.56
Launea mucronata (Forssk.)	Ann.	Med.	_	1.72	_	_	_	_	_	_
Parthenium hysterophorus (L.)	Ann.	Med.	_	_	_	4.99	_	_	_	_
Reichardia picroides (L.) Roth	Per.	Med.	_	_	_	_	_	0.87	_	_
Sonchus oleraceus (L.)	Ann.	Med.	_	1.72	1.67	2.21	2.70		2.23	_
Taraxacum officinale (L.)	Per.	Med.	_	1.47	2.99	_	_	3.79	_	_
Xanthium strumarium (L.)	Ann.	Med.	1.85	_	_	1.39	_	_	_	_

Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy

Table 3. Continued.

Boraginaceae										
Heliotropium bacciferum (Forssk.)	Per.	Fd.	_	1.84	1.98	_	_	_	_	_
Heliotropium europoeum (L.)	Ann.	Fd.	_	6.49	4.94	5.11	_	3.21	5.91	4.86
Heliotropium ovalifolium (Forssk.)	Per.	Med.	1.85	_	1.75	_	_	4.23		_
Heliotropium ramosissimum (Lehm.) Sieber ex DC.	Per.	Med.	_	_	1.98	_	_	_	3.33	_
Heliotropium strigossum (Wild.)	Per.	Med.	_	1.15	2.41	_	_	_	_	_
Brassicaceae										
Lepidium didymum (L.)	Ann.	Fd.	_	_	_	_	_	_	2.23	_
Malcolmia africana (L.)	Ann.	Med.	_	_	_	_	_	_	2.84	_
Moricandia arvense (L.)	Ann.	Fd.	_	2.99	_	_	_	_	_	_
Sisymbrium irio (L.)	Ann.	Fd.	_	_	_	3.83	_	_	_	_
Capparidaceae										
Cleome brachycarpa (Vahl ex DC.)	Per.	Med.	_	_	2.53	_	_	_	2.58	3.77
Caryophyllaceae										
Spergularia marina (L.) Besser.	Ann.	Med.	_	_	_	4.41	1.29	_	_	_
Stellaria media (L.) Vill,Dist.	Ann.	Med.	1.77	_	_	_	_	_	_	1.77
Convolvulaceae										
Convolvulus arvenses (L.)	Per.	Med.	5.06	_	3.42	_	6.09	4.84	2.64	10.39
Convolvulus prostratus (Forssk.)	Per.	Med.	_	_	5.32	_	1.58	3.77	4.70	5.12
Cressa cretica (L.)	Per.	Med.	_	0.57	_	5.92	4.66	_	1.72	_
Ipomea lacunosa (L.)	Ann.	Med.	1.68	_	_	_		_	_	_
Cucurbitaceae										
Citrullus colocynthis (L.)	Per.	Med.	_	1.72	4.28	3.83	_	7.03	3.04	_
Cucumis maderaspatanus (L.)	Per.	Med.	1.77	_	_	_	_	_	_	_
Cucumis melo var.agrestis (Naudin.)	Ann.	Med.	3.10	_	_	_	_	_	1.12	_
Cyperaceae										
Cyperus difformis (L.)	Ann.	Med.	1.59	_	_	_	4.55	_	_	_
Cyperus rotundus (L.)	Per.	Med.	4.14	4.51	3.53	_	5.67	2.93	4.81	3.33
Eleocharis geniculata (L.) Roem. & Schult.	Ann.	Med.	_	_	_	_	3.29	_	_	_
Schoenoplectus lacustris (L.) Palla.	Per.	Med.	_	_	_	3.25	1.12	_	_	_
Scirpus maritimus (L.)	Per.	Med.	_	_	_	_	2.87	_	_	_
Equisetaceae										
Equisetum ramosissimum (Desf.)	Per.	Med.	1.57	_	_	_	_	_	_	_
Euphorbiaceae										
Chrozophora plicata (Vahl) A.Juss. ex Spreng.	Ann.	Med.	_	_	3.53	4.06	_	3.62	4.41	5.53
Euphorbia helioscopia (L.)	Ann.	Med.	_	_	_	_	3.75	_	_	_

Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy

Table 3. Continued.

Euphorbia hirta (L.)	Ann.	Med.	_	1.72	4.86	_	_	_	_	_
Euphorbia maculata (L.)	Ann.	Med.	_	5.86	_	_	_	_	_	_
Euphorbia prostrata (Aiton.)	Ann.	Med.	_	_	1.09	_	_	_	_	_
Euphorbia serpens (Kunth.)	Ann.	Med.	2.13	_	_	_	_	4.28	_	2.00
Phyllanthus maderaspatensis (L.)	Ann.	Med.	_	_	3.74	_	_	_	_	_
Fabaceae										
Medicago polymorpha (L.)	Ann.	Fd.	_	_	_	_	_	_	_	3.99
Melilotus indicus (L.) All.	Ann.	Fd.	_	_	_	_	5.58	4.86	3.70	3.98
Vicia sativa (L.)	Ann.	Med.	_	_	_	_	6.58	3.62	1.52	3.10
Juncaceae										
Juncus maritimus (Lam.)	Ann.	Med.	_	_	_	1.97	0.91	1.50	_	_
Leguminosae										
Lathyrus aphaca (L.)	Ann.	Med.	_	_	_	_	3.92	_	_	_
Molluginaceae										
Glinus lotoides (L.)	Per.	Med.	_	1.72	6.38	_	2.87	3.62	_	8.20
Nyctiginaceae										
Boerhavia diffusa (L.)	Ann.	Med.	4.87	_	_	_	_	_	_	3.32
Orobanchaceae										
Lindenbergia indica (L.)			_	1.72	_	_	_	_		1.33
Oxalidaceae										
Oxalis carniculata (L.)	Ann.	Med.	_	1.15	_	_	_	_	3.39	3.09
Oxalis dillenii (Jacq.)	Ann.	Med.	_	_	_	_	_	4.61	_	_
Papaveraceae										
Fumaria parviflora (Lam.)	Ann.	Med.	_	_	_	_	4.13	_	_	_
Papilionaceae										
Trifolium repens (L.)	Per.	Fd.	2.20	_	_	_	_	_	_	_
Trifolium resupinatum (L.)	Ann.	Fd.	_	_	_	_	_	_	_	4.42
Plantaginaceae										
Nanorrhinum ramosissimum (Wall.)	Ann.	Med.	_	2.36	_	_	_	_	_	_
Veronica polita (Fries.)	Ann.	Med.	_	_	_	_	_	_	_	2.00
Veronica undulata (Wall.)	Per.	Med.	_	_	_	_	_	_	_	2.65
Poaceae										
Aeluropus lagopoides (L.)	Per.	Fd.	_	_	_	4.87	2.38	3.36	_	_
Agrostis capillaris (L.)	Per.	Fd.	1.59	_	_	_	_	_	_	_
Arundo donax (L.)	Per.	Med.	2.28	_	_	_	_	_	_	_
Bromus diandrus (Roth.)	Ann.	Med.	_	_	_	_	_	3.39	2.64	_
Cenchrus ciliaris (L.)	Per.	Fd.	3.25	5.09	5.37	10.80	3.33	3.21	6.39	3.54
Cenchrus longispinus (Hack.)	Ann.	Fd.	_	_	4.17	_	_	2.11	3.24	_
Chloris virgata (Sw.)	Ann.	Fd.	_	_	_	_	_	1.07	_	_

Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy

Table 3. Continued.

Chloris flagellifera (Nees) P.M.Peterson	Per.	Fd.	_	22.92	7.93	5.81	2.91	3.21	3.61	3.34
Cynodon dactylon (L.) Pers.	Per.	Med.	10.51	_	3.85	_	2.00	1.71	1.57	_
Dactyloctenium aegyptium (L.)	Ann.	Fd.	8.66	1.47	3.51	_	_	_	_	_
Dactyloctenium scindicum (Boiss.)	Per.	Fd.	_	4.34	_	_	_	_	_	_
Desmotachya bipinnata (L.)	Per.	Med.	9.09	_	_	5.34	1.33	_	_	_
Dichanthium annulatum (Forssk.) Stapf)	Per.	Fd.	8.10	4.39	2.73	5.46	4.07	4.05	_	3.78
Digitaria adscendens (Kunth) Henr.	Ann.	Fd.	_	_	_	_	_	_	_	2.44
Echinochloa colona (L.)	Ann.	Fd.	1.85	_	5.52	_	_	_	_	_
Eragrostis minor (Host.)	Ann.	Fd.	1.51	11.40	3.62	6.62	_	5.50	4.10	_
Eragrostis tenella (L.) P.Beauv. ex Roem. & Schult.	Ann.	Fd.	_	6.35	_	_	_	_	_	1.77
Imperata cylindrica (L.)	Per.	Fd.	3.08	_	_	4.18	0.91	_	_	_
Pennisetum orientale (L.)	Per.	E	_	_	2.73	_	_	_	_	0.89
Pennisetum setaceum (Forssk.)	Per.	E	_	2.61	_	1.97	_	_	0.92	_
Phalaris minor (Retz.)	Ann.	Fd.	_	_	_	_		_	1.57	2.44
Phragmites australis (Cav.) Trin.Ex Steud.	Per.	Fd.	5.21	_	_	4.88	2.70	1.50	_	_
Polypogon monspeliensis (L.)	Ann.	Med.	_	_	_	_	4.17	_	1.72	_
Saccharum munja (Roxb.)	Per.	E	2.48	_	1.09	4.07	2.70	_	_	0.45
Setaria pumila (Poiret.) Roemer & Schultes	Ann.	Fd.	1.77	_	_	_	_	_	_	_
Sorghum halepense (L.)	Per.	Fd.	_	1.15	5.03	4.06	_	_	3.41	2.90
Sporobolus indicus (L.)	Per.	Med.	_	_	5.00	_	_	_	_	_
Typha angustifolia (L.)	Per.	E	5.90	_	_	5.46	6.49	2.78	_	_
Polygonaceae										
Persicaria glabra (Willd.) M. Gómez	Per.	Med.	5.02	_	_	_	2.84	_	_	_
Polygonum aviculare (L.)	Ann.	Med.	_	_	_	_	_	_	_	4.20
Rumex acetosa (L.)	Per.	Med.	_	_	_	_	4.34	_	_	_
Rumex nepalensis (Spreng.)	Per.	Fd.	1.59	_	_	_	_	_	_	_
Rumex pulcher (L.)	Per.	Med.	_	_	_	_	1.09	_	_	_
Pontederiaceae										
Eichhornia crassipes (Mart.) solms.	Per.	Med.	2.46	_	_	_	_	_	_	_
Portulacaceae										
Portulaca oleracea (L.)	Ann.	Med.	_	_	_	1.86	_	_	_	_
Primulaceae										
Anagallis arvensis (L.)	Ann.	Med.	_	_	_	_	5.04	3.36	3.84	_
Ranunculaceae										
Ranunculus sceleratus (L.)	Ann.	Med.	_	_	_	_	_	3.18	_	_

Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy

Table 3. Continued.

Resedaceae										
Reseda villossa (Coss.)	Per.	Fd.	_	2.30	5.32	_	2.66	_	_	5.74
Rubiaceae										
Galium aparine (L.)	Ann.	Med.	_	_	2.64	_	_	_	_	_
Solanaceae										
Physalis minima (L.)	Ann.	Med.	1.68	_	_	_	_	_	_	_
Solanum americanum (Mill.)	Ann.	Med.	_	_	_	_	_	_	3.81	_
Solanum nigrum (L.)	Ann.	Med.	5.21	_	_	_	_	2.98	3.70	_
Solanum virginianum (L.)	Per.	Med.	4.24	8.91	6.03	_	2.70	2.14	3.04	4.45
Tiliaceae										
Chorchorus depressus (L.)	Per.	Med.	_	0.57	_	_	_	_	_	_
Verbenaceae										
Phyla nodiflora (L.) Greene	Per.	Med.	_	_	_	5.11	3.08	2.50	_	2.67
Zygophyllaceae										
Fagonia cretica (L.)	Per.	Med.	_	7.32	7.70	_	7.04	8.66	6.94	6.62
Tribulus terrestris (L.)	Ann.	Med.	2.99	7.76	2.76	_	_	4.64	3.70	_

The IVI values determined the dominance of species assigned on the basis of these IVI values. *Chloris flagellifera* was the dominant species, followed by *Aerva javanica*, *Cleome brachycarpa,* and *Chenopodiastrum murale*. The dominance of species indicated the fulfillment of suitable and desired conditions for ecological success in different studied sites. The random pattern (the distribution pattern) of plant species in different habitats was assessed, and the result revealed that 100% were randomly distributed (Table 3).

Dera Ghazi Khan Rangelands and their environments are home to a diverse array of plant species that flourish in a variety of habitats. There are two types of habitats in the research area: piedmont and alluvial. Dera Ghazi Khan is located in the Suleman mountain range and is extremely rich in floral variety. We identified the quantity and composition of communities in two historical locations, Piedmont and Riparian habitats, at various altitudinal inclines. The most significant source of habitat for floral variation is topographically diversified landscapes. Species diversity is also dramatically reduced at high altitudes compared to low altitudes [38]. Kenar & Kikvidze [39] reported that biological and evolutionary processes resulted in high species richness and served as corridors of plant and animal migration in the past. Site 1 had the highest density and relative density, and Site 4 had the lowest. Site 2 also had the highest frequency and relative frequency, and Site 6 had the lowest. Site 2 had the highest diversity indices value, and site 1 had the lowest. Poaceae has the greatest species diversity in all study sites due to the production of a large number of reproductive seeds and the many means of seed dispersion through grazing animals, fluvial water, and air. The magnitude of the diversity indices varied somewhat for different species, showing a low or high degree of aggregation where it was significant. Similar results were reported by Wang et al. [40]. Ahmad et al. [38] reported that a total of 76 species were predicted during the summer season. The maximum number of species belong to the Poaceae family, including *Aristida adscensionis, Brachiari areptans, Cenchrus ciliaris, Cenchchrus setiger, Chloris barbata, Cymbopogon jawarancusa, Cynodon dactylon, Cynodon glabratus, Demostachya bipinnata, Dichanthium annulatum, Enneapogon persicus, Eragrostis minor, Eragrostis barrelieri, Panicum hemitomon, Panicum repens, Phalaris minor, Phragmites australis, Saccharum munja, Saccharum officinarum,* and *Saccharum spontaneum*, and these are followed by the Fabaceae and Amaranthaceae families from Dera Ghazi Khan.

Plant scientists worldwide show a common interest in native plant species to predict new inclusive research investigations and ethnopharmacological, ecological, and phytoplankton studies skewed toward charismatic contributions [41, 42]. The phenotypic variation and systematic importance of flowering plants are investigated in different geographical regions of Pakistan to measure the functional diversity in restored ecosystems [43, 44]. As a result, the sources of floral scent variation and floristic diversity provide phylogenetic patterns as per the investigation of earlier researchers [45, 46]. According to Sharma et al. [47], the high species richness in the temperate ecosystem is an indication of better environmental conditions for the survival of plant communities, such as moderate

Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy

temperature, proper seasonal rainfall, and moisture content that are desired for healthy plant growth. Evenness accounts for more variation in the Shannon diversity index (H), which is in agreement with Yadav & Sharma [48]. The highest species richness was also shown by *Chloris flagellifera* in Site 2, with an altitude of 190m. With respect to sampling, it must be noted that even in the case of random sampling, the quadrats may not be independent samples as the variation in species richness in the field may influence the quadrats (that may be species poor or species rich accordingly) situated in the vicinity of each other in species poor or species rich sections [49, 50]. Variances of diversity and equitability were not unexpectedly consistently low for almost all species, as has previously been observed by Ivashchenko et al. [51]. Species diversity may be important because of its possible role in the establishment and coexistence of species though, in some model systems it is found to play hardly any role in these processes [52]. Dominance concentration (D) was found to vary inversely with general diversity (H), which is in agreement with the results of Wang et al. [40], Ivashchenko et al. [51], Hussain et al. [53], Bashir et al. [54], and Wang et al. [69].

Variables influencing species diversity include not just abiotic and environmental characteristics, but also climatic gradients impacted by biotic factors [55]. Sustainable use and conservation of plant biodiversity are important for geo-climatic gradients to drive plant species composition, which plays an important role in how plant species assemble locally into communities [56, 57]. An integrated framework of plant structure and function is dependent on the amount of above-ground vegetation [58], which is more constant owing to some psychological processes, and this is fully dependent on regional climatic and edaphic conditions. Our findings are also consistent with previous research [59] undertaken by several plant ecologists across Pakistan's diverse geographical zones. Herbaceous plants (55.26%) were the most prevalent living form in the alluvial environment of the research region.

The difference in IVI values of different species is due to their environmental conditions and anthropogenic activities around the sites. The wetland habitat is dominated by *Typha angustifolia*, *Schoenoplectus lacustris, Juncus maritimus,* and *Phragmites australis.* In addition, increasing anthropogenic pressures make Choti Zareen biodiversity a fairly sensitive and threatened ecosystem [60, 61]. Fieldwork is essential for phytosociological study and the calculation of diversity. Our devoted findings will help to extend the consensus and concentration of researchers to the herbaceous species of temperate ecosystems in Choti Zareen. In essence, the community under study is an early successional organized community [62, 63] with a low number of species. Species richness, as evaluated by the test proposed by Zhang et al. [64], showed variability within the different sections (or patches) of the community. Zhang et al. [65] explored the phylogenetic diversity and altitudinal patterns of species diversity across temperate mountain forests located in China. [66] examined the phyto-diversity distribution in terms of elevation gradient in Saudi Arabia. [67] conducted a study on the prediction of the soil seed bank of two different geographic arid zones (piedmont and alluvia plains) in Dera Ghazi Khan, Pakistan.

Our results also show consistency in these above-mentioned measures. It is interesting to note that [40, 51, 53, 54, 68, 69] also obtained consistency in the results of the three measures mentioned above. These three similarity measures are the most popular and are recommended for use in ecology.

Ahmad et al. [70] documented that the conservation of phyto-diversity in the Koh-Suleman range mountains has been neglected, especially at lower elevation piedmont zones (Sakhi Sarwar, lower foot hill arid zone) and upper rocky slopes located at high elevations (Fort Manru). There is a need to develop biodiversity conservation strategies for all plant communities and their habitats with the establishment of topographic preferences of these species, especially in the Koh-Suleman range mountains of Gera Ghazi Khan, Pakistan.

Conclusions

The findings of the current study concluded that the study area has 120 herb species, representing 92 genera and 35 families. Poaceaceae and Asteraceae are the dominant families, according to IVI. The dominant species, according to IVI, are *Chloris flagellifera* and *Aerva javanica.* The Shannon diversity index (H) is also moderate for this study area in all sites of herbaceous diversity, but the dominant species, *Chloris flagellifera,* showed the highest diversity, followed by *Aerva javanica*. The result of this research shows that phytosociological diversity is dominant, and their distribution is favored by vegetation that has larger seed production, well-adaptive mechanisms of seed dispersal and competes for basic needs such as soil nutrients. While phytosociological diversity and distribution patterns were distracted by anthropogenic activities and environmental pressure, from this study, it is concluded that the present status of the herbaceous layer in Choti Zareen, Dera Ghazi Khan district, is satisfactory. But in the future, a lot of challenges will be faced with the diversity of herbs if grazing and anthropogenic activity continue in their present form.

Acknowledgment

The authors extend their appreciation to the Deanship of Scientific Research at King Khalid University for funding this work through large group Research Project under grant number R.G.P. 2/223/44.

Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy

Conflict of Interest

The authors declare no conflict of interest.

References

1. OYELOWO O.J., OLADOYE A.O., OJO E.O., OLUBAYO O.O., SONDE B., ADELANI D.O. Phytosociological Assessment and Diversity of Woody Species in Omo Biosphere Reserve, Nigeria. Journal of the Cameroon Academy of Sciences, **19** (2), 125-139, **2023**.
2. AL-SHEHRI M.A., EL-SHEIKH M.A., AL-FARHAN A.H., ARIF I.A., RAJAKRISHNAN R., ALATAR A.A., FAISAL M., BASAHI R.A., AL-ABBADI G.A. Ecology of endangered *Prunus korshinskyi* Hand. Mazz. in Jabal Al-Lauz, Saudi Arabia: Plant associations, size structure, and nutritional screening. Saudi Journal of Biological Sciences, **27** (1), 147-156, **2020.**
3. DAGNE Y., BIRHANU L. Floristic composition and plant community distribution along environmental gradients in Guard dry Afromontane Forest of Northwestern Ethiopia. BMC Ecology and Evolution, **23** (1), 43, **2023**.
4. FIEBERG J.R., VITENSE K., JOHNSON D.H. Resampling-based methods for biologists. Peer Journal, **8,** e9089, **2020**.
5. BAUDOIN W.T.J., FLORE N.J., CLOVIS K.O., ELODIE M.T., SEVERIN K.F.S., JONATHAN N.M., ERIC N.C., BLERIOT F.T.A., LOUISE A.T.M., THEOPHILE F., FRANÇOIS N.V. Assessing floristic diversity and ecological characteristics of mount Mbapit Savannah, Western Highlands of Cameroon. Asian Journal of Biological Sciences, **16** (3), 351, **2023.**
6. BEN-SAID M. Spatial point-pattern analysis as a powerful tool in identifying pattern-process relationships in plant ecology: an updated review. Ecological Processes, **10**, 56, **2021**.
7. O'SHAUGHNESSY K.A., KNIGHTS A.M., HAWKINS S.J., HANLEY M.E., LUNT L., THOMPSON R.C., FIRTH L.B. Metrics matter: Multiple diversity metrics at different spatial scales are needed to understand species diversity in urban environments. Science of the Total Environment, **895**, 164958, **2023**.
8. QIN X., WU H., HUANG X., LOCK T.R., KALLENBACH R.L., MA J., ALI M.P., TU X., CAO G., WANG G., NONG X., MCNEILL M.R., ZHANG Z. Plant composition changes in a small-scale community have a large effect on the performance of an economically important grassland pest. BMC Ecology, **19** (1), 32, **2019**.
9. BOUCHET-VALAT M. General marginal-free association indices for contingency Tables: From the Altham Index to the Intrinsic Association Coefficient. Sociological Methods & Research, **51** (1), 203, **2022.**
10. LOTZE H.K. Marine biodiversity conservation. Current Biology, **31** (19), 1190-1195, **2021.**
11. WANG J., SOININEN J., HEINO J. Ecological indicators for aquatic biodiversity, ecosystem functions, human activities and climate change. Ecological Indicators, **132**, 108250, **2021**.
12. QUINTUS S., ALLEN M.S. Niche Construction and Long-Term Trajectories of Food Production. Journal of Archaeological Research, **32,** 1-53, **2023.** https://doi.org/10.1007/s10814-023-09187-x.
13. KOUL B., BHAT N., ABUBAKAR M., MISHRA M., ARUKHA A.P., YADAV D. Application of Natural Coagulants in Water Treatment: A Sustainable Alternative to Chemicals. *Water,* **14** (22), 3751, **2022**.
14. SRIDHAR K.R., PAVITHRA M. Bioactive Compounds of Ceylon Spinach [*Talinum Triangulare* (Jacq.) Willd.]. In: Murthy H.N., Paek K.Y. (eds) Bioactive Compounds in Underutilized Vegetables and Legumes. Reference Series in Phytochemistry. Springer, Cham. **2021**. https://doi.org/10.1007/978-3-030-57415-4_10.
15. NIU H.X., DUAN Q.J., SHI G.X., WU D.Q., SHAO J., WANG T.M., WANG C.Z. Inhibitory effect of extract of *Coptidis Rhizoma* on invasion of *Candida albicans* hyphae *in vitro*. Zhong Hua Zhong Yi Yao Za Zhi, **44** (1), 125, **2019**. Chinese. doi: 10.19540/j.cnki.cjcmm.20180912.004.
16. CHEN P., XIE R., LU M. Resource Conservation or Environmental Friendliness: How do Urban Clusters Affect Total-Factor Ecological Performance in China? *International Journal of Environmental Research and Public Heal*th, **17** (12), 4361, **2020.**
17. NODZA G.I., ANTHONY R.U., ONUMINYA T.O., OGUNDIPE O.T. Floristic Studies on Herbaceous and Grass Species Growing in the University of Lagos, Nigeria. Tanzania Journal of Science, **47** (1), 80, **2021.**
18. FORD J.D., TILLEARD S.E., BERRANG-FORD L., ARAOS M., BIESBROEK R., LESNIKOWSKI A.C., MACDONALD G.K., HSU A., CHEN C., BIZIKOVA L. Opinion: Big data has big potential for applications to climate change adaptation. Proceedings National Academy of Sciences, 113 **(39)**, 10729, **2016**.
19. GILARDELLI F., VERGANI C., GENTILI R., BONIS A., CHANTELOUP P., CITTERIO S., ENRICO A. Root characteristics of herbaceous species for topsoil stabilization in restoration projects. Land Degradation & Development, **28** (7), 2074, **2017**.
20. ANTÓN-PARDO M. The map of biodiversity: From local to global scales. Mètode Science Studies Journal, **9**, 83-89, **2019**.
21. RAHMAN I.U., HART R.E., AFZAL A., IQBAL Z., BUSSMANN R.W., IJAZ F., KHAN M.A., ALI H., RAHMAN S.U., HASHEM A., ABD_ALLAH E.F., SHER A., CALIXTO E.S. Vegetation-environment interactions: plant species distribution and community assembly in mixed coniferous forests of Northwestern Himalayas. Scientific Reports, **13**, 17228, **2023**.
22. RAHMAN A.U., KHAN S.M., AHMAD Z., ALAMRI S., HASHEM M., ILYAS M., AKSOY A., DÜLGEROĞLU C., KHAN G., ALI S. -Impact of multiple environmental factors on species abundance in various forest layers using an integrative modeling approach. Global Ecology and Conservation, **29**, e01712, **2021**.
23. SELCK C., DOMINGUEZ-VILLAR M. Antigen-Specific Regulatory T Cell Therapy in Autoimmune Diseases and Transplantation. Frontier in Immunology, **12**, 661875, **2021.**
24. MÉDAIL F. Plant Biogeography and Vegetation Patterns of the Mediterranean Islands. The Botanical Review, **88**, 63-129, **2022**.
25. JHARIYA M.K., BANERJEE A., YADAV D.K., RAJ A. Leguminous Trees an Innovative Tool for Soil Sustainability. In: MEENA R., DAS A., YADAV G., LAL R. (eds) Legumes for Soil Health and Sustainable Management. Springer, Singapore, pp. 315-345, **2018**.
26. ORAON M., SHARMA V. Prediction of surface roughness in single point incremental forming of AA3003-O alloy using artificial neural network. International Journal of Materials Engineering Innovation, **9** (1), 1, **2018**.

Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy

27. JIANG L., XU Q. What Do Biodiversity Experiments Tell Us About Biodiversity and Ecological Stability Relationships? In: The Ecological and Societal Consequences of Biodiversity Loss (Eds: LOREAU M., HECTOR A., ISBELL F.), **2022**. https://doi.org/10.1002/9781119902911.ch8.
28. HUGHES A.C. Understanding the drivers of Southeast Asian biodiversity loss. Ecosphere, **8** (1), e01624, **2017**.
29. HERRERO-JÁUREGUI C., SANS G.C., ANDRIES D.M., AGUIAR S., FAHRIG L., MASTRANGELO M. Past and present effects of habitat amount and fragmentation per se on plant species richness, composition and traits in a deforestation hotspot. Biological Conservation, **276**, 109815, **2022**.
30. MOLLASHAHI H., SZYMURA M., SZYMURA T.H. Connectivity assessment and prioritization of urban grasslands as a helpful tool for effective management of urban ecosystem services. PLoS One, **15** (12), e0244452, **2020**.
31. DANGOLI D.R., SHIVAKOTI G.P. Species composition and dominance of plant communities in Western Chitwan, Nepal. Nepal Journal of Science & Technology, **3**, 69, **2001**.
32. CHAUDHRY M.S., SARWAR G., MAJEED A. Natural plant communities in view of response variables and ecological indices from Lalsuhanra biosphere reserve, Pakistan. Pakistan Journal of Biological Sciences, **9** (9), 1628, **2006**.
33. SHANNON C.B., WEAVER W. The Mathematical Theory of Communication. Urbana, Illinois, **1963**.
34. PIELOU E.C. Species-diversity and pattern-diversity in the study of ecological succession. Journal of Theoretical Biology, **10** (2), 370, **1966**.
35. MAGURRAN A.E. Measuring biological diversity. Blackwell Publ., Oxford, UK, **2004**.
36. SOUTHWOOD T.R.E., HENDERSON P. Ecological Methods, 3rd edn. Blackwell Science, Oxford, **2000**.
37. PIELOU E.C. Ecological Diversity. Wiley-Inter-science, New York, pp. 165, USA, **1975**.
38. AHMAD A., SAEED A., GULSHAN A.B., ULLAH A., AHMAD B., ULLAH S., KHAN S.S., ZEB T.A. Phytodiversity of piedmont and alluvial regions of Kohe Suleman mountain range, Dera Ghazi Khan, Punjab Pakistan with emphasis on the altitudinal gradient. Journal of Xi'an Shiyou University Natural Science Edition, **19** (1), 1630, **2023.**
39. KENAR N., KIKVIDZE Z. Climatic drivers of ecological divergence among the genera of the tribe Stipeae in Turkey. Flora, **270**, 151650, **2020**.
40. WANG X., LIU Y., LI Y., JIN J., GUO Q., PEI S. Interspecies Association and Community Stability of Plants in the Core Distribution Area of *Thuja sutchuenensis*. Forests, **14**, 762, **2023**.
41. KANDEL P., GURUNG J., CHETTRI N., NING W., SHARMA E. Biodiversity research trends and gap analysis from a transboundary landscape, eastern himalayas. Journal of Asia-Pacific Biodiversity, **9** (1), 1, **2016**.
42. BRODEUR J., ABRAM P.K., HEIMPEL G.E., MESSING R.H. Trends in biological control: Public interest, international networking and research direction. BioControl, **63** (1), 11, **2018**.
43. HANIF M.A., GUO Z., MONIRUZZAMAN M., HE D., YU Q., RAO X., LIU S., TAN X., SHEN W. Plant taxonomic diversity better explains soil fungal and bacterial diversity than functional diversity in restored forest ecosystems. Plants, **8** (11), 479, **2019**.
44. GUIDEN P.W., BARBER N.A., BLACKBURN R., FARRELL A., FLIGINGER J., HOSLER S.C., KING R.B., NELSON M., ROWLAND E.G., SAVAGE K. Effects of management outweigh effects of plant diversity on restored animal communities in tallgrass prairies. Proceedings National Academy of Sciences, **118** (5), e2015421118, **2021**.
45. ZAFAR I., PERVEZ M.T., RATHER M.A., BABAR M.E., RAZA M.A., IFTIKHAR R. Genome-wide identification and expression analysis of ppos and pox gene families in the selected plant species. *Biosciences, Biotechnology* Research *Asia,* **17** (2), 301, **2020**.
46. ZAFAR I., RUBAB A., ASLAM M., AHMAD S.U., LIYAQAT I., MALIK A., ALAM M., WANI T.A., KHAN A.A. Genome-wide identification and analysis of grf (growth-regulating factor) gene family in camila sativa through in silico approaches. Journal of King Saud University Science, **34** (4), 102038, **2022**.
47. SHARMA C.M., MISHRA A.K., TIWARI O.P., KRISHAN R., RANA Y.S. Effect of altitudinal gradients on forest structure and composition on ridge tops in Garhwal Himalaya. Energy, Ecology and Environment, **2** (6), 404, **2017**.
48. YADAV S., SHARMA L. Phytosociological analysis and species diversity of herbaceous layer in Saliyatoli. International Journal of Advance and Applied Research, **2** (17), 92, **2022**.
49. FONSECA T.C.O., LOBO V.G.R., SCHMIDT A.M. Dynamical non-Gaussian modelling of spatial processes. archivePrefix={arXiv}, **2021**.
50. DING C., ZHOU X., CAO X.J, YANG J. Spatial and mediation analysis of the influences of residential and workplace built environments on commuting by car. *Transportation Research Part A: Policy and Practice,* **171**, **2023**. Article 103662. https://doi.org/10.1016/j.tra.2023.103662
51. IVASHCHENKO K.V., KORNEYKOVA M.V., SAZONOVA O.I., VETROVA A.A., ERMAKOVA A.O., KONSTANTINOV P.I., SOTNIKOVA Y.L., SOSHINA A.S., VASILEVA M.N., VASENEV V.I., GAVRICHKOVA O. Phylloplane Biodiversity and Activity in the City at Different Distances from the Traffic Pollution Source. Plant (Basel), **11** (3), 402, **2022**.
52. MALCOLM K., DIGHTON J., BARKAY T. Mercury affects the phylloplane fungal community of blueberry leaves to a lesser extent than plant age. Mycology, **9** (1), 49, **2018**.
53. HUSSAIN M., KHAN S.M., ABD_ALLAH E.F., UL HAQ Z., ALSHAHRANI T.S., ALQARAWI A.A., UR RAHMAN I., IQBAL M., ABDULLAH1–AHMAD H. Assessment of plant communities and identification of indicator species of an ecotonal forest zone at Durand Line, District Kurram, Pakistan. Applied Ecology and Environmental Research, **17** (3), 6375, **2019**.
54. BASHIR F., ABBAS A., SHAUKAT S.S., SIDDIQUI M.F., QURESHI I.A. Distribution and diversity of marine algae of Sindh coastal area: alpha, beta and gamma diversity. Pakistan Journal of Botany, **54** (6), 2377, **2022**.
55. LI T., XIONG Q., LUO P., ZHANG Y., GU X., LIN B. Direct and indirect effects of environmental factors, spatial constraints, and functional traits on shaping the plant diversity of montane forests. Ecology and Evolution, **10** (1), 557, **2020**.
56. BENITO X., FRITZ S.C., STEINITZ-KANNAN M., TAPIA P.M., KELLY M.A., LOWELL T.V. Geo-climatic factors drive diatom community distribution in tropical

south american freshwaters. Journal of Ecology, **106** (4), 1660, **2018**.
57. ZHOU S., WU N., ZHANG M., PENG W., HE F., GUO K., YAN S., ZOU Y., QU X. Local environmental, geo-climatic and spatial factors interact to drive community distributions and diversity patterns of stream benthic algae, macroinvertebrates and fishes in a large basin, northeast china. Ecological Indicators, **117**, 106673, **2020**.
58. WEIGELT A., MOMMER L., ANDRACZEK K., IVERSEN C.M., BERGMANN J., BRUELHEIDE H., FAN Y., FRESCHET G.T., GUERRERO-RAMÍREZ N.R., KATTGE J. An integrated framework of plant form and function: The belowground perspective. New Phytologist, **232** (1), 42, **2021**.
59. TAO Z., GE Q., WANG H., DAI J. The important role of soil moisture in controlling autumn phenology of herbaceous plants in the inner mongolian steppe. Land Degradation & Development, **32** (13), 3698, **2021**.
60. BIRBEN Ü. The effectiveness of protected areas in biodiversity conservation: the case of Turkey. Cerne, **25** (4), 424, **2019.**
61. TAVŞANOĞLU Ç. Anadolu Bozkır Ekosistemleri Üzerinde İşleyen Müdahale Rejimleri. Kebikec: Insan Bilimleri Icin Kaynak Arastirmali Dergisi, **43**, 259, **2017**.
62. ZANG Z., ZENG Y., WANG D., SHI F., DONG Y., LIU N., LIANG Y. Species-Abundance Distribution Patterns of Plant Communities in the Gurbantünggüt Desert, China. *Sustainability,* **14**, 12957, **2022**.
63. FRĄC M., HANNULA S.E., BEŁKA M., JĘDRYCZKA M. Fungal biodiversity and their role in soil health. Frontiers in Microbiology, **9**, 707, **2018**.
64. ZHANG Z., HAUTIER Y., BAO T., YANG J., QING H., LIU Z., WANG M., LI T., YAN M., ZHANG G. Species richness and asynchrony maintain the stability of primary productivity against seasonal climatic variability. Frontiers in Plant Science, **13**, 1014049, **2022**.
65. ZHANG W., HUANG D., WANG R., LIU J., DU N. Altitudinal patterns of species diversity and phylogenetic diversity across temperate mountain forests of northern China. Plos one, **11** (7), 1, **2016**.
66. MSEDDI K., ALGHAMDI A., ABDELGADIR M., SHARAWY S., CHAIEB M., MILLER T. Phytodiversity distribution in relation to altitudinal gradient in Salma Mountains-Saudi Arabia. Global Ecology and Conservation, **27** (2), 67, **2021**.
67. AHMAD A., SAEED A., GULSHAN A., WALI S., HADI F., ULLAH S., SHER A., RIZWAN M., RAFIQ M. Prediction of soil seed bank of piedmont and alluvial environments of Dera Ghazi Khan, Punjab, Pakistan. Brazilian Journal of Biology, **84** (2), 234, **2021**.
68. SUN Z., ZHANG W., LIU Y., DING C., ZHU W. The changes of phyllosphere fungal communities among three different *Populus* spp. *Microorganisms,* **11**, 2479, **2023**.
69. WANG X., ELLWOOD M.D.F., AI D., ZHANG R., WANG G. Species abundance distributions as a proxy for the niche–neutrality continuum. Journal of Plant Ecology, **11** (3), 445, **2018**.
70. AHMAD A., SAEED A., GULSHAN A.B., YOUSAF W., ZAFAR I. Envisaging natural vegetation in contrasting environments (piedmont and alluvial) of Dera Ghazi Khan, Pakistan. Pakistan Journal of Botany, **55** (6), 2231, **2023**.

Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy • Author Copy