

Original Research

Improving Maize Physiological Attributes by Regulating Urease Activity and Zinc Availability in Rhizosphere Through Bioactivated Zinc-Coated Urea (Engro Zabardast Urea)

Muhammad Hasnain¹, Qudsia Nazir^{2*}, Abubakar Dar³, Azhar Hussain³, Sajid Mahmood⁴, Ifra Saleem¹, Hafiz Tanvir Ahmad^{3, 5}, Syed Shahid Hussain Shah¹, Muhammad Asif Ali¹, Muhammad Naveed⁶, Rashid Iqbal^{7**}, Abd El-Zaher M.A. Mustafa⁸, Mohamed Soliman Elshikh⁸, Allah Ditta^{9, 10***}

¹Engro fertilizers Limited, 7th & 8th floor, the Harbor Front Building, HC-3 Marine Drive Block-4, Clifton Karachi-75600, Pakistan

²Institute of Soil Chemistry and Environmental Sciences, Ayub Agriculture Research Institute, Faisalabad, 38000, Pakistan

³Department of Soil Science, The Islamia University of Bahawalpur-63100, Pakistan

⁴Soil and Environmental Sciences Division, Nuclear Institute of Agriculture and Biology (NIAB), Faisalabad, Pakistan

⁵Provincial Reference Fertilizer Testing Laboratory, Raiwind, Lahore 55150, Pakistan

⁶Institute of Soil & Environmental Sciences, University of Agriculture Faisalabad-38040, Pakistan

⁷Department of Agronomy, Faculty of Agriculture and Environment, The Islamia University of Bahawalpur, Bahawalpur 63100, Pakistan

⁸Department of Botany and Microbiology, College of Science, King Saud University, P.O. 2455, Riyadh 11451, Saudi Arabia

⁹Department of Environmental Sciences, Shaheed Benazir Bhutto University Sheringal, Dir (U), Khyber Pakhtunkhwa 18000, Pakistan

¹⁰School of Biological Sciences, The University of Western Australia, 35 Stirling Highway, Perth, WA 6009, Australia

Received: 17 January 2024

Accepted: 30 April 2024

Abstract

The major cause of reduced Zn availability to plants in Pakistan is unfavorable soil factors (alkaline pH and calcareousness). Different strategies can be adopted, to enhance the Zn bioavailability in the rhizosphere and cereals. Among all strategies, a novel technique is the coating of Zn on macronutrient fertilizers like urea. By adopting this technique, dual benefits can be achieved for example Zn and N become available to plants and on the other hand, the loss of urea can be minimized. Firstly, Engro Zabardast Urea was taken from Engro Fertilizers Pvt. Ltd and evaluated for coating effect on

*e-mail: sbqnazir@gmail.com

**e-mail: rashid.iqbal@iub.edu.pk

***e-mail: allah.ditta@sbbu.edu.pk

urease activity and Zinc release pattern with time (up to 70 days). Then 100%, 90%, 85%, and 80% recommended EZU (Engro Zabardast Urea) were tested to find out the increase in urea efficiency by coating. From the results of the release experiment, it was concluded that EZU @ 90% of the recommended dose showed maximum results of Zn release respective to time and vice versa for urease activity. Secondly, a pot experiment was conducted to evaluate the above-mentioned prepared products on agronomic parameters and Zn and nitrogen concentration in 45 days of maize seedlings. Zn concentration in maize seedlings increased with the application of 90 and 85% of the recommended EZU, respectively. Similar results were found under field conditions with the application of 90% of recommended EZU growth, physiological parameters, and grains and stover yield were improved and Zn contents in grains were enhanced. From the results of all experiments, it can be concluded that EZU is a fantastic product not only to increase Zn bioavailability/fortification but also to reduce nitrogen losses. This product has the potential to reduce the use of plain urea by up to 10%. This ecofriendly approach cannot only reduce extra labor costs but also the ignorance of Zn application by the farming community can also be addressed.

Keywords: EZU, maize seedlings, urease activity, physiological parameters

Introduction

Balanced nutrition is an important factor in enhancing the overall yield of different crops [1]. The farming community of Pakistan prefers to use macronutrients i.e. (N) nitrogen, (P) phosphorous, and (K) potassium fertilizers only [2]. However, after N and P, Zn deficiency is widespread, common, and well reported due to unfavorable Pakistani soil conditions i.e., high CaCO_3 , less organic matter in the calcareous soils, having high pH [3-6]. Zn malnutrition is becoming a severe issue in the developing world and the victims are mostly infants, underaged children, and pregnant women [7]. Moreover, the nitrogen fertilizer losses from the cultivated lands are becoming a major issue due to high temperatures and arid to semi-arid climatic conditions of the country. Nitrogen has a synergistic relationship with Zn and is an essential nutrient for plants and humans [8, 9]. On average $70 \mu\text{g g}^{-1}$ Zn is present in the earth's crust (0.02%) by weight [10, 11] but anthropogenic activities are among the major factors affecting Zn availability and release in the soil [12] because Zn is very less mobile within the soil that's why its deficiency in plants is well reported [13]. Optimal Zn in soil is 0.6-1.0 mg kg⁻¹ (DTPA-extractable), and 10-20 mg kg⁻¹ of dry weight in plants [14, 15]. The limits may differ with soil type and crop rotation [16].

Nearly 70% of soils in Pakistan are deficient in Zn [17, 18]. Soils having characteristics such as saline-sodic, high pH, calcareousness, and high cropping intensity are mostly deficient in Zn contents [19, 20]. Its prevalence in the soil is mostly in the form of Zn minerals which are ZnS, (ZnSO_4) zinkosite, (ZnCO_3) smithsonite, (ZnO) zincite, [$\text{Zn}_3(\text{PO}_4)_2 \cdot 4\text{H}_2\text{O}$] hopeite, (ZnFe_2O_4) franklinite, the availability of Zn from these sources depends upon some natural phenomena, for example, weathering of parent rocks, surface dusting and volcanoes [21-23]. Traditionally Zn is applied to soil through broadcasting with almost 50% use efficiency and the remaining

gets fixed either with minerals or CaCO_3 [24]. Due to its non-availability in Pakistani soils having high pH and calcareousness, the use of coated fertilizers/slow-release fertilizers is becoming popular day by day all over the world. It is among the major strategies to improve environmental quality and sustain agricultural productivity [6, 25-27]. The application of coated fertilizers for better growth and yield is becoming an emerging trend/novel technology in today's agriculture in the future [28]. Various strategies are adopted to increase Zn availability in soil for plants. The second thing is the loss of applied urea in the form of ammonia to the atmosphere is a prominent issue. With the practice of coating, this loss can be minimized up to 15% [29].

Different Zn fertilizers are available to fulfill the requirement of plant Zn but the emerging fertilizer technologies use Zn and urea side by side through Zn coating on urea at varying concentrations [6, 30]. Many benefits can be achieved by coating Zn on urea such as i) the application of Zn and nitrogen to plants side by side because Zn application to crops is neglected among farmers' community, ii) urea loss in the form of ammonia can be minimized up to 10%, iii) slow and continuous supply of nitrogen to plants with very less loss, and iv) Ecofriendly approach with less labor cost [31, 32]. Based on the above discussion, the present study was conducted with the following objectives: i) to test the efficacy of commercialized bioactivated zinc coated urea (EZU) of Engro fertilizers Pvt. Ltd. in terms of soil urease activity and Zn release pattern with specific time intervals, ii) to test the effect of EZU on the growth and Zn contents of maize seedlings in pot studies and iii) to check the improvement in physiological attributes of maize and Zn fortification in grains through application of EZU and common Zn and N sources. It can be hypothesized that the application of EZU may improve the nutritional status of maize and reduce the fixation of Zn and ammonia volatilization in the field.

Experimental

The hypothesis was tested by planning three distinct studies. An incubation study to check the urease activity and release pattern of Zn from EZU at specific time intervals. Pot and field studies at the wirehouse, and field conditions at the Engro center, University of Agriculture, Faisalabad-Pakistan to compare the effect of EZU with ZnSO₄ and urea separately for improving the yield, physiology, and Zn fortification in maize grains (hybrid 4040).

Physicochemical Characteristics of Soil

Soil samples were mix thoroughly after drying, grinding and passing through 2 mm mesh size sieve, determined for physicochemical characteristics: texture, clay loam; organic matter, 0.77% [33]; similarly, pH, 8.22 EC, 1.27 dSm⁻¹ [34]; available phosphorus, 9.17 mg kg⁻¹ [35]; extractable potassium 95 mg kg⁻¹ [34]; and plant available Zn, 0.97 mg kg⁻¹ [36].

Characteristics of EZU Engro Fertilizers Pvt. Ltd.

The Zabardast Urea is a synergetic hybrid of urea, Bioactive Zinc (BAZ)© and Bioactive Coating (BAC)©; a consortium of Zn and other nutrients solubilizing and mobilizing bacteria. BAZ© is organically encapsulated Zn that is less prone to fixation, sandwiching, and trapping in soil structure. BAZ© is gradually released in the rhizosphere as per plant demand that supports an uninterrupted continuous supply of Zn during the crop cycle. In addition, BAC© enhances root growth, mobilizes other nutrients present in the rhizosphere, and triggers induced systemic resistance of plants to healthily pass through stress conditions. The coating cover of BAZ© and BAC© encapsulates urea prills induces a slow N release mechanism, contributes to reducing N losses, and enhances N use efficiency. Collectively, Zabardast urea is a revolutionary fertilizer suitable for all types of soils, climates, and crops [37].

Experimental Description

Experiments were conducted with six treatments along with one control (no Zn) and three replications in the experiments treatment plan as described in Table 1. ZSB was taken from the Department of Soil Science, The Islamia University, Bahawalpur. To conduct the first and second experiments Completely Randomized Design (CRD) was used with three repetitions. Randomized complete block design (RCBD) with three blocks was adopted in the third experiment. Fertilizers were applied as 275, 125, and 75 kg ha⁻¹ NPK recommended doses along with zinc dose as 15 kg ha⁻¹. For the irrigation, tap water was used in the first and second experiments while tube well water was used in the third experiment. In the first experiment, Zn release with time and urease activity was evaluated while in the second experiment maize seedlings of 1.5 months were collected for analysis. In the third experiment, the crop was harvested at maturity, and plant height, biomass, cob length, cob diameter, and yield (stover yield, grain yield, grains Zn concentration, and physiological parameters (with CIRUS) were taken.

Measurement of Zn in Grain

Wet digestion of grain samples was done by following the method described by Jones and Case, [38] and Zn contents in grain were determined from the digested sample on Atomic Absorption Spectrophotometer (PerkinElmer, Analyst 100, Waltham, USA).

Statistical Analysis

The data regarding different parameters was collected and subjected to analysis of variance (ANOVA), through Statistix v. 8.1 (Analytical Software, USA) software. The least significant difference (LSD) test at a 5% probability was used to evaluate/compare the treatment means [39]. The significance of the treatment was presented through alphabetical lettering. The treatment means having the same letters was considered statistically non-significant at p≤0.05.

Table 1. Treatment description of the experiment.

Treatments	Description
T ₀	Control (recommended N, P, and K with no Zn).
T ₁	ZSB (Zinc solubilizing Bacteria) along with (RD) recommended dose of N, P, and K.
T ₂	ZnSO ₄ + urea and recommended P and K.
T ₃	100% of EZU with recommended P and K
T ₄	90% of EZU with recommended P and K
T ₅	85% of EZU with recommended P and K
T ₆	80% of EZU with recommended P and K.

Results

Wirehouse and field area of Engro center, University of Agriculture, Faisalabad was used for Pot and field experiments to find out the efficiency of EZU on growth, yield, and physiology of maize crop and results are given as follows:

Axenic Conditions Study

Zn Release Pattern and Urease Activity

The results regarding Zn release depicted an increasing trend in the release pattern of Zn by application of EZU than control (no Zn) as shown in Fig. 1a). A constant and gradual increase was observed in the pots, which received 90% EZU. On the first day, Zn concentration was 0.76 mg kg^{-1} , which raised to 1.92 mg kg^{-1} Zn at the end of incubation, which was 68% more Zn contents than first-day readings. While an increase of 66% was obtained with the application of 85% EZU. In the treatments where ZnSO_4 and plain urea were applied, 1.1 mg kg^{-1} Zn contents were determined after incubation. Furthermore, 85% EZU produced similar results as ZnSO_4 was applied with plain urea after

60 days, no further increase was observed with time. All the results that were obtained showed a statistically significant interactive effect of EZU. On the other hand, the application of ZSB alone has caused a slow increase (12%) in Zn contents.

As compared to all other treatments, 90% EZU has caused a continuous increase in urease activity throughout the incubation period. At the start of the experiment, the urease activity of $10\text{--}12 \text{ mg NH}_4\text{-N kg}^{-1} \text{ h}^{-1}$ was observed under 90% EZU application and a gradual decline of 5.5% in urease activity was observed. Under control (without Zn) $70 \text{ mg NH}_4\text{-N kg}^{-1} \text{ h}^{-1}$ activity at the end of the incubation. While $66 \text{ mg NH}_4\text{-N kg}^{-1} \text{ h}^{-1}$ activity was observed under 90% EZU application. Furthermore, almost similar and constant results were detected from 50-70 days. The experimental units receiving 85 and 80 EZU showed 66 and $67 \text{ mg NH}_4\text{-N kg}^{-1} \text{ h}^{-1}$ activities, respectively. On the 70th day, the soil analysis showed that with the application of ZnSO_4 and plain urea separately $70 \text{ mg NH}_4\text{-N kg}^{-1} \text{ h}^{-1}$ activity was recorded (Fig. 1b).




Fig. 1. Temporal release of Zn a) and urease activity b) from ZnSO_4 and EZU under controlled conditions.

Pot Experiment

Plant Height and Biomass Production of Maize Seedlings

Different levels of EZU (100, 90, 85, and 80% of EZU) had a significant effect on plant height and biomass (g) of maize seedlings. With the application of different levels of EZU, an increasing trend was obtained in the biomass (g) of maize seedlings. Only the individual application of ZSB showed a minor increase in biomass production. Statistical analysis depicted significance at different levels of EZU; however, the maximum biomass (10.8 g) was recorded under 90% EZU treatment while the minimum height (7.6 g) was recorded without Zn application. With the application of 90% EZU, a 42% increase was observed as compared to control, similar results were found in plant height (Table 2).

Zn and N Concentration in Maize Seedlings

The impact of different levels of EZU (100, 90, 85, and 80% of EZU) showed statistically significant results on Zn and N concentration of maize seedlings. From the results presented, it was clear that 60.7 ($\mu\text{g g}^{-1}$) Zn and 1.56% N were found in maize seedlings with the application of 90% EZU. In the concentration of Zn and N a maximum of 46% and 12% increase was obtained respectively. On the other hand, 43 and 7% increase was obtained in maize seedlings respectively with the application of 85% of EZU (Table 2).

Field Experiment

Effect of EZU on Cob Length and Cob Diameter of Maize

Regarding the effect of different levels of EZU, 90% of EZU depicted a significant ($p < 0.05$) increase in the cob length and cob diameter. There was observed a 6 and 7.6% increase in cob length and cob diameter,

respectively, for T₀ (no Zn). On the contrary, the treatment where ZnSO₄ was applied with plain urea instead of EZU, showed 4 and 6.3% less cob length and cob diameter as compared to 90% EZU. Significant (statistically) results were found in the treatments where 85 and 80% EZU was applied than control and ZnSO₄ and urea application separately (Table 3).

Effect of EZU on the Physiology of Maize

Physiological parameters; net photosynthesis, transpiration rate, substomatal CO₂, stomatal conductance, and chlorophyll contents were determined because Zn has a major role in plant physiology. From the results, it was observed that a significant effect at different levels of EZU was noted. Maximum net photosynthesis, substomatal CO₂ concentration, stomatal conductance, and chlorophyll contents were found in the treatment where 90% EZU was applied, it was 15, 21.5, 31, and 45% increase respectively. An increase of 8.5, 20.6, 14.5, 17, and 39% was noted in net photosynthesis, transpiration rate, substomatal CO₂ concentration, stomatal conductance, and chlorophyll contents, respectively with the application of 85% of EZU (Table 3).

Effect of EZU on Stover Yield, Grain Yield, and Grain Zn and N Contents in Maize

Results of yield attributes depicted statistically significant results at different levels of EZU noted on stover yield, grain yield, and Zn and N contents in grains (Fig. 2). Maximum results were found in the treatment where 90% EZU was applied. Stover and grains yield (8.83 and 6.04 t ha⁻¹) was obtained with the application of 90% EZU. The application of 85% EZU showed 7.9 and 6 t ha⁻¹, stover, and grains yield respectively (Fig. 2-A). On the other hand, a similar trend in grain Zn contents was observed (Fig. 2-B). Zn contents 53.4, 51.2, and 44 $\mu\text{g g}^{-1}$ were observed with the application of 90,

Table 2. Efficacy of EZU against ZnSO₄ in improving Plant height, maize biomass, Zinc, and nitrogen concentration in maize seedlings in a pot experiment.

Treatment	Plant height (cm)	Biomass (g pot ⁻¹)	Zn concentration ($\mu\text{g g}^{-1}$)	N concentration (%)
T ₀	59.1 d	7.6 d	32.5 f	1.37 d
T ₁	62.8 c	9.1 c	54.5 e	1.52 a
T ₂	67.9 ab	10 b	58.4 bc	1.56 a
T ₃	68.4 ab	9.9 b	59.1 b	1.54 a
T ₄	69.4 a	10.8 a	60.7 a	1.56 a
T ₅	67.9 ab	9.2 c	58 c	1.48 b
T ₆	67.9 b	9.0 c	56.8 d	1.41 c
LSD ($p \leq 0.05$)	1.9	0.48	0.96	0.035

In a column, means sharing the same letters are statistically non-significantly different at $p \leq 0.05$.

Table 3. Efficacy of EZU against ZnSO₄ in improving agronomic and physiological attributes of maize under field experiment.

Treatment	Cob length (cm)	Cob diameter (cm)	Pn (mmol m ⁻² s ⁻¹)	E (mmol m ⁻² s ⁻¹)	Ci (mmol m ⁻² s ⁻¹)	Gs (mmol m ⁻² s ⁻¹)	Chlorophyll contents (SPAD value)
T ₀	12.64 c	11.56 d	12.65 e	4.34 d	200.3 d	0.24 c	19.4 d
T ₁	12.67 c	11.73 c	12.94 c	4.56 d	215.3 c	0.26 c	22.0 d
T ₂	12.89 b	11.89 c	14.96 a	5.68 ab	249.7 a	0.34 ab	35.4 a
T ₃	13.6 a	12.54 ab	14.97 a	5.78 a	255.4 a	0.38 a	36.1 a
T ₄	13.5 a	12.7 ab	14.95 a	5.68 ab	255.0 a	0.35 a	35.4 a
T ₅	13.68 a	12.49 b	13.830 b	5.47 b	234.4 b	0.29bc	32.2 b
T ₆	12.8 bc	11.85 c	12.76 d	5.07 c	219.7 c	0.26 c	28.6 c
LSD (<i>p</i> ≤0.05)	0.17	0.16	0.08	0.23	11.00	0.0569	3.06

In a column, means sharing the same letters are statistically non-significantly different at *p*≤0.05. Pn: Net Photosynthesis, E: Rate of transpiration, Ci: Substomatal CO₂ concentration, and gs: Stomatal conductance

85 and 80% EZU. The treatment where ZnSO₄ was applied with plain urea showed 51 µg g⁻¹ Zn which is comparable to 85% EZU (Fig. 2-C).

Nitrogen contents in grains (%) were increased with the application of EZU (Fig. 2-D), and with the application of 90% EZU 30.4% N contents in grains were

increased than control where recommended plain urea was applied with no Zn. Similarly, a 24.8% increase with 90% EZU was obtained as compared to the application of ZSB only. The application of recommended plain urea with ZnSO₄ showed comparable results with the treatment where 80% EZU was applied.




Fig. 2. Efficacy of EZU against ZnSO₄ in improving stover yield (A), Grain Yield (B), Grain Zn contents (C), and Grain N contents (D) under field experiment.

Discussion

Zinc sulfate is the most common Zn source, but its use is limited due to its high price and not availability in time. On the contrary, ZnO is a cheaper source of Zn and contains 80% of Zn but an insoluble source. This insoluble Zn contents in ZnO can be made soluble through Zinc solubilizing bacteria (ZSB) [40]. The solubilization of ZnO with ZSB could be termed as bioactivation of Zn which bacteria do through the production of various organic acids and metabolites [41]. Further, Zn is a vital metallic ion that performs as a cofactor in almost all six classes of enzymes, considered the second most significant metal after iron present in living organisms [42]. However, its deficiency in various crop plants is well reported due to fewer uptakes in Pakistani soils with alkaline pH, calcareousness, and low organic matter. Most of the Pakistani soils are Zn deficient and to overcome Zn deficiency farmers are using commercial fertilizer of Zn in the form of $ZnSO_4$ [6, 43, 44]. To avoid Zn deficiency one effective method is the coating of Zn on urea, in this way, dual benefits can be achieved, the first benefit is the simultaneous application of Zn and nitrogen side by side, and the second benefit is the reduction of urea loss and improvement in nitrogen uptake [45].

In the present study, Zn-coated urea (EZU) was obtained from Engro Fertilizers Pvt. Ltd. and tested on maize crops, first, the Zn release and urease activity were evaluated. We found that their release was slow and steady which indicated that with coating, urea becomes available for a longer period to the plants, and its loss can be minimized up to 15-20% as described by Ali et al. [46]. The effect of EZU on growth and yield attributes of maize crops is more as compared to only the use of Zn as $ZnSO_4$ and plain urea. Application of nitrogen (90%) with EZU showed maximum results in agronomic parameters because of its availability for longer periods, higher uptake in plants, and reduced adsorption on clay and precipitation with other salts as mentioned by Mirbolook et al. [47]. It is evident from the literature that with the application of ZSB, the insoluble Zn compounds can be solubilized in calcareous, high pH soils as these bacteria are blessed with certain mechanisms of metals solubilization mainly organic acids production, siderophore production, and EPS production [48]. Bioactivated zinc-coated urea (EZU) can solubilize Zn by producing organic acids, especially in cereals, and increasing overall growth, yields, and grain Zn concentration through mineral solubilization as described by Ali et al. [43].

Zinc amount can also be saved by the application of Zn-coated urea, as 2.8 kg Zn ha⁻¹ was applied with Zn ZU whereas 6 kg Zn ha⁻¹ is recommended in soil and foliar applications as compared by Prasad et al. [49]. Hence, it is an effective, eco-friendly, and economical fertilizer in poor countries with smallholding farmers [47]. Maximum results in stover and grain yields were obtained in 90% EZU and comparable results were

found in 85% EZU with $ZnSO_4$ and plain urea. From the studies, it was found that with the help of coating almost 10-15% urea could be saved. The present findings are in line with the findings of Pooniya et al. [50]. Many researchers have done work on coated fertilizers and found similar results because with coating the hydrolysis of urea is delayed. We can say that a slow and constant supply of N can be achieved with the application of coated fertilizers [51].

By coating and bioactivation (EZU) urease activity was regulated and slow nitrogen availability was observed. Zinc uptake was also improved by plants. Zn acts as a cofactor in many enzymes that's why with the application of EZU all physiological parameters, for example, net photosynthesis, transpiration, and chlorophyll contents were improved as described by Mumtaz et al. [52]. Plant physiological parameters were improved at maximum level with the application of 90% EZU as compared to plain urea; the possible reason is that Zn acts as a cofactor in carbonic anhydrase enzyme as described by many scientists [48, 53, 54]. Nazir and her coworkers [6] concluded that the farmers of poor/developing communities can get the maximum benefit from the use of this economical, efficient, and eco-friendly bio-activated Zn-coated urea (EZU) by their limited resources and more nutrition in maize grain can be obtained.

Conclusions

It can be concluded from our salient findings that the application of Bioactivated Zn-coated urea (EZU) has a marked effect not only on plant growth but also on (Zn) concentration in plants as well. Different levels of EZU (90, 85, and 80 %) were used but 90% EZU showed maximum increase in maize seedlings Zn concentrations. N acquisition. Moreover, the uptake of Zn and N improved the physiological attributes (photosynthesis, transpiration rate, stomatal conductance, etc.) that are responsible for higher maize yields. On the other hand, we can say that it is an economical, novel, and easy technique as compared to the use of commercial Zn as $ZnSO_4$ and urea distinctly. This approach is very effective in saving 10 to 15% urea without sacrificing the yield. Ongoing research, farmer education, and support through regulatory incentives and research funding are essential for the successful implementation and sustainability of these strategies, contributing to more resilient and productive agricultural systems.

Acknowledgments

The authors extend their appreciation to the Researchers Supporting Project number (RSPD2024R941), King Saud University, Riyadh, Saudi Arabia. Highly acknowledged to Engro Fertilizers Pvt.

Ltd. for administrative and technical support to execute and perform this research.

Conflict of Interest

The authors declare no conflict of interest.

References

- HUANG S., WANG P., YAMAJI N., MA J.F. Plant nutrition for human nutrition: hints from rice research and future perspectives. *Molecular Plant*, **13** (6), 825, **2020**.
- REHMAN A., CHANDIO A.A., HUSSAIN I. JINGDONG L. Fertilizer consumption, water availability, and credit distribution: Major factors affecting agricultural productivity in Pakistan. *Journal of the Saudi Society of Agricultural Sciences*, **18** (3), 269, **2019**.
- RASHID A., RYAN J. Micronutrient constraints to crop production in the Near East: Potential significance and management strategies. In *Micronutrient deficiencies in global crop production*. B. J. Alloway Eds. Springer, Dordrecht, Netherlands, pp149, **2008**.
- ALLOWAY B.J. Soil factors associated with zinc deficiency in crops and humans. *Environmental Geochemistry Health*, **31** (5), 537, **2009**.
- ZEB H., HUSSAIN A., NAVEED M., DITTA A., AHMAD S., JAMSHAD M.U., AHMAD H.T., HUSSAIN B., AZIZ R., HAIDER M.S. Compost enriched with ZnO and Zn-solubilizing bacteria improves yield and Zn-fortification in flooded rice. *Italian Journal of Agronomy*, **13** (4), 310, **2018**.
- NAZIR Q., ARSHAD M., AZIZ T. SHAHID M. Influence of zinc impregnated urea on growth, yield and grain zinc in rice (*Oryza sativa*). *International Journal of Agriculture and Biology*, **18**, 1195, **2016**.
- YADAV A.K., SETH A., KUMAR V. DATTA A. Agronomic Biofortification of Wheat Through Proper Fertilizer Management to Alleviate Zinc Malnutrition: A Review. *Communications in Soil Science and Plant Analysis*, **54** (2), 154, **2023**.
- MAJEED A., RASHID I., NIAZ A., DITTA A., SAMEEN A., AL-HUQAIL A.A., SIDDIQUI M.H. Balanced use of Zn, Cu, Fe, and B improves the yield and sucrose contents of sugarcane juice cultivated in sandy clay loam soil. *Agronomy*, **12** (3), 696, **2022**.
- MONIKA G., KIM S.R.M., KUMAR P.S., GAYATHRI K.V., RANGASAMY G., SARAVANAN A. Biofortification: A long-term solution to improve global health-a review. *Chemosphere*, 314, 137713, **2023**.
- LEACH D.L., BRADLEY D.C., HUSTON D., PISAREVSKY S.A., TAYLOR R.D., GARDOLL S.J. Sediment-hosted lead-zinc deposits in Earth history. *Economic Geology*, **105** (3), 593, **2010**.
- KANIA H. SATERNUS M. Evaluation and Current State of Primary and Secondary Zinc Production - A Review. *Applied Science*, **13** (3), e2003, **2023**.
- CHASAPIS C.T., NTOUPA P.S.A., SPILIOPOULOU C.A., STEFANIDOU M.E. Recent aspects of the effects of zinc on human health. *Archives of Toxicology*, **94**, 1443, **2020**.
- ATSDR, Toxicology profile for Zinc, US Department of Health and Human Services, **2005**.
- FAROOQ M.S., UZAIR M., RAZA A., HABIB M., XU Y., YOUSUF M., YANG S.H., KHAN M.R. Uncovering the research gaps to alleviate the negative impacts of climate change on food security: a review. *Frontiers in Plant Science*, **13**, 927535, **2022**.
- NATASHA N., SHAHID M., BIBI I., IQBAL J., KHALID S., MURTAZA B., BAKHAT H.F., FAROOQ A.B.U., AMJAD M., HAMMAD H.M., NIAZI N.K. Zinc in the soil-plant-human system: A data-analysis review. *Science of the Total Environment*, **808**, e152024, **2022**.
- ASLE-MOHAMMADI Z., KHARAZMI M., SHEIKHI H., MOHAMMADKHANI N., NICOLA S. Foliar Application of Fe, Zn, and Mn as a Practical Strategy to Alleviate the Soil Cu Toxicity and Stimulate the Physiological and Biochemical Properties of Peppermint (*Mentha piperita* L.). *Journal of Soil Science and Plant Nutrition*, **1**, **2023**.
- SAHI A.C., BABAK M., HOSSEIN M.S.H., IMAN J., DADRASNIA A. Effect of zinc supplements on the nutritional status of different wheat genotypes. *Journal of Plant Nutrition*, **1**, **2023**.
- JALAL A., OLIVEIRA C.E.D.S., FERNANDES G.C., DA SILVA E.C., DA COSTA K.N., DE SOUZA J.S., LEITE G.D.S., BIAGINI A.L.C., GALINDO F.S., TEIXEIRA FILHO M.C.M. Integrated use of plant growth-promoting bacteria and nano-zinc foliar spray is a sustainable approach for wheat biofortification, yield, and zinc use efficiency. *Frontiers in Plant Science*, **14**, 1146808, **2023**.
- ALLOWAY B.J. Zinc in soils and crop nutrition, 2nd Ed. International Fertilizer Industry Association, Paris, France, **2008**.
- YOUNAS N., FATIMA I., AHMAD I.A., AYYAZ M.K. Alleviation of zinc deficiency in plants and humans through an effective technique; biofortification: A detailed review. *Acta Ecologica Sinica*, **43** (3), 419, **2023**.
- ALLOWAY B.J. Heavy metals in soils, 2nd Ed. Blackie academic & professional, London, **1995**.
- International Zinc Association IZA. <http://www.zinc.org/sustainability>. (Accessed December, 15, **2011**).
- PARASHAR R., AFZAL S., MISHRA M., SINGH N.K. Improving biofortification success rates and productivity through zinc nanocomposites in rice (*Oryza sativa* L.). *Environmental Science and Pollution Research*, **30** (15), 44223, **2023**.
- KLOFAC D., ANTOSOVSKY J., SKARPA P. Effect of Zinc Foliar Fertilization Alone and Combined with Trehalose on Maize (*Zea mays* L.) Growth under the Drought. *Plants*, **12** (13), p.2539, **2023**.
- MARTINEZ-CUESTA N., CARCIOCHI W., SAINZ-ROZAS H., SALVAGIOTTI F., COLAZO J.C., WYNGAARD N., EYHERABIDE M., FERRARIS G., BARBIERI P. Effect of zinc application strategies on maize grain yield and zinc concentration in mollisols. *Journal of Plant Nutrition*, **44** (4), 486, **2021**.
- HUSSAIN A., ZAHIR Z.A., DITTA A., TAHIR M.U., AHMAD M., MUMTAZ M.Z., HUSSAIN S. Production and Implication of Bio-Activated Organic Fertilizer Enriched with Zinc-Solubilizing Bacteria to Boost up Maize (*Zea mays* L.) Production and Biofortification under Two Cropping Seasons. *Agronomy*, **10** (1), e39, **2019**.
- NATARELLI C.V. LOPES C.M. CARNEIRO J.S. MELO L.C. OLIVEIRA J.E. MEDEIROS E.S. Zinc slow-release systems for maize using biodegradable PBAT nanofibers obtained by solution blow spinning. *Journal of Material Science*, **56**, 4896, **2021**.

28. MIKULA K., IZYDORCZYK G., SKRZYPCZAK D., MIRONIUK M., MOUSTAKAS K., WITEK-KROWIAK A., CHOJNACKA K. Controlled release micronutrient fertilizers for precision agriculture – A review. *Science of the Total Environment*, **712**, e136365, **2020**.
29. KLIMCZYK M., SICZEK A., SCHIMMELPFENNIG L. Improving the efficiency of urea-based fertilization leading to a reduction in ammonia emission. *Science of the Total Environment*, **771**, e145483, **2021**.
30. MUSTAFA A., ATHAR F., KHAN I., CHATTHA M.U., NAWAZ M., SHAH A.N., MAHMOOD A., BATOOL M., ASLAM M.T., JAREMKO M., ABDELSALAM N.R. Improving crop productivity and nitrogen use efficiency using sulfur and zinc-coated urea: A review. *Frontiers in Plant Science*, **13**, e942384, **2022**.
31. SHIVAY Y.S., POONIYA V., PAL M., GHASAL P.C., BANA R., JAT S.L. Coated urea materials for improving yields, profitability, and nutrient use efficiencies of aromatic rice. *Global Challenges*, **3** (12), e900013, **2019**.
32. NAZIR Q., WANG X., HUSSAIN A., DITTA A., AIMEN A., SALEEM I., NAVEED M., AZIZ T., MUSTAFA A., PANPLUEM N. Variation in growth, physiology, yield, and quality of wheat under the application of different zinc coated formulations. *Applied Science*, **11** (11), e4797, **2021**.
33. MOODIE C.D., SMITH H.W., MCCREERY R.A. *Laboratory Manual for Soil Fertility*. Department of Agronomy, State College of Washington Pullman, Washington, USA, **1**, **1959**.
34. ALLISON L.E., BERNSTEIN L., BOWER C.A., BROWN J.W., FIREMAN M., HATCHER J.T., HAYWARD H.E., PEARSON G.A., REEVE R.C., RICHARDS L.A., WILCOX L.V. *Diagnosis and improvement of saline and alkali soils*. USDA Hand Book No. 60. Washington, D.C., USA, **1954**.
35. WATANABE F.S., OLSEN S.R. Test of an ascorbic acid method for determining phosphorous in water and NaHCO₃ extracts. *Soil Science Society of America*, **29**, 677, **1965**.
36. SOLTANPOUR P.N., WORKMAN S.M. Modification of the NaHCO₃ DTPA soil test to omit carbon black. *Communication in Soil Science and Plant Analysis*, **10**, 1411, **1979**.
37. TARIQ M., HAMEED S., MALIK K.A., HAFEEZ F.Y. Plant root-associated bacteria for zinc mobilization in rice. *Pakistan Journal of Botany*, **39** (1), 245, **2007**.
38. JONES JR J.B., CASE V.W. Sampling, handling, and analyzing plant tissue samples. *Soil testing and plant analysis*, **3**, 389, **1990**.
39. STEEL R.G.D., TORRIE J.H., DICKEY D. *Principles and Procedures of Statistics: A Biometrical Approach*. 3rd Edn., McGraw Hill Book Co., Inc., New York, **1997**.
40. PRAJAPATI J., YADAV J., JAISWAL D.K., PRAJAPATI B., TIWARI S., YADAV J. Salt Tolerant Indigenous Zn Solubilizing Bacteria Isolated from Forest Organic Soils Promotes Yield and Root Growth in *Oryza Sativa* under Zinc Deficient Alluvial Soil. *Geomicrobiology Journal*, **39** (6), 465, **2022**.
41. HUSSAIN A., JIANG W., WANG X., SHAHID S., SABA N., AHMAD M., DAR A., MASOOD S.U., IMRAN M., MUSTAFA A. Mechanistic Impact of Zinc Deficiency in Human Development. *Frontiers in Nutrition*, **9**, 717064, **2022**.
42. BROADLEY M.R., WHITE P.J., ZELKO I., LUX A. Zinc in plants. *New Phytologist*, **173** (4), 677, **2007**.
43. ALI M., AHMED I., TARIQ H., ABBAS S., ZIA M.H., MUMTAZ A., SHARIF M. Growth improvement of wheat (*Triticum aestivum*) and zinc biofortification using potent zinc-solubilizing bacteria. *Frontiers in Plant Science*, **14**, 1140454, **2023**.
44. DITTA A., ULLAH N., IMTIAZ M., LI X., JAN A.U., MEHMOOD S., RIZWAN M.S., RIZWAN M. Zn biofortification in crops through Zn-solubilizing plant growth promoting rhizobacteria. In: MAHMOOD Q. (Ed). *Sustainable Plant Nutrition under Contaminated Environments*. Springer Nature Switzerland AG, **2022**. pp. 115.
45. LIU L., LI C., ZHU S., XU Y., LI H., ZHENG X., SHI R. Combined application of organic and inorganic nitrogen fertilizers affects soil prokaryotic communities compositions. *Agronomy*, **10** (1), e132, **2020**.
46. ALI A.M., AZEEM B., ALGHAMDI A.M., SHAHZAD K., AL-ZAHRANI A.A., RASHID M.I., MAHPUDZ A.B., JAMIL A. Optimization of Fluidized-Bed Process Parameters for Coating Uniformity and Nutrient-Release Characteristics of Controlled-Release Urea Produced by Modified Lignocellulosic Coating Material. *Agronomy*, **13** (3), 725, **2023**.
47. MIRBOLOOK A., SADAGHIANI M.R., KESHAVARZ P., ALIKHANI M. New Slow-Release Urea Fertilizer Fortified with Zinc for Improving Zinc Availability and Nitrogen Use Efficiency in Maize. *ACS omega*, **8** (48), 45715, **2023**.
48. HUSSAIN A., WANG X., ZAHIR Z.A., MAHMOOD K., MUMTAZ M.Z., SAQIB M., JAMSHAD M.U., AHMAD H.T. Potential of Integrated Use of *Bacillus* sp. AZ6 and Organic Waste for Zinc Bio-Activation to Improve Physiological Attributes of Maize. *Polish Journal of Environmental Studies*, **31** (2), **2022**.
49. PRASAD R., SHIVAY Y.S., KUMAR D. Zinc fertilization of cereals for increased production and alleviation of zinc malnutrition in India. *Agricultural Research*, **2**, 111, **2013**.
50. POONIYA V., SHIVAY Y.S., PAL M., BANSAL R. Relative performance of boron, sulfur and zinc coatings onto prilled urea for increasing productivity and nitrogen use efficiency in maize. *Experimental Agriculture*, **54** (4), 577, **2018**.
51. BARAL K., SHIVAY Y.S., PRASANNA R., KUMAR D., SRINIVASARAO C., MANDI S., NAYAK S., REDDY K.S. Enhancing physiological metrics, yield, zinc bioavailability, and economic viability of Basmati rice through nano zinc fertilization and summer green manuring in semi-arid South Asian ecosystem. *Frontiers in Plant Science*, **14**, **2023**.
52. MUMTAZ M.Z., AHMAD M., ZAFAR-UL-HYE M., SAQIB M., AKHTAR M.F., ZAHEER M.S. Seed-applied zinc-solubilising *Bacillus* biofertilizers improve antioxidant enzyme activities, crop productivity, and biofortification of maize. *Crop and Pasture Science*, **73** (5), 503, **2022**.
53. YASEEN M., AHMAD A., NAVEED M., ALI M.A., SHAH S.S.H., HASNAIN M., ALI H.M., SIDDIQUI M.H., SALEM M.Z. MUSTAFA A. Subsurface-applied coated nitrogen fertilizer enhanced wheat production by improving nutrient-use efficiency with less ammonia volatilization. *Agronomy*, **11** (12), e2396, **2021**.
54. DACCAK D., LIDON F.C., PESSOA C.C., LUÍS I.C., COELHO A.R.F., MARQUES A.C., RAMALHO J.C., SILVA M.J., RODRIGUES A.P., GUERRA M. LEITÃO R.G. Enrichment of grapes with Zinc-efficiency of foliar fertilization with ZnSO₄ and ZnO and implications on winemaking. *Plants*, **11** (11), e1399, **2022**.