

Original Research

Seedling Survivability and Change of Some Physiological Characters for Drought Resistance in Wheat

İsmet Başer¹, Seher Akseki Semerci², Damla Balaban Göçmen^{1*}

¹ Tekirdağ Namık Kemal University, Agricultural Faculty, Department of Field Crops, Tekirdağ, Turkey

² Tekirdağ Namık Kemal University, Graduate School of Natural and Applied Sciences, Tekirdağ, Turkey

Received: 22 February 2024

Accepted: 3 May 2024

Abstract

Detection of genotypic variation in response to water stress at the seedling stage provides important contributions to plant breeders in the rapid and effective selection of drought-resistant genotypes. The study was conducted in 2021 under in vivo conditions using soil samples taken from the trial area of Namık Kemal University Faculty of Agriculture Department of Field Crops and 45 genotypes, including 39 bread wheat, 1 rye, 1 Spella, 2 Kavlıca, and 2 Einkorn populations. Seedling survivability, stomata density, stomata width and length, seedling development score, canopy temperature, and chlorophyll content were examined to determine the early drought resistance in the experiment. When 45 genotypes in different maturing groups were examined for seedling survivability after drought application, the highest seedling survivability was found in Esperia, Rumeli, Krasunia O'deska, Almeria, Falado, and Rebelde varieties with 5 score values. Adelaide, President, Selimiye, Hakan, Quality, Hamza, LG 59, Golia, Siyez 1, and Siyez 2 genotypes showed the lowest seedling survivability. Esperia, Rumelia, Krasunia O'deska, Almeria, Falado, and Rebelde varieties, which have a high seedling survival rate, have low canopy temperature and high chlorophyll content. The data obtained show that early seedling survivability can be used in the selection of genotypes for drought resistance.

Keywords: bread wheat, chlorophyll content, drought, seedling, survival

Introduction

Wheat (*Triticum aestivum* L.), a major cereal crop, is constrained by drought in numerous regions of the world. Because of severe limitations imposed by drought, the development of cultivars with improved productivity under water stress is important for affected regions. However,

the development of drought-tolerant cultivars is hindered by a lack of understanding of the mechanisms of drought resistance and reliable selection techniques. Because wheat production in our country is mainly dependent on precipitation conditions, it can show significant fluctuations from year to year. In some years, especially with insufficient rainfall, wheat production decreases considerably. As a result, some consider drought as the main environmental stress for different plants, particularly in drought-prone areas [1], the single most critical threat to world food security in the future and the catalyst of important famines in the past

*e-mail: dgocmen@nku.edu.tr
Tel.: +90-545-208-41-67

[2]. Plant drought-stress tolerance, the ability of plants to maintain productivity during periods of drought stress, is a complex phenotype [3]. The decrease in productivity under drought stress conditions is one of the most important factors threatening global food production [4].

Plant drought-stress tolerance, the ability of plants to maintain productivity during periods of drought stress, is a complex phenotype [5]. Plants have numerous mechanisms to tolerate drought stress at the organ system to cellular levels, including adaptations in plant growth and development, resulting in alterations in plant allometry, root system development, and even time to flowering, osmotic adjustment, and optimization of water use [6]. The morphological and physiological responses of crops to drought can explain the large variation in yield under drought stress conditions [7].

Drought stress causes a number of biochemical, morphophysiological, and anatomical disturbances in plants that occur at all stages of plant development, from seed germination to the final stages of the generative phase [8]. Drought tolerance in plants is the result of numerous inter-related mechanisms that operate at the molecular, cellular, and organismal levels. Among them are changes in the state of the stomatal apparatus, and accumulation of various osmolytes in cells [9, 10]. The maximum quantum yield of primary photochemistry in leaves in the dark-adapted state can reflect wheat's light-use efficiency under drought conditions [11]. The relatively low canopy temperature value at the grain-filling stage and the high chlorophyll content during the late grain-filling stage can be used to screen the winter wheat cultivars adapted to dryland ecosystems [12].

Wheat is very sensitive to drought during the early seedling stages and germination. Insufficient available water in the early development stages significantly constrains grain yield through decreasing germination, emergence, and seedling development and establishment [13]. Seedlings of plants are extremely sensitive to drought stress conditions [14]. Drought stress affects water balance, disrupts metabolic reactions at the cellular level, reduces ATP synthesis and respiration, and results in poor seed germination [15].

The aim of the study is to reveal the effectiveness of the survival ratio, canopy temperature, chlorophyll content, and stomatal characteristics of the seedlings after experiencing drought conditions in the differentiation of drought-resistant genotypes of 45 genotypes in different seedling mortality ratios.

Material and Methods

In the study, 45 genotypes of different origins, including 39 bread wheat, 1 rye, 1 Spelta, 2 Kavlıca, and 2 Siyez populations, were used as materials. The genotypes used as material in the experiment are given in Table 1.

The experiment was performed in a covered crate environment in order to determine the early drought resistance levels of 45 genotypes used as material in the experiment. Seedling survival level, seedling development score, canopy temperature, chlorophyll content, stomata density, stomata width, and length were examined.

In order to determine the seedling survival level in the experiment, a 10-meter-long and 1.2-meter-wide crate was constructed (Fig. 1).

Table 1. Genotypes used as material in the experiment

Genotypes	Institution	Growth Habit
Golia	Tareks Agricultural Credit Seed Company	Alternative
Aglika	Tarar Flour and Food Ind. Trade. Co. Ltd.	Winter
Anopa	2A Seed Company	Alternative
Rebelde	Marmara Seed Development Inc.	Winter
Genesi	Tasaco Agroindustry and Trading Inc.	Winter
Kaan	Trakya Agriculture and Veterinary Trade. Co. Ltd.	Winter
Quality	Ata Seed Company	Alternative
Refikbey	Akçakaya Agriculture Ind. Trade. Co. Ltd.	Winter
Rumeli	Trakya Agriculture and Veterinary Trade. Co. Ltd.	Winter
Sarı Mustafa	Sarı Seed Company	Winter
Selimiye	Trakya Agriculture Research Institute	Winter
TT 601	Trakya Agriculture and Veterinary Trade. Co. Ltd.	Winter
Adelaide	Maro Agriculture Building Trade Ind. Inc.	Alternative
Aldane	Trakya Agricultural Research Institute	Winter
Almeria	Alfa Seed Agriculture Marketing Ind. Trade. Co. Ltd.	Alternative



Table 1 cont. Genotypes used as material in the experiment

Genotypes	Institution	Growth Habit
Ambrosia	Pro Gen Seed Company	Alternative
Bc Anica	BC İstitut Agricul. Products Auto Ind. Trade. Co. Ltd.	Alternative
Başkan	Sarı Seed Company	Alternative
Bezostaja 1	Corn Research Institute	Winter
Bora	Tasaco Agroindustry and Trading Inc.	Winter
Energo	ITU Agricultural Practices Co. Ltd.	Winter
Enola	Tarar Flour and Food Ind. Trade. Co. Ltd.	Winter
Esperia	Tasaco Agroindustry and Trading Inc.	Winter
Falado	Syngenta Agroindustry and Trading Inc.	Alternative
Hakan	Trakya Agriculture and Veterinary Trade. Co. Ltd.	Winter
Hamza	Tekcan Seed and Agric. Products Ind. Trade. Co. Ltd.	Alternative
İveta	Tarar Flour and Food Ind. Trade. Co. Ltd.	Winter
Krasunia Odes'ka	Yıldız Agr. Products Seed and Agriculture Ind. Inc.	Winter
LG 59	Limagrain Seed Impr. and Production Ind. Trade. Inc.	Alternative
Maden	Trakya Agriculture and Veterinary Trade. Co. Ltd.	Winter
Masaccio	Pro Gen Seed Company	Alternative
Maya	Trakya Agriculture and Veterinary Trade. Co. Ltd.	Winter
Mihelca	BC İstitut Agricul. Products Auto Ind. Trade. Co. Ltd.	Winter
Misiia Odes'ka	Yıldız Agric. Products Seed and Agriculture Ind. Inc.	Winter
NKU Asiya	Tekirdag Namık Kemal University Faculty of Agri.	Winter
NKU Ergene	Tekirdag Namık Kemal University Faculty of Agri.	Winter
NKU Lider	Tekirdag Namık Kemal University Faculty of Agri.	Winter
Pannonia	Marmara Seed Company	Winter
Prima	BC İstitut Agric. Products Auto Ind. Trade. Co. Ltd.	Winter
Kavlıca white		Population
Kavlıca color		Population
Siyez-1		Population
Siyez-2		Population
Spelta		Population
Ducato		Winter

Soil samples taken from Tekirdag Namık Kemal University, Faculty of Agriculture, Department of Field Crops trial area were filled into the crate with a depth of 20 cm and water was given until the soil was saturated. According to the soil analysis taken from the trial area, the soil has a pH close to neutral, a loamy soil structure, and low organic matter (1.03%). When the soil was at the right level, each genotype was sown with a length of 1 meter and 1-row spacing (26.05.2021), and 500 seeds were used per square meter. Irrigation was applied until the plants had 3 leaves (07.06.2021), and then no irrigation was applied

until it appeared that more than 90% of the seedlings died due to lack of water (27.07.2021).

Plants were subjected to 50 days of drought stress, and when it appeared that most of the plants died due to the lack of water (27.07.2021), irrigation water was given to the crates. Irrigation was repeated at regular intervals and seedling survival ratios were determined again (22.08.2021). Seedling survival after drought was calculated as a percentage [16]. The plants were given a score of 1 for no plant development, 2 for plant development between 1–5%, 3 for plant development between 5.1–10.0%, 4 for



Fig. 1. The seedling development in wheat genotypes



Fig. 2. Drought application in seedling stage and seedling survivability

plant development between 10.1–15.0%, and 5 for plant development exceeding 15% (Fig. 2).

The canopy temperature, chlorophyll content, stomata density, width, and length characteristics of the plants developed at the beginning of the formation of stress images were determined. The chlorophyll content of flag leaves on the main stem of plants was measured with a “Konica Minolta SPAD-502” portable chlorophyll meter. The canopy temperature was determined with an infrared thermometer. Analysis of variance was performed on the data obtained and the Tukey ($P < 0.01$) test was used to determine the differences between genotypes.

Results and Discussion

The seedling development scores of 45 genotypes sown in the rows in the crate environment are given in Table 2.

As seen in the table, seedling development scores varied between 1 and 4. The scores of Adelaide, Başkan, Refikbey, Spelta, Kavlıca color, and Siyez 2 genotypes were the lowest with a score of 1. In terms of seedling development, 15 genotypes in the experiment had a score of 2 while 16 genotypes had a score of 3. Among the genotypes included in the experiment, Maden, Enola, Mihelca, Maya, Ambrosia, Hamza, Aglika, Anapo and LG 59 had a score of 4 for seedling development. According to the data, it was determined that varieties in the early and mid-ripening group, such as Hamza, Aglika, Anopa, and LG 59, had higher seedling development scores [17].

When the survival ratios of the seedlings were evaluated again after the drought in the genotypes included in the experiment, no seedling survivability was observed again in the Adelaide, Başkan, Selimiye, Hakan, Quality, Hamza, LG 59, Golia, Siyez 1, and Siyez 2 genotypes. TT 601, NKU Lider, NKU Ergene, İveta, Maya, Ambrosia, and Kavlıca

Table 2. Seedling development score in the genotypes

Genotypes	Score	Genotypes	Score	Genotypes	Score
Adelaide	1	Golia	2	NKU Ergene	3
Başkan	1	Kaan	2	Hakan	3
Refikbey	1	Kavlıca white	2	Quality	3
Spelta	1	Genesi	2	Masaccio	3
Kavlıca color	1	Rebelde	2	Misiia Odes'ka	3
Siyez-2	1	Ducato	2	Siyez-1	3
Esperia	2	NKU Asiya	3	Maden	4
Selimiye	2	Rumeli	3	Enola	4
Falado	2	Krasunia Odes'ka	3	Mihelca	4
NKU Lider	2	Almeria	3	Maya	4
İveta	2	Bezostaja 1	3	Ambrosia	4
Bora	2	Aldane	3	Hamza	4
Pannonia	2	Prima	3	Aglika	4
Energo	2	Anica	3	Anopa	4
Sarı Mustafa	2	TT 601	3	LG 59	4

white genotypes had a score of 2 with seedling survivability between 1–5%. Prima, Anica, Mihelca, Bora, Pannonia, Masaccio, Sarı Mustafa, Refikbey, Spelta, and Kavlıca color had a score of 3 with seedling survivability between 5.1–10.0%. NKU Asiya, Maden, Enola, Bezostaja 1, Energo, Misiia Odes'ka, Aglika, Kaan, Genesi, and Ducato genotypes had a score of 4 with seedling survivability between 10.1–15.0%. Esperia, Rumeli, Krasunia Odes'ka,

Almeria, Falado, and Rebelde had a score of 5 with seedling survivability ratio of more than 15% (Table 3). The morphological and physiological responses of crops to drought can explain the large variation in yield under drought stress conditions [18]. In the study, seedling weight, root number, root length, root weight, shoot length, and shoot weight characters were determined. It was determined that drought stress applications caused statistically significant decreases

Table 3. Seedling survivability score in the genotypes

Genotypes	Score	Genotypes	Score	Genotypes	Score
Adelaide	1	İveta	2	Maden	4
Başkan	1	Maya	2	Enola	4
Selimiye	1	Ambrosia	2	Bezostaja 1	4
Hakan	1	Kavlıca white	2	Energo	4
Quality	1	Prima	3	Misiia Odes'ka	4
Hamza	1	Anica	3	Aglika	4
Anopa	1	Mihelca	3	Kaan	4
LG 59	1	Bora	3	Genesi	4
Golia	1	Pannonia	3	Ducato	4
Siyez-1	1	Masaccio	3	Esperia	5
Siyez-2	1	Sarı Mustafa	3	Rumeli	5
Aldane	2	Refikbey	3	Krasunia Odes'ka	5
TT 601	2	Spelta	3	Almeria	5
NKU Lider	2	Kavlıca color	3	Falado	5
NKU Ergene	2	NKU Asiya	4	Rebelde	5

in root and shoot characters. Osmotic stress of 1.00 MPa and 0.75 MPa caused statistically significant reductions in root and shoot characters [19].

None of the genotypes with the lowest and highest seedling development scores were among the genotypes showing the highest seedling survivability (Esperia 2, Rumeli 3, Krasunia Odes'ka 3, Almeria 3, Falado 2, and Rebelde 2). With the obtained data, it was concluded that there was no significant relationship between seedling growth score and seedling survivability ratios. A genotype with drought stress tolerance capabilities has a more developed root structure to reduce the pronounced effects of drought by using soil moisture during development and growth. Various methods were investigated to determine wheat genotypes that are drought-resistant

in the seedling period. Selection of wheat varieties on the basis of seedling traits is easy, inexpensive, and less laborious. Similarly, seedling attributes expose moderate-to-high variability with an additive gene effect across environments [20]. It was determined that effective results were obtained in terms of drought resistance by using seedling characteristics and that the material determined for drought resistance in the seedling period could be successfully used in breeding studies [21].

In the analysis of variance performed on canopy temperature, chlorophyll content, and stomatal characteristics of 45 genotypes used in the study, the differences between genotypes were found to be statistically significant and the results of the significance test performed to reveal the differences between genotypes are given in Table 4.

Table 4. Significance groups in canopy temperature, chlorophyll content and stomatal characteristics

Genotypes	Canopy temperature (°C)	Chlorophyll content (SPAD)	Stomata density	Stomata length (μ)	Stomata width (μ)
Başkan	38.67 ^a	31.40 ^{cde}	4.6 ^{gh}	24.6 ^{e-i}	9.4 ^{d-i}
Adelaide	36.46 ^{ab}	33.08 ^{bc-e}	8.50 ^{a-e}	25.4 ^{e-h}	11.0 ^{e-f}
Golia	33.67 ^{abc}	32.90 ^{bc-e}	9.67 ^{abc}	25.4 ^{e-h}	10.6 ^{c-h}
Siyez-1	33.67 ^{abc}	29.43 ^e	5.67 ^{e-1}	18.6 ^{klm}	8.4 ^{fj}
Spelta	32.67 ^{abc}	35.43 ^{bc-e}	4.00 ^l	32.0 ^{ab}	9.4 ^{d-i}
NKU Asiya	31.67 ^{abc}	33.83 ^{bc-e}	6.33 ^{d-1}	32.0 ^{ab}	9.4 ^{d-i}
Kavlıca white	31.33 ^{abc}	37.87 ^{bc-e}	5.67 ^{e-1}	16.6 ^{lm}	8.0 ^{g-i}
Kavlıca color	31.33 ^{abc}	33.83 ^{bc-e}	9.33 ^{a-d}	16.4 ^{lm}	8.0 ^{g-j}
Siyez-2	31.00 ^{abc}	41.50 ^{ab}	10.00 ^{ab}	14.0 ^m	6.6 ^{ij}
Genesi	30.67 ^{abc}	40.30 ^{a-d}	5.67 ^{e-1}	24.6 ^{e-i}	11.4 ^{b-f}
Anica	30.33 ^{abc}	36.73 ^{bc-e}	4.33 ^{hi}	28.0 ^{b-g}	8.6 ^{e-j}
Kaan	30.33 ^{abc}	40.10 ^{a-d}	6.33 ^{d-1}	20.0 ^{h-1}	12.0 ^{a-d}
Pannonia	30.33 ^{abc}	36.90 ^{bc-e}	6.33 ^{d-1}	28.6 ^{b-f}	9.4 ^{d-i}
Masaccio	29.67 ^{abc}	33.87 ^{bc-e}	7.33 ^{b-h}	20.6 ^{h-1}	11.6 ^{a-e}
Energo	29.33 ^{abc}	37.57 ^{bc-e}	5.00 ^{f-1}	23.4 ^{g-k}	11.0 ^{e-g}
Refikbey	29.33 ^{abc}	37.03 ^{bc-e}	6.67 ^{e-1}	29.4 ^{b-e}	11.6 ^{a-e}
Quality	29.00 ^{abc}	33.93 ^{bc-e}	6.00 ^{e-1}	23.6 ^{f-k}	12.0 ^{a-d}
Ambrosia	28.67 ^{abc}	38.63 ^{bc-e}	6.00 ^{e-1}	23.4 ^{g-k}	10.4 ^{c-h}
NKU Lider	28.67 ^{abc}	34.27 ^{bc-e}	7.67 ^{a-g}	25.6 ^{d-h}	14.6 ^a
Sarı Mustafa	28.67 ^{abc}	38.60 ^{bc-e}	9.67 ^{abc}	16.6 ^{lm}	9.0 ^{d-j}
Bezostaja 1	28.33 ^{bc}	31.63 ^{cde}	5.6 ^{e-1}	19.4 ^{jkl}	8.4 ^{fj}
Mihelca	28.09 ^{bc}	41.24 ^{abc}	4.51 ^{gh}	28.2 ^{b-g}	9.0 ^{d-j}
Krasun. Odes'ka	28.00 ^{bc}	36.33 ^{bc-e}	8.67 ^{a-e}	26.6 ^{c-g}	12.6 ^{abc}
LG 59	27.67 ^{bc}	41.03 ^{a-d}	6.67 ^{e-1}	24.0 ^{f-j}	9.4 ^{d-i}
Bora	27.33 ^{bc}	37.13 ^{bc-e}	3.67 ^l	29.4 ^{b-e}	9.4 ^{d-i}
Aglıka	27.00 ^c	41.60 ^{ab}	6.33 ^{d-1}	27.6 ^{b-g}	8.6 ^{e-j}



Table 4 cont. Significance groups in canopy temperature, chlorophyll content and stomatal characteristics

Genotypes	Canopy temperature (°C)	Chlorophyll content (SPAD)	Stomata density	Stomata length (μ)	Stomata width (μ)
Misiia Odes'ka	27.00 ^c	49.03 ^a	9.67 ^{abc}	24.0 ^{fj}	8.6 ^{e-j}
Selimiye	27.00 ^c	29.53 ^e	6.00 ^{e-1}	21.4 ^{h-1}	12.0 ^{a-d}
Anopa	26.67 ^c	37.77 ^{b-e}	10.00 ^{ab}	28.0 ^{b-g}	8.6 ^{e-j}
Hakan	26.67 ^c	37.00 ^{b-e}	4.33 ^{hi}	26.6 ^{c-g}	9.4 ^{d-i}
Prima	26.67 ^c	29.63 ^e	4.00 ⁱ	25.4 ^{e-h}	11.4 ^{b-f}
Esperia	26.33 ^c	35.90 ^{b-e}	5.67 ^{e-1}	36.0 ^a	9.6 ^{e-i}
Maden	26.00 ^c	40.73 ^{a-d}	11.00 ^a	30.63 ^{bcd}	11.4 ^{b-f}
NKU Ergene	26.00 ^c	38.37 ^{b-e}	7.67 ^{a-g}	20.6 ^{h-1}	6.0 ^j
Rumeli	26.00 ^c	37.63 ^{b-e}	7.33 ^{b-h}	24.6 ^{e-i}	11.6 ^{a-e}
Almeria	25.33 ^c	31.20 ^{cde}	5.67 ^{e-1}	20.0 ^{h-1}	8.6 ^{e-j}
Rebelde	25.33 ^c	37.67 ^{b-e}	6.33 ^{d-1}	28.6 ^{b-f}	12.0 ^{a-d}
Maya	25.00 ^c	36.33 ^{b-e}	8.00 ^{a-f}	24.0 ^{fj}	12.6 ^{abc}
TT 601	25.00 ^c	34.20 ^{b-e}	7.33 ^{b-h}	25.6 ^{d-h}	9.4 ^{d-i}
Hamza	24.67 ^c	37.63 ^{b-e}	4.00 ⁱ	24.4 ^{e-j}	11.6 ^{a-e}
İveta	24.67 ^c	37.90 ^{b-e}	5.67 ^{e-1}	21.4 ^{h-1}	9.6 ^{e-i}
Aldane	24.33 ^c	30.20 ^e	8.00 ^{a-f}	31.0 ^{abc}	9.6 ^{e-i}
Ducato	24.33 ^c	36.87 ^{b-e}	5.67 ^{e-1}	19.4 ^{ijkl}	8.6 ^{e-j}
Enola	24.00 ^c	35.03 ^{b-e}	8.00 ^{a-f}	27.6 ^{b-g}	7.6 ^{hij}
Falado	23.67 ^c	30.97 ^{de}	6.67 ^{c-1}	24.6 ^{e-i}	14.4 ^{ab}

The identical letters indicate statistical groups of identical values with a $P < 0.01$ confidence level by the Tukey Test.

According to the average values, the canopy temperature varied between 38.67–23.67°C in 45 genotypes. Among the genotypes in the study, the lowest canopy temperature was in Falado cultivar with 23.67°C, followed by Ducato and Aldane with 24.33°C, Iveta with 24.67°C, TT 601 and Maya with 25.00°C, and Almeria and Rebelde with 25.33°C. The highest canopy temperature was 38.67°C in Başkan cultivar, followed by the genotypes such as Adelaide with 36.45°C, Golia with 33.67°C, Siyez 1 with 33.67°C, Spelta with 32.67°C, NKU Asiya with 31.67°C, Kavlıca white and Kavlıca color with 31.33°C, Siyez 2 with 31.00°C.

When the genotypes were analyzed for chlorophyll content, it varied between 49.03–29.43. The highest chlorophyll content was found in Misiia Odes'ka at 49.03, followed by Aglika, Siyez 2, Mihelca and LG 59 at 41.60, 41.50, 41.24 and 41.03 values. The lowest chlorophyll content was found in Siyez 1 population at 29.43, followed by Selimiye with 29.53 and Prima at 29.63. When the genotypes were analyzed for stomatal characteristics, the stomata density varied between 4.00 and 11.00, stomata width between 6.0 and 14.6 μ and stomata length between 14.0 and 36.0 μ. In crops, recent studies are showing that engineering plants to reduce stomatal number may be an effective tool to improve plant WUE and drought tolerance without yield reductions

[22]. While the highest stomata density was 11.00 in Maden cultivar, it was followed by Siyez 1 and Anapo varieties at 10.00. In terms of the stomata density, Misiia Odes'ka and Sarı Mustafa at 9.67 and Kavlıca at 9.33 were ranked next. The lowest stomata density was 4.0 in Prima, 4.33 in Hakan, Hamza, Spelta and Anica cultivars. The lowest stomata density was in Prima cultivar at 4.0, and Hakan, Hamza, Spelta and Anica cultivars at 4.33. Plants' first physiological response to water scarcity is stomatal closure, which reduces photosynthetic activity by slowing the rate of carbon dioxide entry into mesophyll cells [23].

The highest value for stomata length was found in the Esperia cultivar with 36.0 μ, followed by Spelta and NKU Asiya with 32.0 μ. The least stomata length was found in the Siyez 2 genotype with 14.0 μ, followed by Kavlıca color, Kavlıca white and Sarı Mustafa genotypes with 16.4, and 16.6 μ. The highest value for stomata width was found in the NKU Lider cultivar with 14.6 μ, followed by Falado with 14.4 μ, Maya, Krasunia Odes'ka, Selimiye and Rebelde cultivars with 12.6 μ. The lowest stomata width was found in the NKU Ergene cultivar with 6.0 μ, followed by Siyez 2 population with 6.6 μ and Enola genotypes with 7.6 μ.

The reduction in the length and weight of the hypocotyl in seedlings due to water stress can be mitigated with

the application of gibberellic acid, which helps to maintain the internal water balance and the protein synthesis in drought stressed plants [24]. Hence, certain genotypes and cultivars, which are drought susceptible and unable to adjust to environmental conditions, resulted in low water-use efficiency [25].

Conclusions

When 45 genotypes in different maturing groups were examined for seedling survivability after drought application, 11 genotypes, including Adelaide, Başkan, Selimiye, Hakan, Quality, Hamza, LG 59, Golia, Siyez 1, and Siyez 2, did not show seedling survivability, while Esperia, Rumeli, Krasunia Odes'ka, Almeria, Falado, and Rebelde were the genotypes that showed the highest seedling survivability with a score of 5. Genotypes with low canopy temperature and high chlorophyll content showed higher seedling survivability. The data obtained show that early seedling survivability can be used in the selection of genotypes for drought resistance.

Conflict of Interest

“The authors declare no conflict of interest”.

References

- DIATTA A.A., FIKE J.H., BATTAGLIA M.L., GALBRAITH J., BAIG M.B. Effects of biochar on soil fertility and crop productivity in arid regions: A review. *Arabian Journal of Geosciences*, **13**, 595, **2020**.
- OKORIE V.O., MPHAMBUKELI T.N., AMUSAN S.O. Exploring the political economy of water and food security nexus in BRICS. *Africa Insight*, **48** (4), 21, **2019**.
- NGUMBI E., KLOPPER J. Bacterial-mediated drought tolerance: current and future prospects. *Applied Soil Ecology*, **105**, 109, **2016**.
- FAHAD S., BAJWA A.A., NAZİR U., ANJUM S.A., FAROOQ A., ZOHAİB A., SADİA S., NAŚİM W., ADKİNS S., SAUD S., İHSAN M.Z., ALHARBY H., WU C., WANG D., HUANG J. Crop production under drought and heat stress: plant responses and management options. *Frontiers in Plant Science*, **29** (8), 1147, **2017**.
- NGUMBI E., KLOPPER J.W. Bacterial-mediated drought tolerance: current and future prospects. *Applied Soil Ecology*, **105**, 109, **2016**.
- MEENA K.K., SORTY A.M., BİTLA U.M., CHOUDHARY K., GUPTA P., PAREEK A., SİNGH D.P., PRABHA R., SAHU P.K., GUPTA V.K., SİNGH H.B., KRİSHANANİ K.K., MİNHAS P.S. Abiotic stress responses and microbe-mediated mitigation in plants: the omics strategies. *Frontiers Plant Science*, **9** (8), 172, **2017**.
- PUANGBUT D., JOGLOY S., VORASOOT N., SONGSRİ P. Photosynthetic and physiological responses to drought of Jerusalem artichoke genotypes differing in drought resistance. *Agricultural Water Management*, **259** (2), 107252, **2022**.
- TİWARİ R.K., LAL M.K., KUMAR R., CHOURASİA K.N., NAGA K.C., KUMAR D., DAS, S.K., ZİNTA G. Mechanistic insights on melatonin-mediated drought stress mitigation in plants. *Physiologia Plantarum*, **172** (2), 1212, **2021**.
- SİNGH S., PRAKASH P., SİNGH A.K. Salicylic acid and hydrogen peroxide improve antioxidant response and compatible osmolytes in wheat (*Triticum aestivum* L.) under water deficit. *Agricultural Research*, **10** (2), 175, **2021**.
- KOLUPAEV Y.E., YASTREB T.O., RYABCHUN N.I., KOKOREV A.I., KOLOMATSKA V.P., DMİTRİEV A.P. Redox homeostasis of cereals during acclimation to drought. *Theoretical and Experimental Plant Physiology*, **35** (2), 133, **2023**.
- YU S., ZHANG N., KAİSER E., Lİ G., AN D., SUN Q., CHEN W., LİU W., LUO W. Integrating chlorophyll fluorescence parameters into a crop model improves growth prediction under severe drought. *Agricultural and Forest Meteorology*, **303**, 108367, **2021**.
- LU Y., YAN Z., Lİ L., GAO C., SHAO L. Selecting traits to improve the yield and water use efficiency of winter wheat under limited water supply. *Agricultural Water Management*, **242**, 106410, **2020**.
- RICHARDS R.A., LUKACS Z. Seedling vigour in wheat sources of variation for genetic and agronomic improvement. *Australian Journal of Agricultural Research*, **53** (1), 41, **2002**.
- MAHPARA S., ZAİNAB A., ULLAH R., KAUSAR S., BİLAL M., LATİF M.I., ARİF M., AKHTAR I., ALHASHİMİ A., ELSHİKH M.S., ZİVCAK M., ZUAN A.T.K. The impact of PEG-induced drought stress on seed germination and seedling growth of different bread wheat (*Triticum aestivum* L.) genotypes. *Plos. One*, **17** (2), 29, **2022**.
- UPADHYAYA N.M., MAGO R., PANWAR V., HEWITT T., LUO M., CHEN J., SPERSCHNEIDER J., NGUYENPHUC H., WANG A., ORTIZ D., HAC L., BHATT D., LI F., ZHANG J., AYLİFFE M., FIGUEROA M., KANYUKA K., ELLIS J.G., DODDS P.N. Genomics accelerated isolation of a new stem rust a virulence gene-wheat resistance gene pair. *Nature Plants*, **7**, 1220, **2021**.
- WINTER S.R., MUSICK J.T., PORTER K.B. Evaluation of creening techniques for breeding drought-resistant winter wheat. *Crop Science*, **28** (3), 512, **1988**.
- YILDIZ C.F. Determination of yield and quality parameters of some bread wheat varieties in Southern Marmara ecological conditions. Bursa Uludağ University, Institute of Science, Field Crops Department, Master's Thesis, **2023**.
- PUANGBUT D., JOGLOY S., VORASOOT N., SONGSRİ P. Photosynthetic and physiological responses to drought of Jerusalem artichoke genotypes differing in drought resistance. *Agricultural Water Management*, **259**, 107252, **2022**.
- BAŞER İ., SEMERCİ S.A., GÖÇMEN D.B., BİLGİN O., BALKAN A. Variability for drought stress effects on seedling growth in bread wheat (*Triticum aestivum* L.) genotypes. *Ekin Journal of Crop Breeding and Genetics*, **9** (1), 24, **2023**.
- AHMED H., KHAN A.S., KHAN S.H., KASHİF M. Genome wide allelic pattern and genetic diversity of spring wheat genotypes through SSR markers. *International Journal of Agriculture and Biology*, **19** (6), 1559, **2017**.
- FAROOQ M.O., KASHİF M. Accessing potential of seedling traits for screening of wheat genotypes under drought conditions. *Agrobiological Records*, **6**, 1, **2021**.
- BERTOLINO L.T., CAINE R.S., GRAY J.F. Impact of stomatal density and morphology on water-use efficiency

- in a changing world. *Frontiers Plant Sciences*, **10**, 225, **2019**.
23. GOMEZ-CANDON D., MATHIEU V., MARTÍNEZ S., LABBE S., DELALANDE M., REGNARD J.L. Unravelling the responses of different apple varieties to water constraints by continuous field thermal monitoring. *Scientia Horticulturae*, **299**, 111013, **2022**.
24. TARDIEU F., SIMONNEAU T., MULLER B. The physiological basis of drought tolerance in crop plants: A scenario-dependent probabilistic approach. *Annual Review of Plant Biology*, **69**, 733, **2018**.
25. DING Z., ALI E.F., ELMAHDY A.M., RAGAB K.E., SELEIMAN M.F., KHEIR A.M.S. Modelling the combined impacts of deficit irrigation, rising temperature and compost application on wheat yield and water productivity. *Agricultural Water Management*, **244**, 106626, **2021**.