

Original Research

Phytochemical Screening of Antioxidant and Antibacterial Activities of Marine Algae Extracts

Djazia Kherraz-Chemlal*, Fatima Khelil, Meriem Mazouzi, Amaria Matallah-Boutiba,
Mohamed Bouderbala

Environmental Monitoring Network, Department of Biology, Faculty of Science,
University Ahmed Ben Bella Oran 1, Algeria

Received: 21 January 2024

Accepted: 08 May 2024

Abstract

This work consists of the evaluation of the phytochemical compositions and the study of the antioxidant and antimicrobial activities of three algae (green algae: *Ulva lactuca*, brown algae: *Cystoseira amentacea* var. *stricta*, and red algae: *Corallina elongata*) collected from the Algerian west coast.

For the extraction we used the maceration method and the antioxidant activity was determined using the scavenging activity of DPPH free radical and ferric-reducing antioxidant power assay for the antimicrobial activity of the extracts of the three algae, we used the diffusion method. The results of the dosages of polyphenols and flavonoids indicate the richness of the species *C. elongata* in polyphenols with a value of (219.43 mg EAG/g) and of *Ulva lactuca* in flavonoids with (65.17 mg EQ/g). As for the measurement of anti-radical activity, we note that the best antioxidant activity is that of the red algae extract *C. elongata* with an IC50 value of 275.70 µg/ml. For the antibacterial activity, the results obtained show that the crude extracts of *Ulva* and *Cystoseira* act in the same way on *Bacillus cereus*, *Bacillus subtilis*, and Methicillin-resistant *Staphylococcus aureus* (SARM) with diameters of (8 mm), (10 mm) and (22 mm) respectively for *Ulva* and (20 mm), (14 mm) and (21 mm) respectively for *Cystoseira*. As for *Corallina* extract, the inhibition zones vary from (16 mm) against *B. subtilis*, to 8mm for *Klebsiella* and *Staphylococcus*, passing through an inhibition zone of (12 mm) for *E. coli*.

Keywords: *Cystoseira amentacea*, *Ulva lactuca*, *Corallina elongata*, polyphenols, flavonoids

Introduction

The marine environment is an ecosystem made unique because of the diversity of the organisms it supports

[1]. From estuaries to abyssal trenches to hydrothermal vents, algae demonstrate incredible ecological plasticity. The latter constitutes an enormous reservoir of potentially active natural molecules [2, 3]. Studies have made it possible to isolate and identify a very large number of new molecules of great structural originality, many of which have interesting biological activity. Furthermore, the plant origin of the active ingredients isolated from algae

*e-mail: kherraz_djazia@yahoo.fr

constitutes a real and important advantage at a time when ingredients of animal origin are irrelevant [4]. Faced with the appearance of resistant forms of several bacteria to certain antibiotics, the search for new active molecules has become a necessity [5]. In addition, natural antioxidants and antimicrobial agents may provide solutions to the public's health issues because they can constitute a safer and more effective alternative to synthetic products whose toxicity is proven [6]. Likewise, these bioactive compounds are of great interest in the medical field, which is in an ongoing search for more powerful antibiotics to combat the antibiotic resistance of pathogens [7, 8].

Algeria is a country with a maritime coastline stretching over 1622 km. It constitutes a rich source of marine algae, which, however, remains little studied and is not yet exploited by pharmaceutical or cosmetic companies, while algae constitute a considerable economic development issue. The existing work on the algal flora of Algeria is essentially of the floristic inventory type. This is why our study aims at another aspect of greater interest and of growing interest which is the composition of polyphenols and flavonoids as well as the study of the antioxidant and antimicrobial potential of marine algae, in particular in *Ulva lactuca*, *Cystoseira amentacea* var. *stricta*, and *Corallina elongata*.

Materials and Methods

Harvesting and Processing Algae

The harvest of algae was carried out on the Algerian West Coast. The algae were identified by specialists in the field using the standard identification keys. Algal specimens were collected by hand and stored in plastic bins, filled with seawater, until processed the same day in the laboratory. After removing debris, small crustaceans, and other algae, the thalli are rinsed with distilled water several times to clean the epiphytes completely. These algae are dried in the open air and away from light for 7 to 10 days until complete dehydration, then ground into a very fine powder, then sifted and stored in a dark and dry place [9].

Extraction

The extraction method adopted was maceration by mixing two organic solvents. 50 g of each powdered algae is added to 400 ml of two solvents: 200 ml of methanol and 200 ml of toluene. The first is polar, and the second is non-polar. The resulting mixture was stirred continuously for 48 hours in the absence of light at room temperature, then filtered with Wattman N°1 paper. The filtrate obtained was evaporated under reduced pressure at a temperature of 40°C using a "Büchi" type rotary evaporator [10].

Each residue was weighed, and the extract recovered with a few drops of methanol was stored in sterile pill bottles at 4°C until its use for the different dosages [11].

Determination of Yield

The determination of the yield of the dry extract is performed by applying the following formula:

$$R (\%) = (P1 - P2/PE) \times 100$$

knowing that:

P1: Weight of the balloon after evaporation.

P2: Weight of the empty flask before evaporation.

PE: Weight of dry algal material.

Dosage of Flavonoids

A volume of 1 ml of the extract is added to 1 ml of 2% aluminum trichloride. The mixture was shaken and incubated for 30 min at room temperature in the dark, then the absorbance was measured at 430 nm on a spectrophotometer, with quercetin used as a reference standard. The results are expressed in micrograms of quercetin equivalent per milligram of dry extract ($\mu\text{g EQ/mg extract}$) [12].

Dosage of Polyphenols

Add 200 μl of the extract in a tube to the mixture (1 ml of Folin-Ciocalteu reagent diluted 10 times and 0.8 ml of 7.5% sodium carbonate). Shake the tube and store at room temperature for 30 minutes. Use a spectrophotometer to measure the absorbance at 765 nm against a blank. Use gallic acid as a positive control and run the calibration curve in parallel under the same operating conditions. Results are expressed in milligrams (mg) of gallic acid equivalent per gram of dry algae (mg EAG/g DM).

Dosage of Chlorophyll a, b, and Total Carotenoids

The determination of chlorophyll and carotenoids was carried out according to the spectral method [13, 14]. It uses characteristic wavelengths which correspond to the maximum absorbances of chlorophylls and carotenoids. 3 g of crushed algae were added to 30 ml of methanol. After adding 300 mg of magnesium sulfate, the mixture was incubated for 24 hours at 4°C. The mixture obtained was filtered and then centrifuged at 5000 rpm for 10 minutes. The supernatant was collected and the absorbances were read at 666, 653, and 470 nm for chlorophylls a, b, and total carotenoids, respectively. The chlorophyll and carotenoid contents were calculated in $\mu\text{g/ml}$ according to the following formulas:

$$Ca (\mu\text{g/ml}) = 15.65 A_{666} - 7.340 A_{653}$$

$$Cb (\mu\text{g/ml}) = 27.05 A_{653} - 11.21 A_{666}$$

$$Cc (\mu\text{g/ml}) = 1000 A_{470} - 2.860 Ca - 129.2 Cb/245$$

Evaluation of Antioxidant Activity

Trapping of the Free Radical DPPH (2,2-Diphenyl-1-Picrylhydrazyl)

This method is based on measuring the ability of antioxidants to trap the DPPH radical [13]. A volume of 50 µl of different concentrations of each extract is added to 1950 µl of the freshly prepared methanolic solution of DPPH (0.025 g/l). Concerning the negative control, the latter is prepared in parallel by mixing 50 µl of methanol with 1950 µl of a methanolic solution of DPPH at the same concentration used. The positive control is represented by a methanolic solution of a standard antioxidant (Ascorbic acid). After incubation in the dark for 30 min and at room temperature, the absorbance reading is carried out at 515 nm using a spectrophotometer against the white, according to the formula:

$$\text{DPPH (\%)} = (\text{Ac} - \text{At}) / \text{Ac} \times 100$$

DPPH (%): Percentage reduction in DPPH.

Ac: absorbance of the negative control

At: absorbance of the extract

The IC₅₀ value is the extract concentration, which ensures the 50% reduction in DPPH, determined graphically by linear regression, for each extract from the curve of the percentage reduction as a function of concentration [15].

Evaluation of Antibacterial Activity

The bacterial activity was evaluated on seven bacteria (*Pseudomonas sp*, *Escherichia coli*, *Bacillus subtilis*, *Bacillus cereus*, *Methicillin-resistant Staphylococcus aureus* or *SARM*, *Klebsiella sp*, *Staphylococcus aureus*).

In order to evaluate the antimicrobial activity of the extracts of the three algae, we used the diffusion method on the Muller Hinton agar medium [14]. Sterile Wattman filter paper discs measuring 6 millimeters in diameter are impregnated with 40 µl of the different extracts. Using sterile forceps, the discs are placed on the surface of a medium seeded with a microbial suspension with an optical density of 0.5 McFarland (10⁸CFU/ml). After diffusion, the plates are incubated for 18 to 24 hours at 37°C. Discs impregnated with different solvents will serve as controls.

All analyses were performed in triplicates and the experimental data were expressed as means±standard deviation using Microsoft Office Excel 2007.

Results and Discussion

Extraction Yields

Yields of crude extracts were determined relative to 50 g of plant material, and the results are shown in Fig. 1.

The method of extracting bioactive materials from plants plays an important role in providing consumers with a high-quality herbal product, and modern "advanced" extraction methods are superior to conventional methods. Extraction is considered one of the most crucial procedures in the manufacture of herbal products, which affects the active ingredients in the sample both qualitatively and quantitatively [16].

The best yield extraction obtained is that of *Cystoseira amentacea var. stricta* with a value of 6.4%, followed by that of *Ulva* (4.50%), and the lowest is recorded by the extract of *Corallina elongata*, with a rate of 1.2%. Results that are close to ours put forward a yield value of 2.53% for the methanolic extraction of *Corallina elongata* [17].




Fig. 1. Extraction yields of the algae.

Table 1. Total phenol and flavonoid contents of crude extracts of *Ulva lactuca*, *Cystoseira amentacea* var. *stricta* and *Corallina elongata*.

Crude extrats	Polyphenols (mg EAG/g extract)	Flavonoid (mg EQ/g extract)
<i>Ulva lactuca</i>	79.77±0.72	65.17±4.19
<i>Cystoseira amentacea</i>	73.92±0.53	36.2±3.33
<i>Corallina elongata</i>	219.43±5.79	61.6±1.37

Calculations of extraction yields are based on several factors, namely the extraction temperature, the initial plant material, and humidity [18]. This difference between species can also be attributed to the polarity of the compounds present in these algae, in addition to the nature of the solvent used and the chemical properties of the molecules to be extracted [19]. Likewise, the extraction method (maceration, decoction, infusion) also plays an important role in determining the yield and the chemical composition of the prepared extract [20].

Dosage of Phenolic Compounds

The contents of polyphenols and total flavonoids in our samples are reported in Table 1. A single dissolvent cannot extract all classes of phenolic compounds simultaneously and at a maximum concentration [21]. Therefore, we have used a mixture of two solvents.

From the results of a dosage of polyphenols we find that the methanol-toluene extract of *Corallina elongata* is rich in total polyphenols with a value of 219.43 mg/g, followed by the green algae extract *Ulva lactuca* with 79.7743 mg/g, while the extract of the brown algae *Cystoseira amentacea* is lower than the other two with a content of 73.92 mg/g. These results are not in agreement with other studies [22], which indicate that brown algae have higher polyphenolic contents compared to green and red algae. On the other hand, the values found are close to those [23] which put forward more or less comparable concentrations in *Cystoseires*: *Cystoseira crinita* (56.5 mg/g), *C. sedoides* (50.3 mg/g), *C. compressa* (61 mg/g), and which [24] recorded values of 37.5 mg/g of polyphenols in *Corallina elongata*.

According to the literature, methanolic extracts are the richest in phenolic compounds, therefore methanol remains the best solvent for extracting these compounds. This is due to the ability of methanol to inhibit the action

of polyphenol oxidase, which causes the oxidation of polyphenols in plant tissues [25, 26].

A wide range of algae species, including *Ulva fasciata*, were screened for antimicrobial activity. Researchers stated that drying reduced the bioactive compounds in the chosen algal species, and they found that methanol: toluene was the best solvent for extracting the antimicrobial compounds from fresh algae, and that the antimicrobial activity from chosen algae was due to a lipophilic compound that was stable over a broad range of temperatures (30–60°C) [27].

Many works suggest that maceration time affects the extraction rate [28–30] and the increasing temperature leads to an increasing extraction rate [31].

For flavonoids, we see that the *Ulva* extract has a higher content of flavonoids with a value of 65.17 mg/g, coming just after the *Corallina* red algae extract with 61.6 mg/g, the highest content. The lowest was recorded in *Cystoseira* with a value of 36.2 mg/g.

There exists little work on the flavonoid content in marine algae [32–34] in contrast to the numerous studies on flavonoids from terrestrial plants. A study on the distribution of flavonoids in 27 species of marine algae (6 *Chlorophyta*, 11 *Phaeophyta*, and 10 *Rhodophyta*) showed that they had a completely different flavonoid composition from fruits and vegetables [35].

The dosage of flavonoids is very important because they are a class of secondary plant phenols possessing potent antioxidant activity, thereby scavenging a wide range of reactive oxygen species and inhibiting lipoperoxidation, making them a potential therapeutic agent against a wide variety of diseases [36].

Dosage of Chlorophyll a, b, and Total Carotenoids

The results of the dosage of chlorophyll obtained are summarized in Table 2 and indicate that the brown alga *Cystoseira amentacea* var. *stricta* is richer in chlorophyll a and b (20.49 and 15.10 µg/ml) respectively than the green alga *Ulva lactuca* (7.96 and 14.12 µg/ml) and the red alga *Corallina elongata* (7.54 and 11.61 µg/ml). On the other hand, the highest carotenoid level is recorded by *U. lactuca*. These variations are probably linked to the species, geographic distribution, and age of the algae [17]. The effect of the solvent is not spared, although methanol is among the best solvents extractable from chlorophylls.

The main function of polar solvents is to improve and facilitate the permeability of the cell wall of the chemical

Table 2. Contents of chlorophyll a, b and total carotenoids in the extracts.

Algae	Contents	Chlorophyll a (µg/ml)	Chlorophyll b (µg/ml)	Total caroténoids (µg/ml)
<i>U. lactuca</i>		7.96±1.80	14.12±2.11	5552.76±5.60
<i>C. elongata</i>		7.54±1.17	11.61±0.90	884±3.30
<i>C. amentacea</i>		20.49±1.44	15.10±3.30	2215.43±3.67

Table 3. Percentages of DPPH inhibition by algae extracts.

	Concentration of extracts ($\mu\text{g/ml}$)					IC ₅₀ ($\mu\text{g/ml}$)
	750	500	350	250	50	
	Percentage reduction of DPPH %					
<i>U. lactuca</i>	68.06 \pm 1.59	62.34 \pm 2.01	66.55 \pm 2.19	45.09 \pm 2.20	23.46 \pm 1.64	329.38 \pm 2.39
<i>C. elongata</i>	72.43 \pm 3.04	65.12 \pm 2.50	54.32 \pm 4.08	56.7 \pm 3.79	30.54 \pm 1.70	275.70 \pm 2.93
<i>C. amentacea</i>	66.02 \pm 4.55	63.31 \pm 0.70	58.9 \pm 0.47	32.98 \pm 0.63	35.8 \pm 2.15	352.91 \pm 1.58

Diameter [mm]




Fig. 2. Antibacterial effects of crude extracts

substances, which allows better contact between the solvent and the solute, thus increasing the percentage yield of the extraction [37].

Evaluation of Antioxidant Activity

According to the results obtained for anti-radical activity presented in Table 3, the increase in anti-radical activity is proportional to the increase in the concentration of the extract. At the lowest concentration of 50 ($\mu\text{g/ml}$), the reduction rate of *Ulva* was the lowest with a value of 23.46% followed by *Corallina elongata* extract with 30.54%, and finally the extract from *Cystoseira* with 35.8%. At moderately high concentrations of 350 ($\mu\text{g/ml}$), the extracts show a progressive increase in DPPH reduction varying between 66.55%, 54.32%, and 58.9%, for *Ulva lactuca*, *Cystoseira amentacea*, and *Corallina elongata*, respectively. At the highest concentrations (750 $\mu\text{g/ml}$) we recorded values of 68.06%, 72.43%, and 66.02% for *Ulva lactuca*, *Corallina elongata*, and *Cystoseira amentacea*, respectively.

According to our results, the IC₅₀ has a value of 275.7 $\mu\text{g/ml}$ for *Corallina elongata*, 329.38 $\mu\text{g/ml}$ for

Ulva lactuca, and 352.91 $\mu\text{g/ml}$ for *Cystoseira amentacea var stricta*, and these values remain significantly higher than that of ascorbic acid (63.24 $\mu\text{g/ml}$), which is in agreement with the low percentages of inhibitions recorded by the different extracts.

The most effective antioxidant power was recorded with the red algae *Corallina*. Our results show very high IC₅₀s. The latter was also reported by others [17] who recorded IC₅₀s at 1787.7 $\mu\text{g/ml}$, and 3450 $\mu\text{g/ml}$ for methanolic extracts of *Corallina elongata* and *Cystoseira stricta*, respectively. We know that the IC₅₀ is inversely linked to the antioxidant capacity of a compound. However, an extract with low antioxidant activity cannot be considered a poor source of antioxidants, because an extract is composed of chemicals with different functional groups and polarities, allowing it to behave differently depending on the reaction mixture [38].

Because of different protocols and reaction conditions, IC₅₀ values for the same extracts vary a lot. Thus, it turns out to be more difficult to compare the results of different laboratories [39]. Light, oxygen, and pH of the reaction mixture also affect the absorbance of the DPPH solution [40].

Antimicrobial Activity

A study of 82 different marine macroalgae from the Chlorophyceae, Phaeophyceae, and Rhodophyceae classes were screened for antimicrobial activity [41]. Among the selected taxa, Phaeophyceae were reported to be the most active, with 84% of the algae from this taxa showing antimicrobial activity.

The results obtained for the antibacterial activity, reported in the histogram in Fig. 2, and expressed in terms of the diameter of the inhibition zone, showed that all bacteria are sensitive to one or other of the extracts except *Pseudomonas* which revealed its resistance for all three algae. The crude extracts of *Ulva* act on *B. cereus*, *B. subtilis*, and SARM with diameters of (8 mm), (10 mm), and (22 mm), respectively. This is consistent with the work [42] that noted the activity of methanolic extracts of *U. lactuca* on SARM with inhibition zones greater than 10 mm. Unlike those [43], who showed that antimicrobial activity is absent in *Ulva* algae extracts. A previous study [44] reported that the methanolic and aqueous extracts of 19 marine algal species collected along the western coast of Libya against 4 Gram-positive and 4 Gram-negative bacteria revealed that the observed inhibition zones values with methanolic extract of *U. lactuca* ranged between 11 mm for *S. aureus* and *Bacillus subtilis* and 14 mm for *P. aeruginosa* and *Klebsiella spp.* They also reported that *Cystoseira crinite* (Phaeophyceae) of the 19 examined algae, was the most potent against tested isolates.

The same results are noted for *Cystoseira* extract. Indeed, the latter acts on *B. cereus*, *B. subtilis*, and SARM with inhibition zones of (20 mm), (14 mm), and (21 mm), respectively. Researchers [42] who worked on two brown algae *Cystoseira humilis* and *C. bifurcata* showed that they had good activity against SARM with an inhibition zone diameter greater than 15 mm. *Cystoseira* crude extract has no effect on *Klebsiella*, *Staphylococcus*, *E. coli*, and *Pseudomonas*, contrary to other researchers [45], who reported that different *Cystoseira* species exhibited potent antibacterial activity against Gram-negative bacteria: *Salmonella typhi*, *E. coli*, *P. aeruginosa*, and *Klebsiella sp.*

The extracts of the two algae *Ulva* and *Cystoseira* have no effect on *E. coli*. These results are consistent with others [42, 46] which noted that only a few species of algae presented weak inhibitory activity against *E. coli*. However, a study [47] reported that the antibacterial activity of methanolic extracts of *U. lactuca* and *Enteromorpha compressa* (Chlorophyta), against 4 Gram-negative and 2 Gram-positive isolates showed that *U. lactuca* extract exhibited the most inhibitory effect for *K. pneumonia*, *S. aureus*, *P. vulgaris*, *B. subtilis*, and *P. aeruginosa*, followed by *E. compressa*. Nonetheless, the methanolic extract of *U. lactuca* was inactive against *E. coli*. Reports [48] have been made about the antibacterial activity of methanol/methylene chloride of *Ulva lactuca*, *Codiumto mentosum*, and *Hypnea musciformis* collected from the Suez Canal, Egypt. The latest study showed that *U. lactuca* has inhibitory activity only against *Salmonella typhimurium*,

K. pneumonia, *E. coli*, *Shigella boydii* and *S. aureus* with inhibitory zones ranging between 6–9 mm.

As for the *Corallina* extract, its effectiveness is noted against most strains except SARM and *Pseudomonas*. The inhibition zones vary from (16 mm) against *B. subtilis*, to 8mm for *Kleibseila* and *Staphylococcus*, passing through an inhibition zone of (12 mm) for *E. coli*.

Our results agree with those [49] which reported that high activity was shown on *E. coli* (32.00±1.73 mm) and *E. faecalis* (21.66±0.57 mm) with a methanolic extract of the red alga *Corallina officinalis*. Works [50] about the antimicrobial activities of some marine algae from the Mediterranean Sea (Algeria) show that the chloroform extracts of *Ulva lactuca* and *Corallina elongata* had the highest activity against *E. coli* and *Salmonella sp.*

Conclusions

The marine environment is a rich source of biological and chemical diversity. This diversity is at the origin of new chemical compounds likely to be used in industrial applications such as pharmaceutical products, cosmetics, and nutritional supplements. Finding new sources of purely natural bioactive substances is the main goal of most researchers today. This work is aimed at exploring the natural potential of marine algae from the west coast of Algeria. The various activity tests performed on the three species of algae show that the extracts studied contain a mixture of bioactive substances with a broad spectrum of activities which must be carefully studied and deepened in order to be able to use them for therapeutic purposes (antioxidants, antibiotics).

Acknowledgment

The authors are grateful to the staff of the faculty of Science, University Ahmed Ben Bella Oran 1, Algeria, for providing facilities to conduct the research.

Conflict of Interest

The authors declare that they have no conflicts of interest.

References

1. GAMFELDT L., LEFCHECK J.S., BYRNES J.E., CARDINALE B.J., DUFFY J.E., GRIFFIN J.N. Marine biodiversity and ecosystem functioning: what's known and what's next. *Oikos*, **124** (3), 252, **2015**.
2. BLUNT J.W, COPP B.R., MUNRO MURRAY H.G., NORTHCOTEC P.T., PRINSEP M.R. Marine natural products. Royal Society of Chemistry. Natural Products Reports, **32** (3), 382, **2015**.
3. MAURYA I.K., DILAWARI R., SINGH D., SINGH R.P. Bioactive compounds from extremophiles. Microbial Versatility in Varied Environments: Microbes in Sensitive Environments, 117, **2020**.

4. DUVAL R.E., GRARE M., DEMORE B. Fight against antimicrobial resistance: we always need new antibacterials but for right bacteria. *Molecules*, **24** (17), 3152, **2019**.
5. TERRENI M., TACCANI M., PREGNOLATO M. New Antibiotics for Multidrug-Resistant Bacterial Strains: Latest Research Developments and Future Perspectives, **26** (9), 2671, **2021**.
6. GOPEECHUND A., BHAGOOI R., NEERGHEEN V.S., BOLTON J.J., BAHORUN T., OZTURK M., EGAMBERDIEVA D., PEŠIĆ M. Macroalgae: status and future perspectives. In: *Biodiversity and Biomedicine*. Elsevier, Academic Press, USA, 257, **2020**.
7. GANESAN A.R., TIWARI U., RAJAURIA G. Seaweed nutraceuticals and their therapeutic role in disease prevention. *Food Science and Human Wellness*, **8** (3), 252, **2019**.
8. BIRIS-DORHOI E.S., MICHIU D., POP C.R., ROTAR A.M., TOFANA M., POP O.L., SOCACI S.A., FARCAS A.C. Macroalgae- a Sustainable Source of Chemical Compounds with Biological Activities. *Nutrients*, **12** (10), 3085. **2020**.
9. SAIM S., SAHNOUNI F., BOUHADI D., KHARBOUCHE S. The Antimicrobial Activity of Two Marine Red Algae Collected from Algerian West Coast. *Trends in pharmaceutical Sciences*, **7** (4), 233, **2021**.
10. NATARAJ G., VELMURGAN P., JAGADEESAN G., MANOHARAN A.L., THANGARAJ P. A comparative study of solvent extraction methods for a weed *Sphaeranthus amaranthoides* Burm. f.: phytochemical evaluation using chromatography (GC/MS and HPLC) and in vitro anti-diabetic activity. *Biomass Conversion and Biorefinery*, **1**, **2023**
11. JIMENEZ-ZAMORA A., DELGADO-ANDRADE C., RUFIAN-HENARES J.A. Antioxidant capacity, total phenols and color profile during the storage of selected plants used for infusion. *Food Chemistry*, **199**, 339, **2016**.
12. OREOPOULOU A., TSIMOGIANNIS D., OREOPOULOU V. Extraction of polyphenols from aromatic and medicinal plants: an overview of the methods and the effect of extraction parameters. *Polyphenols in plants*, 243, **2019**.
13. BUTNARIU M. Methods of analysis (extraction, separation, identification and quantification) of carotenoids from natural products. *Journal of Ecosystem and Ecography*, **6** (2), 1, **2016**.
14. CHAZAUX M., SCHIPHORST C., LAZZARI G., CAF-FARRI S. Precise estimation of chlorophyll a, b and carotenoid content by deconvolution of the absorption spectrum and new simultaneous equations for Chl determination. *Plant Journal*, **109** (6), 1630, **2022**.
15. DIOMANDE A., YOUSOUF S., HONORA D.T., ADAMA B., WITABOUNA K.M. Antioxidant power and phenolic compound content of two species of the genus *Albertisia*: *Albertisia cordifolia* (Mangenot and J. Miège) Forman and *Albertisia scandens* (Mangenot and J. Miège) Forman (Menispermaceae). *European Scientific Journal*, **14** (30), 128, **2018**.
16. LEHFA F., BELKHODJA H., SAHNOUNI F. Phytochemical Screening, Antioxidant and Anti-inflammatory Activities of Polyphenolic Extracts of Strawberry-tree Fruits (*Arbutus unedo* L.). *Journal of Applied Biotechnology Reports*, **10** (2), 992, **2023**.
17. OUCIF H., ADJOUT R., SEBAHI R., BOUKORT F., ALI-MEHIDI I., ABI-AYAD S.M.A. Comparison of in vitro antioxidant activity of some selected seaweeds from Algerian West Coast. *African Journal of Biotechnology*, **16** (26), 1474, **2017**.
18. JHA A.K., SIT N. Extraction of bioactive compounds from plant materials using combination of various novel methods: A review. *Trends in Food Science & Technology*, **119**, 579, **2022**.
19. SUJATHA R., SIVA D., NAWAS P.M.A. Screening of phytochemical profile and antibacterial activity of various solvent extracts of marine algae *Sargassum swartzii*. *World Scientific News*, (115), 27, **2019**
20. TEFIANI C., RIAZI A., YOUCEFI F., AAZZA S., GAGO C., FALEIRO M.L., PEDRO L.G., BARROSO J.G., FIGUEIREDO A.C., MEGÍAS C., CORTÉS-GIRALDO I., VIOQUE J., MIGUEL M.G. *Ammoides pusilla* (Apiaceae) and *Thymus munbyanus* (Lamiaceae) from Algeria essential oils: chemical composition, antimicrobial, antioxidant and antiproliferative activities. *Journal of Essential Oil Research*, **27** (2), 131, **2015**.
21. MOKRANI A., MADANI K. Effect of solvent, time and temperature on the extraction of phenolic compounds and antioxidant capacity of peach (*Prunus persica* L.) fruit. *Separation and Purification Technology*, **162**, 68, **2016**
22. PARTHIBAN C., SARANYA C., SOMASUNDARAM S.T., ANANTHARAMAN P. Antioxidant activities of some selected seaweeds from Tuticorin coast, Tamilnadu, India. *International Journal of Phytopharmacy Research*, **5** (1), 36, **2014**.
23. MHADHEBI L., CHAIEB K., BOURAOUI A. Evaluation of antimicrobial activity of organic extracts of six marine algae from Tunisian Mediterranean coasts. *International Journal of Pharmacy and Pharmaceutical Sciences*, **4** (1), 534, **2014**.
24. PINTEUS S., SILVA J., ALVES C., ILONA A., FINO N., RODRIGUES A., MEND PEDROSA R. Cytoprotective effect of seaweeds with high antioxidant activity from the Peniche coast (Portugal). *Food Chemistry*, **218**, 591, **2017**.
25. SINGH R., UPADHYAY A.K., CHANDRA P. Sodium chloride incites reactive oxygen species in green algae *Chlorococcum humicola* and *Chlorella vulgaris*: implication on lipid synthesis, mineral nutrients and antioxidant system. *Bioresource Technology*, **270**, 489, **2018**.
26. BOBO-GARCÍA G., ARROQUI C., MERINO G. Anti-browning compounds for minimally processed potatoes: A review. *Food Reviews International*, **36** (5), 529, **2020**
27. CHANDRASEKARAN M., VENKATESALU VENUGOPALAN R.A.J., ADAIKALA G. Antibacterial activity of *Ulva fasciata* against multidrug resistant bacterial strains. *International Journal of Natural Sciences*, **14**, **2014**.
28. CHEN Q., FUNG K.Y., LAU Y.T., NG K.M., LAU D.T. Relationship between maceration and extraction yield in the production of Chinese herbal medicine. *Food and Bioproducts Processing*, **98**, 236, **2016**.
29. JOVANOVIĆ A.A., ĐORĐEVIĆ V.B., ZDUNIĆ G.M., PLJEVLJAKUŠIĆ D.S., ŠAVIKIN, K.P., GOĐEVAC D.M., BUGARSKI B.M. Optimization of the extraction process of polyphenols from *Thymus serpyllum* L. herb using maceration, heat-and ultrasound-assisted techniques. *Separation and Purification Technology*, **179**, 369, **2017**.
30. HAYOUNI E.A., ABEDRABBA M., BOUIX M., HAMDI M. The effects of solvents and extraction method on the phenolic contents and biological activities in vitro of Tunisian *Quercus coccifera* L. and *Juniperus phoenicea* L. fruit extracts. *France. Food Chemistry*, **105** (3), 1126, **2007**.

31. HAYAT J., AKODAD M., MOUMEN A., BAGHOUR M., SKALLI A., EZRARI S. Phytochemical screening, polyphenols, flavonoids and tannin content, antioxidant activities and Ftir characterization of *Marrubium vulgare* L. from 2 different localities of Northeast of Morocco. *Heliyon*, **6** (11), **2020**.
32. BEN SAAD H., GARGOURI M., KALLEL F., CHAABENE R., BOUDAWARA T., JAMOUCI K., BEN AMARA I. Flavonoid compounds from the red marine alga *Alsidium corallinum* protect against potassium bromate-induced nephrotoxicity in adult mice. *Environmental Toxicology*, **32** (5), 1475, **2017**.
33. FERNANDO I.P.S., LEE W., AHN G. Marine algal flavonoids and phlorotannins; an intriguing frontier of biofunctional secondary metabolites. *Critical Reviews in Biotechnology*, **42** (1), 23, **2022**.
34. OUATTARA A., YAO K., KOMOE K., KPAN W.B., KONE M.W., OUATTARA D., TOURE S. Teneur en Composés Phénoliques et Activités Antioxydantes de *Sargassum fluitans* et *Sargassum natans*, deux Algues Brunes Invasives des Côtes Ivoiriennes. *European Scientific Journal*, **18** (33), 313, **2022**.
35. FREILE-PELEGRÍN Y., ROBLEDO D. Bioactive phenolic compounds from algae. Bioactive compounds from marine foods. *Plant and animal sources*, 113, **2013**.
36. PARSAEIMEHR A., LUTZU G. Algae as a novel source of antimicrobial compounds: current and future perspectives. *Antibiotic Resistance Mechanisms and New Antimicrobial Approaches*, 377, **2016**.
37. PATLE T.K., SHRIVAS K., KURREY R., UPADHYAY S., JANGDE R., CHAUHAN R. Phytochemical screening and determination of phenolics and flavonoids in *Dillenia pentagyna* using UV-vis and FTIR spectroscopy. *Spectrochimica Acta, Part A: Molecular and Biomolecular Spectroscopy (SAA)*, **242**, 118717, **2020**.
38. TUSEVSKI O., KOSTOVSKA A., ILOSKA A., TRAJKOVSKA L., GADZOVSKA SIMIC S. Phenolic production and antioxidant properties of some Macedonian medicinal plants. *Central European Journal of Biology*, **9** (9), 888, **2014**.
39. APAK R.A., ÖZYÜREK M., GÜÇLÜ K., ÇAPANOĞLU E. Antioxidant activity/capacity measurement. 2. Hydrogen atom transfer (HAT)-based, mixed-mode (electron transfer (ET)/HAT), and lipid peroxidation assays. *Journal of Agricultural and Food Chemistry*, **64** (5), 1028, **2016**.
40. APAK R.A. Current issues in antioxidant measurement. *Journal of Agricultural and Food Chemistry*, **67** (33), 9187, **2019**.
41. SALVADOR N., GOMEZ GARRETA M.A., LAVELI L., RIBERA M.A. Antimicrobial activity of Iberian macroalgae. *Scientia Marina*, **71** (1), 101, **2007**.
42. YOUNES F., ETAHIRI S., ASSOBBEI O. Activité antimicrobienne des algues marines de la lagune d'Oualidia (Maroc): Criblage et optimisation de la période de la récolte. *Journal of Applied Biosciences*, **24**, 1543, **2009**.
43. KUMAR A.K., RENGASAMY R. Evaluation of antibacterial potential of seaweeds occurring along the coast of Tamil Nadu, India against the plant pathogenic bacterium *Xanthomonas oryzae* pv. *oryzae* (Ishiyama) Dye. *Botanica Marina*, **43**, 409, **2000**.
44. ALGHAZEER R., WHIDA F., ABDUELRHMAN E., GAMMOUDI F. AZWAI S. Screening of Antibacterial Activity in Marine Green, Red and Brown Macroalgae from the Western Coast of Libya. *Natural Sciences*, **5**, 7, **2013**.
45. CHIHEB I., RIADI H., MARTINEZ-LOPEZ J., DOMINGUEZ SEGLAR J.F., GOMEZ VIDAL J.A., BOUZIANE H., KADIRI M. Screening of antibacterial activity in marine green and brown macroalgae from the coast of Morocco. *African Journal of Biotechnology*, **8** (7), 1258, **2009**.
46. FEBLES C.I., ARIAS A., HARDISSON A., LOPEZ A.S. Antimicrobial activity of extracts from Canary species of *Phaeophyta* and *Chlorophyta*. *Phytotherapy Research*, **9** (5), 385, **1995**.
47. ELNABRIS K.J., ELMANAMA A.A. CHIHADDEH W.N. Antibacterial Activity of Four Marine Seaweeds Collected from the Coast of Gaza Strip, Palestine. *Mesopotamian Journal of Marine Science*, **28** (1), 81, **2013**.
48. HAMZA E.H., TEMRAZ T.A. AHMED S.A. Bioactivity of Some Egyptian Seaweeds Extract. *Catrina*, **11** (1), 17, **2015**.
49. TASKIN E., OZTURK M., TASKIN E., KURT O. Antibacterial activities of some marine algae from the Aegean Sea (Turkey). *African Journal of Biotechnology*, **6** (24), 2746, **2007**.
50. ZOUAOU B., GHALEM B.R. The phenolic contents and antimicrobial activities of some marine algae from the Mediterranean Sea (Algeria). *Russian Journal of Marine Biology*, **43**, 491, **2017**.