

Original Research

Molecular Identification and Genetic Characterization of *Trypanosoma Evansi* from Cattle in Makkah, Saudi Arabia

Haleema H. Albohiri, Muslimah N. Alsulami*

Department of Biology, College of Science, University of Jeddah, Jeddah, Saudi Arabia

Received: 18 March 2024

Accepted: 27 May 2024

Abstract

Surra is an infectious disease caused by *Trypanosoma evansi* (*T. evansi*), which is spread in developing countries. This study aimed to estimate the prevalence and genetic characterization of *T. evansi* in cattle in Makkah, Saudi Arabia. Blood samples from 150 cattle were collected and examined by a blood smear and PCR targeting the RoTat 1.2VSG gene. The findings revealed that the prevalence of *T. evansi* varied depending on the detection test, ranging from 20.7% to 25.3% by blood smear examination and PCR. RoTat 1.2VSG sequences analysis of two *T. evansi* isolates revealed little difference compared to similar sequences in the database. The phylogenetic tree revealed that local *T. evansi* isolates in this study were clustered with other *T. evansi* sequences from Egypt, India, Nigeria, Kenya, and Pakistan. The sequences in the study and the Egyptian sequence (MG674485, camel strain) were grouped into a single clade with little genetic variation. In conclusion, the results of this study urge for the introduction and adoption of appropriate control strategies to reduce the effect of *T. evansi* infection on cattle production in Saudi Arabia.

Keywords: *Trypanosoma evansi*, Cattel, RoTat 1.2VSG gene, Saudi Arabia

Introduction

Trypanosoma evansi (*T. evansi*), a haemoprotozoan parasite, is the causative agent of surra or trypanosomiasis in tropical and subtropical regions and affects a wide range of domestic and wild animals, including cattle, camels, horses, and various wildlife species [1-3]. *T. evansi* is a serious and potentially fatal disease causing substantial economic losses in animal production due to decreased productivity, and loss of

weight. Furthermore, *T. evansi* has been linked to failure in vaccination, which can substantially impact global trade in live animals and their products, in addition to the widespread utilization of trypanocides [4-6].

Globally, it has been reported in Africa, Asia, Europe, and South America that the susceptibility to *T. evansi* infection differs among host species, and the prevalence varies geographically [5]. In Asia, *T. evansi* is prevalent in the Arabian Peninsula, including Saudi Arabia, the United Arab Emirates, Jordan, Lebanon, Syria, Iraq, and Pakistan [7-9].

Diagnosis of *T. evansi* can be challenging due to the intermittent presence of the parasite in the bloodstream and low parasitemia due to early or chronic infections

*e-mail: mnal-sulami@uj.edu.sa

[10, 11]. Several diagnostic methods can be used, including microscopic examination of blood smears, serological tests, and PCR amplification of parasitic DNA [12]. Although the blood smear examination and serological assay have been used for years in the standard diagnosis of *T. evansi*, both assays presented have low sensitivity to detect and differentiate between the various *Trypanosoma* species in animals. Therefore, different molecular assays have been developed to overcome the limitations of parasitological and serological methods [12, 13] and improve the diagnosis of the disease in both early and chronic infections [14, 15]. Moreover, it was found that PCR showed high sensitivity and specificity, being able to detect *T. evansi* at very low levels, ranging from 1 to 10 trypanosomes per one milliliter of blood [16].

Further, *T. evansi* can be amplified and sequenced using several primer sequences targeting different regions or genes, including kinetoplast DNA, the internal transcribed spacer area, ribosomal DNA, and variable surface glycoprotein (VSG) genes with varying degrees of sensitivity and specificity [10, 17-19]. According to [18, 20], the RoTat 1.2VSG sequence is distinct and specific for the identification and characterization of *T. evansi*.

In Saudi Arabia, *T. evansi* infection is endemic in camels and recently the parasite has also been reported in horses, either microscopically or genetically [7, 21- 26]. A 2018 study using PCR and blood smear for detection of *T. evansi* in 4.3% and 5.6% of blood samples from stray dogs by PCR and blood smear, respectively [27]. A 2009 study examining blood samples from camels, cattle, sheep, and goats found *T. evansi* infection in camels, but this study did not detect *T. evansi* in cattle, sheep, and goats by microscopy depending on the region [22]. In cattle, the high seroprevalence of *T. evansi* was reported from India and Egypt [2, 28] and a high molecular prevalence of *T. evansi* in blood samples obtained from Egyptian cattle has been reported [10]. Therefore, these studies conducted in Saudi Arabia and other countries indicate that this parasite may exist in cattle.

Additionally, the livestock sector in Saudi Arabia has a positive long-run response to Agricultural Growth Domestic Product, where it might be considered as the leading subsector in the economy [29]. Furthermore, Saudi Arabia leads the livestock sector in the Gulf Cooperation Council (GCC) region, particularly in the production of animal products. Therefore, cattle are essential for food production and economic stability in Saudi Arabia [30]. There are no published data on *T. evansi* for cattle in Saudi Arabia. Therefore, the current study aimed to use blood smear examination and PCR to determine the prevalence of *T. evansi* in naturally infected cattle from Makkah city, and to genetically characterize *T. evansi* isolates utilizing the RoTat 1.2 VSG gene.

Methods

Study Area and Sampling

The study was performed at the AIKai'ah slaughtering house, one of the hajj abattoirs, which was chosen because it processes all kinds of animals, in Makkah City (21.4241°N, 39.8173°E) in Saudi Arabia (Fig. 1). According to the Koppen-Geiger climate classification, the climate is a hot desert type. During the year 2023, 150 blood samples were randomly collected from cattle submitted for slaughter. Blood samples were collected from the jugular vein and transferred into EDTA tubes (BD Vacutainer® Tube, Gribbles Pathology, VIC, Australia). The samples were then sent in a cool container to the Parasitology Laboratory of the Biological Department, College of Science, University of Jeddah, for microscopic examination. A thin blood smear was used to confirm the presence of *T. evansi*. In brief, a drop of EDTA blood was placed and spread on a clean slide. The smear was dried, followed by fixation with absolute methanol for 3 minutes. Afterward, excess methanol was removed, and the thin smear was stained by Giemsa stain diluted at a ratio of 1:20 for 20 minutes. Finally, the blood film was examined under magnifications of 100× (using oil immersion microscopy) to detect trypanomastigotes [31].

DNA Extraction

Total genomic DNA (gDNA) was extracted from 150 blood samples (200 µl) using the DNeasy Blood and Tissue Kit (Qiagen, Hilden, Germany), following the protocol described by the manufacturer, and the gDNA was stored at -20°C until PCR testing.

PCR

PCR was performed on all gDNA extracts for the detection of partial fragment (488 bp) of *T. evansi* type A using the RoTat 1.2 VSG primer pairs: forward primer: 5'-GCCACCACGGCGAAAGAC-3'; reverse primer: 5'-TAATCAGTGTGGTGTGC-3' [32]. PCR reaction was performed in a 25 µL reaction volume containing 12.5 µL of 2X GoTaq® Green Master Mix (Promega, USA), 1 µL (0.5 µM) of each primer, and 2 µL of DNA extract and 8.5 µL of ddH₂O. The PCR protocol was one cycle for 3 min at 94°C, 30 cycles (94°C for 60 s, 58°C for 60 s, and 72°C for 60 s), and one cycle at 72°C for 5 minutes. Negative (no DNA template) control was included. The PCR result that was amplified was placed on a 1.5% agarose gel, stained with 0.5 µg/ml of ethidium bromide, and examined using a gel documentation system (Biospectrum UVP, UK).

Sequence Analysis and Phylogenetic Analysis

Twelve positive RoTat 1.2VSG PCR samples were chosen based on the density of the band and PCR




Fig. 1. Site of *T. evansi* in cattle collected from Al Kakee's Slaughter, Makkah, Saudi Arabia (General Authority of Survey and Geospatial Information).

products were purified by QIAquick PCR Purification Kit (Qiagen, Germany). Sanger sequencing of purified DNA was performed in an automated DNA sequencer (ABI 3730XL, Solgent Co. Ltd., South Korea). The sequences were read and edited manually using DNA BaserV3 software (Heracle BioSoft S.R.L./ Romania) to fix any potential calling errors. The Basic Local Alignment Search Tool (BLAST) (<http://www.ncbi.nlm.nih.gov/BLAST>) was used to search for sequence similarity. The RoTat 1.2VSG sequences were aligned using Clustal W in MEGA X software (www.megasoftware.net) with sequences available in the GenBank database. For phylogenetic tree construction, the Neighbor-Joining (NJ) method was used in MEGA X with 1000 thorough bootstrap replicates. The evolutionary distances are expressed in base substitutions per site and were calculated using the Maximum Composite Likelihood technique. There were 19 nucleotide sequences in this investigation. All gaps and missing data were removed (complete deletion option). Of the 12 partial sequences, two were identified in the study and have been deposited in GenBank under the accession number PP105580 and PP105581.

Statistical Analysis

The kappa agreement test between blood smear and PCR was calculated using online statistical tools (<http://vassarstats.net>). The assessment of the sensitivity and specificity of blood smear compared to PCR as a reference test were calculated using MedCalc Software Ltd. (<https://www.medcalc.org/calc>; Version 20.115).

Results

PCR and Microscopic Analysis of Stained Blood Smears

According to microscopic analysis of stained blood smears, the overall prevalence of *T. evansi* among 150 cattle was 20.7% (31/150) and reached 25.3% (38/150) by PCR targeting RoTat 1.2VSG. Furthermore, the 31 blood smear-positive samples were also PCR positive. Based on the positive and negative results, the Kappa agreement between the two tests (blood smear and PCR) showed very substantial agreement (0.71; 95% CI = 0.63-0.88) (Table 1). In addition to the kappa results, the blood smear showed 91.33% accuracy, 73.68% sensitivity, and 97.32% specificity in detecting *Brucella* antibodies in cattle compared to the results of PCR as a reference test (Table 2).

Phylogenetic Analysis of RoTat 1.2VSG Gene of *T. Evansi*

Sequences (PP105580, PP105581) were searched in the GenBank database that confirmed the PCR amplicons' identity as *T. evansi* with an identity of 99% (3 nt substitution). The BLAST search revealed that the closest hit for all *Trypanosoma* sequences was *T. evansi* with 97.5-100% nucleotide identity and the genetic divergence ranged from 0.01 to 1.97 (Fig. 2).

Phylogenetic analysis (Fig. 3) of the partial amplified RoTat 1.2VSG gene of *T. evansi* revealed a clustering of our isolates with groups with similar sequences previously identified in camels, cattle, *Panthera leo*, and *Panthera tigris tigris* from Egypt, India, Pakistan, Kenya, and Nigeria with 59% nodal support. The sequences presented in this study are closely related

Table 1. Kappa value, standard error, and 95% confidence interval for the test agreement (Blood smear vs. PCR) for *T. evansi* diagnosis in 150 cattle.

Variable		PCR		Total	K-value	Standard Error	0.95% Confidence Interval
Blood smear		Positive	Negative				
	Positive	28	3	31	0.71	0.065	0.63-0.88
	Negative	10	109	119			
Total	38	112	150				

Table 2. Estimated sensitivity and specificity of blood smear compared to PCR results.

Statistic	Value %	95% CI
True positive	28	
False positive	3	
True negative	109	
False negative	10	
Sensitivity	73.68	56.90-86.60
Specificity	97.32	92.37-99.44
Positive predictive value*	90.32	75.05-96.66
Negative predictive value*	91.60	86.48-94.89
Accuracy*	91.33	85.64-95.30

*, these values are based on the prevalence of *T. evansi*.

to *T. evansi* isolate found in Egyptian camels (Accession number: MG674185) with 96% nodal support. They also showed a little genetic divergence of 1.84-1.97 from the RoTat 1.2VSG gene of the *T. evansi* found in horses

(Accession number: KU589274-MT501210, India) and buffalo (Accession number: EF495337, India) (Fig. 2). The resulting pattern supports the soundness of RoTat 1.2VSG analysis as a particular marker of *T. evansi* type A and identifies minor changes between closely associated sequences. A partial sequence from the ITS1 region of *T. evansi* (Accession number MH247175.2) was included as an outgroup for phylogenetic analysis (Fig. 3).

Discussion

Molecular biology facilitates epidemiological surveys of infectious diseases, particularly trypanosomiasis in domestic animals; nonetheless, *T. evansi* in livestock has received little attention until recently. During the previous period, Saudi Arabia's cattle industry experienced significant expansion to fulfill the rising demand for its meat.

In our study, the prevalence of *T. evansi* infection in cattle was 20.7% using blood smear examination, which is higher than the 2.75% reported in cattle and buffaloes from India by conventional parasitological methods [15]

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1 PP105581 <i>T. evansi</i> (Cattle-Saudi Arabia)		0.01	3.69	5.96	3.69	3.69	3.69	3.69	3.69	3.69	3.69	3.69	3.69	6.43	6.13	3.69	0.02	3.28
2 PP105580 <i>T. evansi</i> (Cattle-Saudi Arabia)	0.01		3.68	5.95	3.68	3.68	3.68	3.68	3.68	3.68	3.68	3.68	3.68	6.43	6.13	3.68	0.01	3.28
3 MK367833 <i>T. evansi</i> (Camel-Kenya)	1.16	1.15		6.72	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	6.27	6.20	0.00	3.70	2.65
4 KU589274 <i>T. evansi</i> (Horse-India)	1.95	1.97	1.86		6.72	6.72	6.72	6.72	6.72	6.72	6.72	6.72	6.72	0.45	0.74	6.72	5.98	9.98
5 OMI72432 <i>T. evansi</i> (Camel-Nigeria)	1.16	1.15	0.00	1.86		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	6.27	6.20	0.00	3.70	2.65
6 OMI72427 <i>T. evansi</i> (Camel-Nigeria)	1.16	1.15	0.00	1.86	0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	6.27	6.20	0.00	3.70	2.65
7 OMI72425 <i>T. evansi</i> (Camel-Nigeria)	1.16	1.15	0.00	1.86	0.00	0.00		0.00	0.00	0.00	0.00	0.00	0.00	6.27	6.20	0.00	3.70	2.65
8 MW881772 <i>T. evansi</i> (<i>Panthera leo</i> -Pakistan)	1.16	1.15	0.00	1.86	0.00	0.00	0.00		0.00	0.00	0.00	0.00	0.00	6.27	6.20	0.00	3.70	2.65
9 MW881771 <i>T. evansi</i> (<i>Panthera leo</i> -Pakistan)	1.16	1.15	0.00	1.86	0.00	0.00	0.00	0.00		0.00	0.00	0.00	0.00	6.27	6.20	0.00	3.70	2.65
10 MW865397 <i>T. evansi</i> (<i>Panthera tigris tigris</i> -Pakistan)	1.16	1.15	0.00	1.86	0.00	0.00	0.00	0.00	0.00		0.00	0.00	0.00	6.27	6.20	0.00	3.70	2.65
11 KF726106 <i>T. evansi</i> (Cattle-Egypt)	1.16	1.15	0.00	1.86	0.00	0.00	0.00	0.00	0.00	0.00		0.00	0.00	6.27	6.20	0.00	3.70	2.65
12 JX888091 <i>T. evansi</i> (Camel-Egypt)	1.16	1.15	0.00	1.86	0.00	0.00	0.00	0.00	0.00	0.00	0.00		0.00	6.27	6.20	0.00	3.70	2.65
13 JX134605 <i>T. evansi</i> (Camel-India)	1.16	1.15	0.00	1.86	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00		6.27	6.20	0.00	3.70	2.65
14 EF495337 <i>T. evansi</i> (Buffalo-India)	1.84	1.86	1.58	0.16	1.58	1.58	1.58	1.58	1.58	1.58	1.58	1.58	1.58		0.02	6.27	6.47	7.55
15 MT501210 <i>T. evansi</i> (Horse-India)	1.84	1.86	1.52	0.18	1.52	1.52	1.52	1.52	1.52	1.52	1.52	1.52	1.52	0.02		6.20	6.17	7.46
16 OR039342 <i>T. evansi</i> (Cattle-India)	1.16	1.15	0.00	1.86	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.58	1.52		3.70	2.65
17 MG674185 <i>T. evansi</i> (Camel-Egypt)	0.03	0.02	1.24	1.94	1.24	1.24	1.24	1.24	1.24	1.24	1.24	1.24	1.24	1.95	1.95	1.24		3.28
18 MZ032003 <i>T. evansi</i> (Camel-India)	1.46	1.39	0.88	1.75	0.88	0.88	0.88	0.88	0.88	0.88	0.88	0.88	0.88	1.44	1.50	0.88	1.34	

Fig. 2. Divergence between partial RoTat 1.2VSG *T. evansi* sequences (Accession number: PP105580 - PP105581) and available reference *T. evansi* nt sequences of RoTat 1.2VSG gene. The number of base substitutions for each site between sequences is shown.




Fig. 3. Phylogenetic tree based on partial RoTat 1.2VSG gene nt sequences from *T. evansi* isolates from cattle in Saudi Arabia and nt sequences worldwide, available in GenBank. The evolutionary history was inferred using the Neighbor-Joining method. Two sequences of *T. evansi* obtained in the present study are represented by a black circle.

and 1% in cattle from Indonesia [33] but lower than the 42.2% found in Egyptian cattle by ELISA [28]. The PCR revealed a higher infection rate (25.3%) in cattle, which is lower than the 30.4% found in Egyptian cattle by PCR [10] and higher than the infection rate (3%) reported in Indonesia [33]. Several factors explain the difference in the detection rate of *T. evansi* infection by blood smear screening and PCR. Among these factors is the low sensitivity of the blood smear, with a detection limit of 10^5 trypanosomes/ml [5, 34]. During the chronic phase or the presence of the parasite in low quantities, PCR showed the ability to detect 1-20 parasites/ml of blood [35, 36] in studies that supported our results.

The sensitivity of blood smear (73.68%) was close to the 76% observed in Egypt [37] and higher than the 27.02% recorded in India [38]. However, the specificity of the blood smear (97.32%) was lower than the 100% reported in Egypt [37]. Although PCR increased *T. evansi* detection in blood samples that tested negative by blood smear, we found substantial agreement ($\kappa = 0.71$, 95% CI = 0.63-0.88) between blood smear and PCR results. A previous study in Egypt found almost perfect kappa agreement ($\kappa = 0.83$) between the results of a blood smear, CATT, and PCR tests [37]. Another study in Pakistan reported poor agreement between blood smears and PCR [19]. This variation is due to changes in the number of samples, location, farm management, and animal conditions.

This study tried to genetically characterize the prevalence of *T. evansi* in cattle in Saudi Arabia. Thirty-eight of the 150 cattle sampled in this study were positive for *T. evansi* by RoTat 1.2VSG PCR. Previously in Saudi Arabia, *T. evansi* infection has been reported in camels, horses, and dogs either microscopically

or genetically [7, 21-26]; nevertheless, we could not find any sequences of *T. evansi* targeting RoTat 1.2VSG gene indexed in GenBank. Moreover, no published data on *T. evansi* are available for cattle in Saudi Arabia. Therefore, this is the first study reporting the parasite in Saudi Arabia microscopically and molecularly using the RoTat 1.2 VSG gene. Our findings are supported by a previous study from Egypt, where the presence of the *T. evansi* parasite was first reported in cattle raised alongside camels [10] and infection was confirmed by PCR targeting the ITS and RoTat 1.2 VSG genes and Sanger sequencing of the VSG gene.

Analysis of nucleotide sequence pairwise alignment of RoTat 1.2VSG from *T. evansi* ($n = 2$) revealed a high degree of identity (>99%) in between and ranged 97.9-100% with those retrieved from the GenBank. This pattern indicated that the RoTat 1.2VSG gene was useful and informative for the genetic characterization of *T. evansi* isolates and to distinguish it from other members of the *Trypanozoon* subgenus [32, 39]. Recently, [40] reported a non-RoTat 1.2 *T. evansi* in Kenya and classified it as *T. evansi* type B. The findings presented here revealed that the *T. evansi* isolates found in cattle have the RoTat 1.2 VSG encoding gene. Although the RoTat 1.2VSG region is substantially conserved across *T. evansi* strains [20], two *T. evansi* sequences have been created with little variation, most likely due to the influence of geography, host, chronic disease, and treatment [39, 41].

Phylogenetic analyses of the partial RoTat 1.2VSG gene (Fig. 3) showed that the two obtained *T. evansi* sequences were clustered into one clade and revealed intraspecies genetic difference in at least two groups within one host (cattle) and were clustered close

to the *T. evansi* sequence (Accession number MG674158) isolated from camels in Egypt sequences. Previously, the genetic diversity of *T. evansi* isolated from camels and cattle has been reported using the RoTat 1.2VSG gene [10, 38]. Further, the genetic diversity of *T. evansi* isolated from camels using the ITS-1 and ESAG6 genes has also been reported [42, 43].

Conclusions

The results of this study showed that *T. evansi* is circulating in cattle from Makkah for the first time. Two sequences were identified, based on RoTat 1.2VSG, with little genetic difference. The correlation between blood smear results and PCR results confirmed the estimate made by blood smear examination. For continuous screening, PCR testing is the preferred and recommended diagnostic method. However, it is necessary to genetically characterize the circulating *T. evansi* species.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflict of interest

There are no conflicts of interest.

References

- ALGEHANI A.M.G., JABER F.A., KHAN A., ALSULAMI M.N. Review on trypanosomiasis and their prevalence in some country on the Red Sea. *Brazilian Journal of Biology*, **83**, 2021.
- CHANDU A.G.S., SENGUPTA P.P., JACOB S.S., BORTHAKUR S.K., PATRA G., ROY P. Mining the pervasiveness of surra in different animal species of Northeastern states of India: Assam, Mizoram and Tripura. *Journal of Parasitic Diseases*, **45**, 330, 2021.
- JABER F.A., ALGEHANI A.M., ALMALKI G., ALBOHIRI H.H., ALSULAMI M.N. Histological and Immunohistochemical Assays of *Trypanosoma evansi* Infected Camel Hepatic Tissues. *Polish Journal of Environmental Studies*, **32** (5), 4617, 2023.
- EREQAT S., NASEREDDIN A., AL-JAWABREH A., AL-JAWABREH H., AL-LAHAM N., ABDEEN Z. Prevalence of *Trypanosoma evansi* in livestock in Palestine. *Parasites and vectors*, **13**, 1, 2020
- AREGAWI W.G., AGGA G.E., ABDI R.D., BÜSCHER P. Systematic review and meta-analysis on the global distribution, host range, and prevalence of *Trypanosoma evansi*. *Parasites and vectors*, **12**, 1, 2019.
- MAMOUDOU A., NJANLOGA A., HAYATOU A., SUH P.F., ACHUKWI M.D. Animal trypanosomosis in clinically healthy cattle of north Cameroon: epidemiological implications. *Parasites & Vectors*, **9**, 1, 2016.
- AL MALKI J.S., HUSSIEN N.A. Molecular characterization of *Trypanosoma evansi*, *T. vivax* and *T. congolense* in camels (*Camelus dromedarius*) of KSA. *BMC Veterinary Research*, **18** (45), 2022.
- SALAH A.A., ROBERTSON I.D., MOHAMED A.S. Prevalence and distribution of *Trypanosoma evansi* in camels in Somaliland. *Tropical animal health and production*, **51** (8), 2371, 2019.
- TEHSEEN S., JAHAN, N., DESQUESNES M., SHAHZAD M.I., QAMAR M.F. Field investigation of *Trypanosoma evansi* and comparative analysis of diagnostic tests in horses from Bahawalpur, Pakistan. *Turkish Journal of Veterinary and Animal Sciences*, **41** (2), 288, 2017.
- ELHAIG M.M., SELIM A., MAHMOUD M.M., EL-GAYAR E.K. Molecular confirmation of *Trypanosoma evansi* and *Babesia bigemina* in cattle from lower Egypt. *Pakistan Veterinary Journal*, **36**, 409, 2016.
- SADEK A., EL-KHABAZ K., EL-GENEDY S., EL-GIOUSHY M. Comparative diagnostic performance of microscopic examination, polyclonal antigen-elisa, and polymerase chain reaction for the detection of *Trypanosoma evansi* in camels (*Camelus dromedarius*). *Advances in Animal and Veterinary Sciences*, **9**, 1004, 2021.
- MAMMAN S.A., DAKUL D.A., YOHANNA J.A., DOGO G.A., REUBEN R.C., OGUNLEYE O.O., TYEM D.A., PETER J.G., KAMANI J. Parasitological, serological, and molecular survey of trypanosomosis (*Surra*) in camels slaughtered in northwestern Nigeria. *Tropical Animal Health and Production*, **53**, 1, 2021.
- BARGHASH S.M. Molecular changes in *Trypanosoma evansi* after treatment against trypanosomosis. *Annals of Parasitology*, **66**, 2020.
- MULENGA G.M., NAMANGALA B., CHILONGO K., MUBAMBA C., HAYASHIDA K., HENNING L., GUMMOW B. Challenges in the diagnostic performance of parasitological and molecular tests in the surveillance of African trypanosomiasis in eastern Zambia. *Tropical Medicine and Infectious Disease*, **6** (68), 2021.
- SIVAJOTHI S., RAYULU V.C., MALAKONDAIAH P., SREENIVASULU D. Diagnosis of *Trypanosoma evansi* in bovines by indirect ELISA. *Journal of Parasitic Diseases*, **40**, 141, 2016.
- DABO N.T., MAIGARI A.K. Soft options for effective diagnosis of African animal trypanosomiasis: a review. *International Journal of Medical Evaluation and Physical Report*, **2** (2), 1, 2017.
- ELWATHIG M., FAYE B., THEVENON S., RAVEL S., BOSSARD G. Epidemiological surveys of camel trypanosomosis in Al-jouf, Saudi Arabia based on PCR and ELISA. *Emirates Journal of food and Agriculture*, **212**, 2016.
- TONG Q., CHEN R., KONG Q., GOOSSENS J., RADWANSKA M., LOU D., LU S. DNA detection of *Trypanosoma evansi*: Diagnostic validity of a new assay based on loop-mediated isothermal amplification (LAMP). *Veterinary parasitology*, **250**, 1, 2018.
- IHASSAN-KADLE A.A., IBRAHIM A.M., NYINGILILI H.S., YUSUF A.A., VIEIRA T.S., VIEIRA R.F. Parasitological, serological and molecular survey of camel trypanosomiasis in Somalia. *Parasites & vectors*, **12**, 1, 2019.
- SUGANUMA K., NARANTSATSRAL S., BATTUR B., YAMASAKI S., OTGONSUREN D., MUSINGUZI

- S.P., INOUE N. Isolation, cultivation and molecular characterization of a new *Trypanosoma equiperdum* strain in Mongolia. *Parasites and vectors*, **9**, 1, **2016**.
21. AL-KHARUSI A., ELSHAFIE E.I., BAQIR S., FARAZ A., AL-ANSARI A., BURGER P., ROBERTS D. (2022). Detection of *Trypanosoma* infection in dromedary camels by using different diagnostic techniques in northern Oman. *Animals*, **12** (11), 1348, **2022**.
 22. PERVEEN N., MUZAFFAR S.B., AL-DEEB M.A. Ticks and tick-borne diseases of livestock in the Middle East and North Africa: A review. *Insects*, **12** (1), 83, **2021**.
 23. AMIN Y.A., NOSEER E.A., FOUAD S.S., ALI R.A., MAHMOUD H.Y. Changes of reproductive indices of the testis due to *Trypanosoma evansi* infection in dromedary bulls (*Camelus dromedarius*): semen picture, hormonal profile, histopathology, oxidative parameters, and hematobiochemical profile. *Journal of Advanced Veterinary and Animal Research*, **7** (3), 537, **2020**.
 24. ALANAZI A.D., PUSCHENDORF R., SALIM B., ALYOUSIF M.S., ALANAZI I.O., AL-SHEHRI H.R. Molecular detection of equine trypanosomiasis in the Riyadh Province of Saudi Arabia. *Journal of Veterinary Diagnostic Investigation*, **30**, 942, **2018**.
 25. ALI A., DERAR D.R. Ovary and ovarian bursa in dromedary camels: clinical relevance of the topographical features. *Anatomia, Histologia, Embryologia*, **49** (3), 325, **2020**.
 26. METWALLY D.M., AL-TURAIKI I.M., ALTWAIJRY N., ALGHAMDI S.Q., ALANAZI A.D. Molecular identification of *trypanosoma evansi* isolated from arabian camels (*camelus dromedarius*) in Riyadh and Al-Qassim, Saudi Arabia. *Animals*, **11**, 1149, **2021**.
 27. ALANAZI A.D. Parasitological and molecular detection of canine trypanosomiasis from Riyadh Province, Saudi Arabia. *Journal of Parasitology*, **104**, 539, **2018**.
 28. FERREIG R.M., MOHAMED S.G.A., MAHMOUD H.Y.A.H., ABOULAILA M.R., GUSWANTO A., NGUYEN T.-T., MOHAMED, A.E.A., INOUE N., IGARASHI I., NISHIKAWA Y. Seroprevalence of *Babesia bovis*, *B. bigemina*, *Trypanosoma evansi*, and *Anaplasma marginale* antibodies in cattle in southern Egypt. *Ticks and tick-borne diseases*, **8**, 125, **2017**.
 29. EMAM A.A., ABASS A.S., ELMULTHUM N.A., ELRASHEED M. Status and prospects of agricultural growth domestic product in the Kingdom of Saudi Arabia. *Sage Open*, **11**, 2158244021100545, **2021**.
 30. MANSOUR A.T., AHMED A.E. Role of Livestock on Food Security in Saudi Arabia, In: *Food and Nutrition Security in the Kingdom of Saudi Arabia*, Vol. 1: National Analysis of Agricultural and Food Security. Springer, 141, **2024**.
 31. EREQAT S., NASEREDDIN A., AL-JAWABREH A., AL-JAWABREH H., AL-LAHAM N., ABDEEN Z. Prevalence of *Trypanosoma evansi* in livestock in Palestine. *Parasites & vectors*, **13**, 1, **2020**.
 32. BOULANGÉ A., PILLAY D., CHEVTZOFF C., BITEAU N., DE GRAÇA V.C., REMPETERS L., BALTZ T. Development of a rapid antibody test for point-of-care diagnosis of animal African trypanosomosis. *Veterinary parasitology*, **233**, 32, **2017**.
 33. SETIAWAN A., NURCAHYO W., PRIYOWIDODO D., BUDIATI R.T., SUSANTI D.S.R. Genetic and parasitological identification of *Trypanosoma evansi* infecting cattle in South Sulawesi, Indonesia. *Veterinary World*, **14** (1), 113, **2021**.
 34. OKELLO I., MAFIE E., EASTWOOD G., NZALAWAHE J., MBOERA L.E. African animal trypanosomiasis: A systematic review on prevalence, risk factors and drug resistance in sub-Saharan Africa. *Journal of Medical Entomology*, **59** (4), 1099, **2022**.
 35. CHAU V.V.N., CHAU B.L., DESQUESNES M., HERDER S., LAN P.H.N., CAMPBELL J.I., BAKER S.A. clinical and epidemiological investigation of the first reported human infection with the zoonotic parasite *Trypanosoma evansi* in Southeast Asia. *Clinical Infectious Diseases*, **62** (8), 1002, **2016**.
 36. ELHAIG M.M., SALLAM N.H. Molecular survey and characterization of *Trypanosoma evansi* in naturally infected camels with suspicion of a Trypanozoon infection in horses by molecular detection in Egypt. *Microbial pathogenesis*, **123**, 201, **2018**.
 37. SELIM A., ALAFARI H.A., ATTIA K., ALKAHTANI M.D.F., ALBOHAIRY F.M., ELISOHABY I. Prevalence and animal level risk factors associated with *Trypanosoma evansi* infection in dromedary camels. *Scientific Reports*, **12**, 8933, **2022**.
 38. KYARI F., MBAYA A.W., BIU A.A., ADAMU L., DENNIS O.O. Seroprevalence of *Trypanosoma evansi* in camels using CATT/T. *evansi* technique in Borno and Yobe states, Nigeria. *Parasite Epidemiology and Control*, **13**, e00209, **2021**.
 39. MOHAMED M.E.A., MOHAMED Y.O., ELSHAFIE E.I., JABIR ALI ALHARBI Y., AL-MEKHLAFI H.M. Molecular detection of *Trypanosoma evansi* in camels (*Camelus dromedarius*) in southwestern Saudi Arabia. *The Thai Journal of Veterinary Medicine*, **49** (1), 93, **2019**.
 40. UZCANGA G.L., PÉREZ-ROJAS Y., CAMARGO R., IZQUIER A., NODA J.A., CHACÍN R., BUBIS J. Serodiagnosis of bovine trypanosomosis caused by non-tsetse transmitted *Trypanosoma (Duttonella) vivax* parasites using the soluble form of a Trypanozoon variant surface glycoprotein antigen. *Veterinary Parasitology*, **218**, 31, **2016**.
 41. ALFALEH F.A., ELHAIG M.M. Molecular prevalence, associated risk factors and genetic characterization of *Trypanosoma evansi* in camels. *Microbial pathogenesis*, **175**, 105967, **2023**.
 42. BARGHASH S.M., DARWISH A.M., ABOU-ELNAGA T.R. Molecular characterization and phylogenetic analysis of *Trypanosoma evansi* from local and imported camels in Egypt. *Journal of Phylogenetics and Evolutionary Biology*, **4**, 18, **2016**.
 43. SARKHEL S.P., GUPTA S.K., KAUSHIK J., SINGH J., SAINI V.K., KUMAR S., KUMAR R. Intra and inter species genetic variability of transferrin receptor gene regions in *Trypanosoma evansi* isolates of different livestock and geographical regions of India. *Acta Parasitologica*, **62** (1), 133, **2017**.