ONLINE PUBLICATION DATE:

Original Research

Analysis of the Land Ecological Security Pattern in the Zhejiang Urban Circle under Small Sample Scenarios

Qing He¹, Xiong Zou¹, Cheng Zhang^{2*}

¹School of Law, Fuzhou University, Fuzhou 350108, China 2School of Business, Taizhou University, Taizhou, 318000, China

Received: 27 January 2024 Accepted: 5 June 2024

Abstract

With continuous socioeconomic development, the land ecological security pattern is becoming increasingly severe. In this article, an evaluation system for land ecological security patterns is constructed based on existing research. Python's TensorFlow framework is used to construct a support vector machine (SVM) based on the land ecological security evaluation method. Finally, based on a small sample scenario, the land ecological security pattern of the urban agglomeration in Zhejiang Province is evaluated and analyzed, and optimization strategies are proposed. The results indicated (1) from 2015 to 2021, the land ecological security levels of most cities in Zhejiang Province gradually decreased and were generally low, with most cities moving from relatively safe levels to critical security levels; (2) significant spatial changes occurred in the overall land ecological pattern of the Zhejiang urban agglomeration, which changed from "higher in the north and south, lower in the middle" to "lower in the southwest, slightly higher in other areas, and extremely low in a few areas (Huzhou city, Zhoushan city)"; and (3) considering social indicators, the land ecological security pattern in the urban ecosystem of Zhejiang Province primarily deteriorated because the per capita cultivated area and the proportion of cultivated land in the natural indicators significantly decreased, and the per capita residential and urban construction land areas significantly increased.

Keywords: land ecological security pattern, Zhejiang urban circle, support vector machine, optimization strategy

Introduction

Land is a precious resource that humans rely on for survival, and all human production and living activities utilize land resources [1]. However, with the rapid socioeconomic development of China, the ecological landscape is becoming increasingly severe. With the requirements of the "Five in One" development strategy and sustainable development, as well as the continuous promotion of the integration of urban agglomerations into the construction of a "two-oriented society" process [1, 2], improving and addressing land ecological security issues has received widespread attention [3]. Therefore,

^{*}e-mail: 201910002@fzu.edu.cn

as the first step in solving this problem, evaluating and analyzing the land ecological security patterns

reasonably is particularly important. Due to the late start in exploring land ecological security evaluation research in China, this research is still in the exploratory stage [4]. Existing research has primarily constructed an urban land ecological security evaluation system considering three dimensions: natural factors, economic factors, and social factors [5-8]. Reference [9] revealed the differentiation information of indicators by establishing a matter element model and evaluating the ecological level of urban land using empirical research methods. Zhang constructed an evaluation index system consisting of a target layer, project layer, factor layer, and indicator layer from three perspectives: resource and environmental pressure, resource and environmental conditions, and human environment response. They subsequently established a comprehensive model for analyzing cities such as Beijing and Shanghai [10]. In recent years, with the continuous development of machine learning, methods integrating land ecological security and machine learning have also received increased attention from scholars. For example, in reference [11], a BP neural network was used in the machine learning framework Keras to evaluate the land ecological security level of the Wuhan urban agglomeration from 2010 to 2017. Based on these findings, ArcGIS software was used to spatially identify the evaluation results, and corresponding land ecological security optimization strategies were proposed.

However, existing research methods often involve simply introducing machine learning methods without identifying or selecting machine learning methods that are suitable for land ecological security evaluation. The available sample size for learning in the land ecological security field is small, which is not suitable for most machine learning methods, and relatively little research has been conducted on Zhejiang Province [12-15]. Therefore, based on existing research, in this article, evaluation indicators that include naturalness, economy, and sociality are first constructed. Then, the advantages and disadvantages of the three types of machine learning methods are compared, and a support vector machine suitable for small samples with high data dimensions is selected as the machine learning method. Finally, the land ecological security pattern of the urban agglomeration in Zhejiang Province is evaluated and analyzed, and optimization strategies are proposed.

Methods

Construction of the Evaluation Index System

"Land Ecological Security" refers to the ability of land-based ecosystems to maintain their structural and functional integrity in the face of environmental changes and human impacts. This concept emphasizes the importance of sustaining land health and productivity to ensure that essential ecological processes, such as nutrient cycling, soil formation, and habitat provision, are preserved. It encompasses strategies to mitigate risks such as soil degradation, contamination, and habitat loss, aiming to protect and enhance the resilience of land areas so they can continue to support biodiversity, agricultural productivity, and human well-being. Based on this definition, the urban land ecosystem is a complex system of interactions between various biotic and abiotic elements in urban areas that include the mutual influences of natural ecology and human construction. These influencing factors are mainly categorized into natural factors, social factors, and economic factors, which are coupled with, influence, and provide feedback to each other, forming a complex urban land ecosystem. Therefore, the evaluation indicators selected in this article can be divided into natural evaluation indicators, social evaluation indicators, and economic evaluation indicators. Compared with the original index selection scheme, this paper uses the Construction Indicators of Ecological Counties, Ecological Cities, and Ecological Provinces (Revised Draft) and the Series Standards of National Garden Cities as guidance and directly selects the indicators of the Chinese government when assessing the level of land ecological security. Based on this, it can be better to provide a more comprehensive understanding of the drivers that affect land ecological security in urban environments, and the final selected indicators are shown in Table 1.

Data Collection and Preprocessing

To analyze the spatiotemporal pattern of land ecological security, data samples must be sampled in both spatial and temporal dimensions. To ensure that the temporal and spatial dimension indicators show significant changes, the temporal sampling interval is selected to be 2 years in this paper, and the spatial dimension is divided by the actual administrative regions under its jurisdiction.

Due to the large number of indicators mentioned above, which come from different fields and have different dimensions, to eliminate the influence of dimensions for unified analysis, the maximum value is used as the standard for data normalization. Notably, to enable the analysis of the spatiotemporal distribution pattern of soil ecological security, the maximum value selection must include indicator data from the entire year. The specific formula is shown in Equation (1):

$$X_i^* = \frac{X_i}{X_{\max}} \tag{1}$$

In this equation, X_i^* represents the normalized indicators, Xi represents the data before normalization, and Xmax represents the maximum total sample value.

Col	Anuiys	is of in	e Lunu	Ecologica	l Security			3				
JOL	Table 1. Summary of evaluation metric selection.											
uth	Form				Reason	Specific indicators						
hor Copy • Author Copy • Author Copy • Author Copy • Author Copy	ui Naturalness		urba an	n land use on d systematica nd, greening	tors evaluate the impact of ecosystems scientifically illy, mainly considering rate, and overall ecological vironment.	Per capita arable land, proportion of arable land area, green coverage rate in built-up areas, forest coverage rate, industrial wastewater discharge, industrial smoke (powder) dust discharge, waste treatment rate, harmless garbage treatment rate, and centralized sewage treatment rate						
uthor			resilie	nce of urban	valuate the adaptability and land ecosystems to human ies and development.	Population density, natural population growth rate, employment rate, urbanization level, per capita living area, per capita water resources, and proportion of urban construction land						
Copy • A	Economy		of	urban land ec pment and tra	tors examine the impacts cosystems on economic ack corresponding feedback echanisms	Per capita GDP, disposable income of rural residents, proportion of tertiary industry, per capita GDP, energy consumption per 10000 yuan of GDP, total amount of road freight transportation						
uthor (Table 2.	Learnin	g sample	classificatior	15.							
IA.	Level	Evaluation value range		Security level	Ecologie	Sample data	Evaluation data					
or Cop)	1	[0.8	-1.0]	Safe	biodiversity, and fully fund land's resilience to disturb	tions with intact ecosystems, high ctioning ecological processes. The pances is very high, ensuring long- sustainability	Maximum value	1.0				
y • Auth	2	[0.6	-0.8)	More secure	Though minor disturband are healthy with good bi processes. The land is well fi	Third quartile	0.7					
1or Cop	3	[0.4	-0.6)	Critical safe	Immediate measures mig Biodiversity and ecologica	increasingly susceptible to risks. ht be required to prevent decline. al functions are maintained but are from potential stresses	Second quartile	0.5				

Less

secure

Unsafe

Moreover, to correctly evaluate the land ecological

security pattern, in this article, land ecological security

registration is divided into five levels, and 21 indicators are selected. The samples were manually divided and

constructed based on the quartiles of each indicator.

The minimum value corresponds to the unsafe level, the

first quartile corresponds to the relatively unsafe level,

the second quartile corresponds to the critical safe level,

the third quartile corresponds to the relatively safe level,

and the maximum value corresponds to the security

expansion is needed for the samples. Because the

quantiles of the sample data are not simply linearly

related, segmented interpolation is used to expand

Table 2 shows that the constructed learning samples contain only five pieces of data. To meet the sample size requirements of machine learning, interpolation

level. The specific details are shown in Table 2.

restoration efforts to regain any ecological functionality. the learning sample size from 5 to 50. Segmented interpolation [16] is a method adopted to mitigate the disadvantages of higher-order interpolation polynomials. The interpolation interval is divided into several subintervals, and a low-degree interpolation polynomial is constructed for each subinterval. This approach yields a better approximation, especially as the number of nodes gradually increases and the interpolation curve becomes more stable at the edges. At this point, the learning samples have been constructed.

Ecosystems are degraded with reduced biodiversity and

diminished ecological functions. They are vulnerable to

disturbances and the environmental resilience is low. Active

restoration efforts are necessary to prevent further decline These are severely degraded landscapes where ecological processes are disrupted or nearly non-functional. Biodiversity

is critically low or absent, and the land requires extensive

Comparison of Machine Learning Methods

In the field of machine learning, support vector machines (SVMs), random forests, and neural networks are widely used in data modeling and classification tasks. These three methods represent different learning

0.3

0.0

First quartile

Minimum

value

4

5

[0.2-0.4)

[0.0-0.2)

paradigms and algorithm strategies, and each has unique advantages and disadvantages [17].

Neural networks [18-20] simulate the structure of human brain neurons and have strong learning ability and adaptability. Neural networks perform well when handling complex nonlinear relationships, but their adaptability to small sample sizes and high-dimensional datasets is relatively weak. Neural networks require a large amount of training data, and selecting and adjusting the model parameters is relatively cumbersome. The random forest algorithm is an ensemble learning-based method that improves the robustness and generalizability of a model by constructing multiple decision trees and synthesizing their prediction results. The random forest algorithm is suitable for processing high-dimensional data and large-scale datasets and has high efficiency during training [21].

SVMs are renowned for their powerful classification performance in high-dimensional spaces. In SVMs, an optimal hyperplane is identified between data points to effectively separate different categories, as shown in Fig. 1. SVMs perform well in small sample scenarios and can map data to high-dimensional space through kernel functions to achieve linear separability, giving them good generalizability in nonlinear high-dimensional feature scenarios [22-24].

In summary, considering that 22 indicators were selected for the analysis of land ecological security patterns and given the high feature dimensions, only 5 effective source data points, and only 50 data points after interpolation expansion, the ability of SVMs to process small samples of high-dimensional feature data highly aligns with the needs of the scenario in this article. Therefore, in this article, SVM is chosen as the machine learning method for subsequent research.

Analysis

Introduction to the Analysis Area

Zhejiang Province is located in the southeastern coastal region of China and belongs to the subtropical monsoon climate zone, covering an area of approximately 109000 square kilometers. Its flat terrain, well-developed water system, and abundant




Fig. 1. Principle diagram of the SVM.

natural resources effectively support transportation and economic development. Zhejiang Province governs 11 prefecture-level administrative units, including Hangzhou, Wenzhou, Ningbo, and Jinhua.

As one of China's strong economic provinces, Zhejiang has become an important center of the Chinese economy due to its large economic output and sustained and stable economic growth rate. Social development in Zhejiang is showing a trend of diversification and openness, with a high level of urbanization and close economic connections forming between cities. This development trend has attracted many people to Zhejiang for development. As of the end of 2021, the total population of Zhejiang Province had reached approximately 53 million.

However, with the rapid advancements of industrialization and urbanization, excessive land development and pollution have become the main factors restricting sustainable development. The decreasing agricultural land area and increasing ecosystem damage and soil pollution, among other issues, are becoming increasingly prominent, posing major challenges to land ecological security. Therefore, an in-depth analysis of the ecological security patterns in the region must be urgently conducted. This analysis helps to comprehensively illustrate the importance of rational utilization of land resources and healthy maintenance of ecosystems and provides a scientific basis for formulating sustainable development strategies.

The relevant data in this article is sourced from official reports such as the Zhejiang Statistical Yearbook, the China Urban Statistical Yearbook, the 14th Year Plan for Energy Development in Five Zhejiang Province, and the Zhejiang Low Carbon Development Report. Some of the data are sourced from the official websites of various statistical bureaus in Zhejiang Province. To form the basic data of this article, the above sources are summarized, and indicators are extracted.

SVM Multi-Classification Model Structure

SVM is usually a binary classification algorithm, but it can be roughly extended to multiclassification problems. In this paper, the SVM multiclassification OAA (One Against-ALL) method, which has high recognition accuracy in small sample environments, is adopted. The OAA method includes two main steps: First, a binary classifier is trained for each category, treating that category as a positive example and all the other categories as negative examples. During testing, the category corresponding to the classifier with the highest decision value is selected as the final classification result. The multiclassification problem is then converted into multiple binary classification problems. Because land ecological security is classified into five levels in this article, five SVM binary classifiers must be constructed with the same structure but different data labels.

The key parameters of the SVM classifier are set as follows:

(1) Kernel function: Commonly used kernel functions include the linear kernel function, polynomial kernel function, and radial basis function (RBF). In nonlinear situations, the best-performing RBF kernel function is typically used. The corresponding kernel function is represented as follows:

$$K(Z_i, Z_j) = \exp\left(-\frac{\left\|Z_i - Z_j\right\|}{2\sigma^2}\right)$$
(2)

In the equation, Z_j and Z_i are input vectors, and σ is the Gaussian function standard deviation.

(2) Penalty parameter C: Parameter C controls the degree of punishment for misclassification. A smaller C value can lead to smoother decision boundaries and tolerate more misclassification; this is suitable for situations with high data noise. A larger C value forces the model to classify each sample as accurately as possible, which may lead to overfitting. In this article, the optimal C value of 0.1 is selected through cross-validation.

(3) Hyperparameters: SVM uses an optimization algorithm called sequential minimum optimization (SMO). In contrast to traditional neural network training, standard SVM libraries and implementations do not require setting parameters similar to the number of training iterations and batch size in neural networks. Personalized configurations must be constructed only for hyperparameters similar to gradient descent, including the learning rate and number of iterations. In this article, the default hyperparameter values are adopted.

Results

The Temporal Trend of the Land Ecological Security Pattern in Zhejiang Province

In this article, the TensorFlow framework in Python is used to construct an SVM multiclassifier, with learning samples used as training samples for SVM multiclassifier training. 22 indicator vectors from various cities in Zhejiang Province in 2015, 2017, 2019, and 2021 were used as data inputs to determine the land ecological security level of the Zhejiang urban agglomeration, as shown in Table 3.

As shown in Table 3, the land ecological security level did not significantly change in most cities from 2015 to 2017; however, in 2019, the land ecological security levels of all cities except Jinhua City and Taizhou City decreased. In 2021, the land ecological security levels significantly changed only in Jiaxing City. Overall, from 2015 to 2021, the land ecological security level in Jiaxing City first increased, then increased, and finally recovered to a relatively safe level. The security levels of Taizhou City and Jinhua City significantly improved from 2015 to 2021, increasing by one level and reaching the critical safe level. The security levels of all other cities decreased by one level, and the land ecological security levels were generally low. The land ecological security levels of Huzhou City and Zhoushan City even decreased to the least safe level.

The Spatial Distribution Trend of Land Ecological Security Patterns in Zhejiang Province

To analyze the spatial distribution trend of the land ecological security pattern in Zhejiang, the evaluation results in Table 3 were combined with a spatial distribution map of the land ecological security pattern in Zhejiang, as shown in Fig. 2.

Tab	ole 3.	Summary	of	evalu	ation	metric	selection	n.
-----	--------	---------	----	-------	-------	--------	-----------	----

City	2015	2017	2019	2021	
Hangzhou City	4	4	3	3	
Ningbo City	2	2	3	3	
Wenzhou City	4	4	3	3	
Jiaxing City	4	4	3	4	
Huzhou City	2	2	1	1	
Shaoxing City	4	4	3	3	
Jinhua City	2	2	3	3	
Quzhou City	4	4	2	2	
Zhoushan City	2	2	1	1	
Taizhou City	2	2	3	3	
Lishui City	2	2	2	2	




Fig. 2. Spatial distribution of land ecological security patterns in Zhejiang Province.

The graph shows that from 2015 to 2017, for the north-south pattern, the land ecological spatial security level in northern Zhejiang Province was greater than that in central and southern Zhejiang Province, excluding Huzhou City in the northernmost part and Wenzhou City in the southernmost part. Similarly, for the east-west pattern, the land ecological spatial security level of the western region of Zhejiang Province was slightly greater than that of the eastern region. From 2019 to 2021, the land ecological space security level in the southwest was lower than that in the central, southern, and northern regions, except for Huzhou, while in the easternmost part, the level in Zhoushan city was generally lower than that in the other regions.

Reasons for Changes in the Ecological Security Pattern of Land in Zhejiang Province

To analyze in depth the reasons for the changes in the land ecological security pattern of the Zhejiang urban agglomeration, the change trends are examined using naturalness, economy, and sociality indicators. First, the land ecological security level change trends corresponding to the three categories of indicators are trained based on the SVM model, as shown in Table 4.

The results in Table 4 show that the naturalness index is strongly correlated with the overall land ecological security level, and the spatiotemporal distribution patterns of the two indices are consistent; that is, the change in the naturalness index is the dominant factor in the changes in the overall land ecological security level. The source data collected in this article indicate that between 2015 and 2021, the per capita arable land area and the proportion of arable land area in most cities with declining ecological security levels for the land significantly decreased, while the emissions of industrial wastewater and industrial smoke (powder) dust slowly increased.

The trend of changes in the economic indicators shows that all cities are slowly improving, which is inconsistent with the distribution pattern of the overall land ecological security level. That is, the impact of economic indicators on the overall land ecological security level is relatively small. Analyzing the source data collected in this article reveals that from 2015 to 2021, the per capita GDP, disposable income of rural residents, proportion of the tertiary industry, per capita GDP, and total road freight volume in economic indicators increased annually, while the trend of energy consumption per 10 thousand yuan of GPD was consistent with the trend of changes in land ecological security levels.

The trend of changes in social indicators shows that except for Jinhua and Shaoxing, the distributions of land ecological security levels in other cities are relatively consistent, indicating that social indicators have a certain impact on the overall land ecological security level. Analyzing the source data collected in

6

Table 4. Trends of changes in three major indicators.												
City	Naturalness indicators				Economic indicators				Social indicators			
City	2015	2017	2019	2021	2015	2017	2019	2021	2015	2017	2019	2021
Hangzhou City	4	4	3	3	2	2	2	4	4	4	4	3
Ningbo City	2	2	3	3	2	2	2	4	4	4	4	4
Wenzhou City	4	4	3	3	2	2	2	4	3	4	3	3
Jiaxing City	4	4	3	4	2	2	3	4	2	2	3	3
Huzhou City	2	2	1	1	3	3	3	4	2	2	2	2
Shaoxing City	4	4	3	3	2	2	2	4	2	2	3	3
Jinhua City	2	2	3	3	2	2	2	4	3	2	3	3
Quzhou City	4	4	2	2	3	3	3	4	2	2	2	2
Zhoushan City	2	2	2	2	4	4	4	4	3	3	3	3
Taizhou City	2	2	3	3	2	2	2	4	2	2	2	2
Lishui City	2	2	2	2	3	3	3	4	2	2	2	2

Table 4. Trends of changes in three major indicators.

this article showed that the per capita water resources in social indicators are positively correlated with the land ecological level but strongly negatively correlated with the proportion of urban construction land and per capita residential area.

Conclusions and Optimization

Conclusions

In this article, a land ecological security pattern evaluation system that includes natural, economic, and social indicators is constructed. Using the TensorFlow framework in Python, an SVM multiclass trainer is designed, and the learning samples are interpolated and expanded to serve as training input. Finally, the model is used for land ecological security pattern evaluation in the Zhejiang urban agglomeration. The conclusions of this paper are as follows:

(1) The land ecological security level in the Zhejiang urban agglomeration generally declined between 2015 and 2021, and the land ecological security levels in most cities were relatively low, mostly at the critical safe level.

(2) The land ecological security pattern of the Zhejiang urban agglomeration has changed from the original pattern of "higher in the north and south, lower in the central level" to the current pattern of "lower in the southwest, slightly higher in other areas, and extremely low in a few areas (Huzhou city, Zhoushan city)".

(3) The natural indicators, some social indicators such as per capita water resources and per capita living area, and the economic indicator of energy consumption per 10000 yuan of GDP are important factors in land ecological security pattern changes.

Optimization Strategy

(1) Improving the legal system: Improving the legal system is the primary task for ensuring land ecological security. The government should formulate and strengthen relevant regulations, regulate land use behavior, and increase the costs of illegal activities. By revising laws such as the Land Management Law and the Environmental Protection Law, punishments for illegal development and damage to the ecological environment can be made more severe. Moreover, these regulations should be more strongly enforced to ensure the laws are effectively implemented. In addition, social participation should be established, and the breadth and depth of supervision should be improved.

(2) Enhancing public awareness: Public environmental awareness participation and are crucial for ensuring land ecological security. The government should take various measures, such as conducting environmental protection publicity and education activities, strengthening media publicity, and increasing public awareness of the importance of land ecology. Ecological civilization education in schools should be strengthened, and students' environmental awareness should be cultivated. Moreover, community residents should be encouraged to actively participate in environmental protection activities; volunteers should be organized to conduct public welfare work such as afforestation and ecological restoration; and an atmosphere of joint participation in protecting the ecological environment should be created across overall society.

(3) Establishing warning mechanisms and scientifically formulating land use plans: To detect and respond to potential ecological risks in a timely manner, a sound land ecological security warning mechanism should be established. Advanced technologies such as remote sensing and geographic information systems can be introduced to monitor and evaluate the ecological status of land in real time, and to provide early alerts for potential problems. Moreover, the government should scientifically formulate land use plans, reasonably plan urban expansion and rural construction, and mitigate the adverse effects of excessive development on the ecological environment. Community planning standards should be developed to ensure that land resources are rationally used while preserving public spaces such as green spaces and parks and improving the ecological integrity of urban communities. We suggest strengthening crossdepartmental cooperation, forming a comprehensive land use plan, and ensuring coordination between ecological needs and economic development.

Acknowledgments

The general project of the National Social Science Fund "Study on the Rule Perfection and Arrangement of Environmental Credit Evaluation System of Chinese Enterprises" (23BFX099)

Conflict of Interest

The authors declare no conflicts of interest.

References

- ZOU C., YE X., YANG S. Construction of ecological security patterns based on ecological protection redlines in Jiangxi Province, China. In 2018 IEEE International Geoscience and Remote Sensing Symposium, IEEE: Valencia, Spain, 7468, 2018.
- CHEN S. Quantum computer assisted dynamics modeling for the unified configuration analysis of cultivated land and ecological land with data classification algorithms. In 2022 4th International Conference on Smart Systems and Inventive Technology (ICSSIT), IEEE: Tirunelveli, India, 1531, **2022**.
- YOU Y., SUN J., YAN H., JIANG J. Ensemble belief rule-based model for ecological security of land resource evaluation. In 2019 5th International Conference on Big Data and Information Analytics (BigDIA), IEEE: Kunming, China, 108, 2019.
- LI Q.S., LI J. Evaluation of land ecological security in Wuhan city circle based on principal component analysis. Wuhan Vocational College of Technology, 19 (06), 96, 2020.
- CHEN Z., MA L., GUO F., CHEN Y. Internet of things technology in ecological security assessment system of intelligent land. IEEE Access, 8, 99772, 2020.
- YUAN S.F. Study on regional land ecological security evolvement: a case of Hangzhou. In 2011 Seventh International Conference on Natural Computation, IEEE: Shanghai, China, 4, 2278, 2011.
- 7. ZHU J. Evaluation of land ecological security based on set pair analysis. In 2010 International Conference on

Intelligent Computation Technology and Automation, IEEE: Changsha, China, 2, 233, 2010.

- ZHU J. An application of attribute recognition theory in land ecological security evaluation a case study on areas along the Changjiang River in Anhui Province. In 2010 International Conference on E-Health Networking Digital Ecosystems and Technologies (EDT), IEEE: Shenzhen, China, (1), 278, 2010.
- LUO W.B., WU C.F., WANG Y.J. Evaluation on urban land ecological level based on matter element analysis: a case of Hangzhou City in Zhejiang Province. China Land Science, 22 (12), 31, 2008.
- ZHANG L.R. Early warning research of wuhan urban agglomeration land ecological security. Huazhong Agricultural University, 1, 2012.
- JIANG C., LUO Z., ZHAO J., ZHAO Y. Dynamic evaluation of land ecological security in Jiangxi Province based on DPSIR-TOPSIS model. Jiangsu Agricultural Sciences, 46 (17), 294, 2018.
- SU S., CHEN X., DEGLORIA S.D., WU J. Integrative fuzzy set pair model for land ecological security assessment: a case study of Xiaolangdi Reservoir Region, China. Stochastic Environmental Research and Risk Assessment, (1), 98, 2009.
- RU X., JIANG T., WU H., PENG L. Land use/cover change analysis and evaluation of ecological carrying capacity in urban ecological environment. In 2011 International Conference on Remote Sensing, Environment and Transportation Engineering, IEEE, 4141, 2011.
- TANG Y. Regional land resource eco-security evaluation. In 2011 International Symposium on Water Resource and Environmental Protection, IEEE: Xi'an, China, 2820, 2011.
- 15. YING J., CUI H.S., SONG X.F. Evaluation of land ecological security in Hubei Province. In 2013 International Conference on Management Science and Engineering 20th Annual Conference Proceedings, IEEE: Harbin, China, 2236, 2013.
- SASAKI Y., YU J., ISHIKAWA Y. Road segment interpolation for incomplete road data. In 2019 IEEE International Conference on Big Data and Smart Computing (BigComp), IEEE: Kyoto, Japan, 1, 2019.
- LI H. Computer network connection enhancement optimization algorithm based on convolutional neural network. In 2021 International Conference on Networking, Communications and Information Technology (NetCIT), IEEE: Manchester, UK, 281, 2021.
- HONGO S., ISOKAWA T., MATSUI N., NISHIMURA H., KAMIURA N. Constructing convolutional neural networks based on quaternion. In 2020 International Joint Conference on Neural Networks (IJCNN), IEEE: Glasgow, UK, 1, 2020.
- GAO K., ZHANG Q., YU L., HUO L. A muti-branch convolutional netural network for mobile platform. In 2019 34rd Youth Academic Annual Conference of Chinese Association of Automation (YAC), IEEE: Jinzhou, China, 357, 2019.
- 20. SHI X., LI L. Loop closure detection for visual SLAM systems based on convolutional netural network. In 2021 IEEE 24th International Conference on Computational Science and Engineering (CSE), IEEE: Shenyang, China, 123, 2021.
- LATIFAH A.L., SHABRINA A., WAHYUNI I.N., SADIKIN R. Evaluation of random forest model for forest fire prediction based on climatology over Borneo. In 2019 International Conference on Computer, Control,

Informatics and its Applications (IC3INA), IEEE: Tangerang, Indonesia, 4, **2019**.

- 22. LI J., ZHAO Y., GAO X., ZHAO W. Feature extraction and SVM parameters optimization based on ICQPSO. In 2020 International Conference on Computer Network, Electronic and Automation (ICCNEA), IEEE: Xi'an, China, 329, 2020.
- CHARAN K.S., CHOKKALINGAM S.P., SUNDARI K.K.S. Efficiency of decision tree algorithm for brain tumor MRI images comparing with SVM algorithm.

In 2022 14th International Conference on Mathematics, Actuarial Science, Computer Science and Statistics (MACS), IEEE: Karachi, Pakistan, 1, **2022**.

24. DUTTA K.K., SWAMY S.A., BANERJEE A., RASHI B.D., CHANDAN R., VAPRANI D. Kannada character recognition using multi-class SVM method. In 2021 11th International Conference on Cloud Computing, Data Science & Engineering (Confluence), IEEE: Noida, India, 405, 2021.