

Original Research

Spatial Heterogeneity and Determinants of Soil Organic Carbon Density Across Varied Karst Landscapes in Southwest China

Jiang Hong¹, Wen Hu², Hefeng Wan¹, Xianfei Huang^{3, 4*}

¹Guizhou Academy of Sciences, Guiyang 550005, China

²College of Resources and Environment Engineering, Guizhou University, Guiyang 550025, China

³Institute for Forest Resources and Environment of Guizhou, Key Laboratory of Forest Cultivation in Plateau Mountain Areas, College of Forestry, Guizhou University, Guiyang 550025, China

⁴Guizhou Provincial Key Laboratory for Environment, Guizhou Normal University, Guiyang 550025, China

Received: 3 March 2024

Accepted: 9 July 2024

Abstract

This study examined the influence of environmental factors on soil organic carbon density (SOCD) in five regions of southwest China, including karst canyon, karst peak cluster depression, karst plateau, karst trough valley, and non-karst areas. Significant differences in SOCD were found across these landscapes. The study quantitatively analyzed the vertical distribution and key drivers of SOCD in these landscapes. Specifically, soil organic carbon density value in soil thickness of 0 to 0.40 m is: karst trough valley>karst peak cluster depression>karst plateau>karst canyon>non-karst regions; soil organic carbon density value in soil thickness of 0.40 to 1.00 m is: karst plateau>karst trough valley>karst peak cluster depression>karst canyon>on-karst regions. Soil organic carbon density in non-karst regions is markedly positively related to slope gradient, slope aspect, and slope position and negatively correlated with land use and soil thickness. The study also found that the main factors influencing SOCD differ by region. SOCD was influenced by soil thickness in karst canyons, karst peak cluster depressions, and karst plateaus. SOCD was influenced by slope aspects in karst trough valleys.

Keyword: topographical features, organic carbon density, karst, non-karst, influence factors

Introduction

Karst is a term that refers to a special landscape containing caves and an extensive underground water system [1], and it is one of the most ecologically fragile landform types with a very fragile geological setting

and an intense karst process [2, 3]. In this region, carbonate rocks are usually dissolved under the work of acidic waters, which are derived from CO₂ present in the soil and in the air, forming complex above-underground geographic structures [4, 5]. Therefore, studies on resources and pollutants are facing huge challenges, and some studies have been carried out during the past two decades [6, 7]. With the growing development intensity of humans exploiting nature, population pressure and unreasonable land use changed the landscape structure

*e-mail: hxfswj@gznu.edu.cn

of the karst region, accelerating the emergence of ecological and environmental problems in the region such as soil degradation, fertility depletion, land desertification, and biodiversity reduction, and caused a prominent contradiction between ecological protection and social and economic development [8]. The fragility of the karst ecosystem determines its vulnerability and difficult recovery from desertification [9]. Research focused on a realistic strategy in the karst area has been widely concerned. It is a very prominent environmental issue of ecological construction in western China and also one of the major obstacles of karst regions in southwestern China in sustainable development [10].

As the center of the karst region in southern China, the topography of Guizhou province is more complex compared with other provinces, including at the same time mountainous regions, hills, basins, and even high mountains, valleys, and so on [11]. Among them, the Guizhou karst landform is the most representative [12]. Karst landform in the province can be divided into four types; karst canyon, karst peak cluster depression, karst plateau, and karst trough valley [13]. The total area of the karst landform is 109084 km² or 62% of the total land area. Strong water erosion and interstitial, variable, and tilted surface uplift of earth crust and complex geological process of the epigenetic zone, since Neogene in rock desertification processes form rich and colorful karst landform in Guizhou and soil conservative water, fertilizer, and the productivity of land, differ in different karst landforms. Thus, the carbon cycle characteristics are different [14, 15].

Soil carbon pool is the biggest carbon pool in terrestrial ecosystem carbon pools, and SOCD is an important indicator in evaluating soil organic carbon condition as well as the core content of global carbon cycle research. SOCD has important relations to nutrient supply and prevention of soil erosion [16, 17]. Carbon fixed during photosynthesis in plants enters the soil organic carbon pool through plant residue decomposition and soil microorganisms metabolic processes. Accumulating soil organic carbon enhances soil structure, sustains soil fertility, and aids in reducing atmospheric CO₂ levels. The decrease in soil organic carbon storage can directly cause the soil quality decline, appearing as a rapid decrease in the ability of soil to supply nutrients to plants, soil tilth, and air and water permeability [18]. Soil organic carbon content generally decreases with depth, being higher in the surface soil and decreasing with depth. Moreover, soil type, climate, vegetation type, and land use practices are additional factors influencing the distribution of soil organic carbon. For instance, forest soil in humid climates typically contains higher organic carbon, whereas grassland soil has lower levels. The nature of karst rock desertification is that the soil quality changes, mainly in the physical, chemical, and biological properties [19]. Studies on soil ecosystem degradation caused by karst rock desertification by domestic and foreign experts mainly focus on aspects

of the reason for soil rocky desertification, characteristics of soil degradation, the vegetation recovery of degraded ecosystems, etc. [19, 20]. While studies on the carbon cycle in soil and the distribution of soil organic carbon in different karst landforms and rock desertification backgrounds are few [21]. The carbon transfer process in karst systems is dominated and controlled by soil carbon, and SOCD becomes the dynamic mechanism driving and restricting carbon transfer in surface-layer karst systems [22]. These processes are in dynamic change due to changes in soil genetic characteristics and environmental conditions. Therefore, conducting comprehensive research on the spatial variation of soil organic carbon and human influences on it in karst regions is crucial for understanding soil carbon cycling mechanisms and directing land use planning and environmental conservation efforts. Systematically analyzing the spatial distribution of organic carbon and assessing human impacts can establish a scientific basis for future soil management and ecological restoration in karst regions [23].

Through taking SOCD in typical karst ecosystems and non-karst ecosystems in southwest China as research subjects. This research discussed the distribution characteristics of organic carbon density and its internal connection with karst topography and soil environmental factors. The distribution pattern of SOCD in karst ecosystems and its driving mechanism were illuminated. The response regularity and its internal mechanism of SOCD in the degradation, recovery, and evolution processes of the karst rock desertification ecosystem were illuminated. We also provided references for the reconstruction of rock desertification ecosystems and coping with the source reduction and sinks of carbon cycles in global climate change.

Materials and Methods

Study Region Overview

This paper chose 4 typical karst regions and 1 non-karst region, 5 research subjects in total. The non-karst region is located in Zunyi County, Zunyi City, Guizhou Province, Study region I, and 4 typical karst regions of different topographical features are respectively located in karst Grand Canyon, Guanling County, Guizhou Province, Study region II, karst peak cluster depression, Libo County in Guizhou Province, Study region III, karst plateau, Puding County, Guizhou Province, Study region IV, karst trough valley, Yinjiang County, Guizhou Province, Study region V. Specific geographical location and basic information were shown in Fig. 1.




Fig. 1. Distribution of studied region and sampling sites.

Zunyi County (non-karst region) is located in the northeast of Yunnan-Guizhou Plateau, with geographic coordinates of $105^{\circ}36'-108^{\circ}13'E$ and $27^{\circ}8'-29^{\circ}12'N$. It belongs to the Baipu tributary area of the Yangtze River basin, with no large river water system in the region. The terrain is high in the northwest and low in the southeast, with the Lou Mountains and its north-south branch as its skeleton. It belongs to the subtropical monsoon climate zone, with an average annual temperature of $14.7^{\circ}C$ and an average annual precipitation of 1200 mm. Subtropical evergreen broad-leaved forest and mixed broadleaf-conifer forest where native vegetation was mostly destroyed and now the remaining are few trees and shrubs, herbs, vines, etc. Guanling County (karst canyon) belongs to the Beipanjiang water system, the upper reaches of the Pearl River. The river valley is deep, and the basin is small in area. The terrain undulation is large, with an elevation of 450 m to 1450 m and a relative height difference of up to 1000 m. The average annual precipitation of the region is 1100 mm, and precipitation is mainly distributed from May to October, accounting for 83% of the total rainfall throughout the year. The vegetation type was similar to Zunyi County (non-karst region). Libo County (karst peak cluster) is located at $107^{\circ}52'10''-108^{\circ}05'40''E$ and $25^{\circ}09'20''-25^{\circ}20'50''N$, belonging to the Longjiang Water System, a tributary of the Pearl River basin. The elevation ranges from 430 to 1078 m, with most of the elevation around 800 m. Except for a very small amount of sand shale exposed in some parts, most parts of the region are the exposed karst peak cluster and peak forest landforms composed of pure dolomite and limestone. The vegetation types were evergreen broad-leaved forests and broad-leaved coniferous forests. And there were also rich in vegetation due to the complete protection of the local area. Study Area IV: The Houzhaihe catchment of Puding County (karst

plateau) covers an area of 81 km^2 , with geographic coordinates of $105^{\circ}40'43''-105^{\circ}48'2''E$ and $26^{\circ}12'29''-26^{\circ}17'15''N$. The elevation at the watershed between the Yangtze River and the Pearl River water systems ranges from 1223.4 to 1567.4 m. There are karst caves, subterranean streams, and other surface types under the ground. The air pressure is between 806.1 and 883.8 hpa. The evergreen conifer forest is an evergreen deciduous broad-leaf mixed forest where the original forest vegetation is reclaimed as agricultural land. Yinjiang County (karst trough valley) is located at $108^{\circ}17'52''-108^{\circ}48'18''E$ and $27^{\circ}35'19''-28^{\circ}20'32''N$, belonging to the Wujiang water system of the Yangtze River basin. The maximum east-west distance is 62.5 km, and the maximum north-south distance is 75.8 km. It is located in the transitional slope zone from Yunnan-Guizhou Plateau to Xiangxi Hill and Sichuan Basin, between the low-mountain hilly area in eastern Guizhou and the middle-mountain canyon in northeastern Guizhou. The highest peak of the Wuling Mountains, Mount Fanjing, is located in its eastern part, forming a terrain that is high in the east and low in the west, sloping from southeast to northwest. The vegetation types were evergreen conifer broad-leaf mixed forest. It belongs to the subtropical warm humid climate zone, with an average annual temperature of $16.8^{\circ}C$ and an annual precipitation of around 1100 mm in Table 1.

Soil Sampling

We sampled in different sampling areas and set 1×1 m quadrates in each area. A five-point sampling method is used to excavate soil profiles in the quadrate, and the soil samples in one quadrate are uniformly mixed into one sample and stored. Soil samples were collected through the layered sampling method. Soil samples were sampled by bottom-up layered sampling of

Table 1. Main overview of the research area.

Study region	Abbreviation	Soil erosion rate
Study region I (non-karst region)	ZYFK	24.30%
Study region II (Karst canyon)	HJDXG	49.60 %
Study region III (Karst peak cluster)	LBFC	69.75 %
Study region IV (Karst plateau)	PDJY	54.30%
Study region V (Karst trough valley)	YJCG	59.32%

the soil profile. The excavation depth of the soil profile was no greater than 100 cm. The soil layer was excavated from the basement rock or parent material to 100 cm underground. It was divided into a total of 10 layers, including 0 to 10 cm, 10 to 20 cm, 20 to 30 cm, 30 to 40 cm, 40 to 50 cm, 50 to 60 cm, 60 to 70 cm, 70 to 80 cm, 80 to 90 cm, and 90 to 100 cm. The background information of each sampling point should be recorded, along with the environmental information such as slope gradient, elevation, aspect, and slope position. A total of 126 sampling points were surveyed and collected, and a total of 968 soil samples were collected [24].

Soil Analysis Methods

The soil samples were naturally air-dried at room temperature and screened at 150 μm . Then, total SOC levels were determined by $\text{K}_2\text{Cr}_2\text{O}_7$ oxidation at 170-180°C followed by titration with 0.10 mol L^{-1} FeSO_4 , and 10% repetition was conducted (Herold et al. 2014).

Soil acreage was calculated using GIS information and field survey data. Bulk density was measured layer by layer from the top to the bottom of the soil profile using the cutting-ring method. Soil thickness was recorded according to the type of ecological niche using an iron measuring rod that was 60 or 120 cm long, based on the soil mass at different depths. The boulder content was surveyed using a linear transect. Due to the existence of complex landscapes in the karst area, the length of the linear transect was set at 10 m. Although more accurate information could be obtained from a longer transect, establishing a longer transect would have required excessive effort. Grid cells with rock coverage were surveyed using a tape measure.

Statistical Analysis

Data were managed and treated with Microsoft Excel 2007 (Microsoft, Redmond, WA, USA). Statistical analysis was performed with the Statistical Package for the R language (R 5.0, IBM, Armonk, NY, USA) and ArcGIS mapping software (ArcMap 10.3, ESRI, Redlands, CA, USA).

Results

Main Soil Properties Under Different Topographical Features

Karst development area has obvious stripes for being controlled by geological structure forming cross distribution of two landforms of karst and non-karst. Due to the common interbedding of carbonate formation and fragmental rocks and the fact that the whole sediment structure is mainly carbonate, carbonate formation is discretely striped in horizontal distribution after fold fracture and denudation erosion. It manifests as two different landforms evolving and developing together and intercrossing with each other. Five study regions represent 5 typical landform regions in the southwest and 4 study regions chosen represent respectively 4 typical landform types in the southwest; karst plateau mountains, plateau basins, plateau canyons, and plateau trough valleys. Specific basic characteristics are shown in Table 1. This study studied the distribution of basic attributes of different karst landforms in the study region scale, and the specific results are shown in Table 2. The basic attributes of 968 soil samples from 126 sampling points in 5 landform types have different variations.

Due to special geological and climatic conditions, the soil environment of the karst region has essential features of bedrock exposure, small soil stock, discrete distribution, complex, diverse microrelief, etc. Thus karst soil thickness, rock outcrops, and gravel content are all higher than those of non-karst regions. The gravel content in different karst landforms shows that karst trough valley > karst canyon > karst peak cluster depression > karst plateau. The soil thickness shows that: karst plateau > karst canyon > karst trough valley > karst peak cluster depression. The gravel content and organic carbon content show that karst peak cluster depression > karst canyon > karst trough valley > karst plateau. The inspection result of multiple comparisons Duncan shows that soil organic carbon contents of different landform rock desertification are obviously different, which shows that karst landform has significant influences on soil organic carbon content ($P < 0.05$).

Characteristics of SOCD Profile in Different Landforms

This research studied the spatial distribution of SOCD in different landforms in the southwest from two aspects of karst and non-karst, and the specific results are shown in Fig. 2. The maximum value of organic carbon density (0-100 cm) in the non-karst region is 61.22 $\text{kg}\cdot\text{m}^{-2}$ and the minimum value is 0.25 $\text{kg}\cdot\text{m}^{-2}$, and the average value (0-100 cm) is 5.61 $\text{kg}\cdot\text{m}^{-2}$ while the maximum value of organic carbon density (0-100 cm) in the karst region is 94.11 $\text{kg}\cdot\text{m}^{-2}$, the minimum value is 1.22 $\text{kg}\cdot\text{m}^{-2}$,

Table 2. Main soil properties in the study area under different geomorphological characteristics.

Research area	Number	Rock outcrops (%)	Soil thickness (cm)	Gravel content in (%)	SOC (g.kg ⁻¹)	SOCD (kg.m ⁻²)
ZYFK	27	6±1.21	81.64±12.34	5.64±1.47	9.53±1.12	4.98±1.23
HJDXG	25	30±2.34	46.93±29.61	13.94±2.45	16.41±2.12	5.61±1.73
LBFC	25	28±3.45	39.08±25.42	15.21±2.32	17.43±2.68	6.34±1.58
PDJY	22	12±1.23	80.26±19.74	11.66±1.56	13.53±1.39	7.58±2.17
YJCG	27	32±1.12	45.22±14.3	13.01±2.3	16.63±1.86	6.57±1.60

and the average value (0-100 cm) is 5.01 kg·m⁻². The characteristics of the SOCD profile in different landforms are shown in Fig. 2. From Fig. 2, it can be seen that the vertical variation features of SOCD in different landforms are different; on the whole, SOCD of the karst region is higher than that of the non-karst region; SOCD content of soil thickness of 0-0.40 m rapidly rises along with soil depth increases; and SOCD value of soil thickness of 0-0.40 m is YJCG>LBFC>PDGY>HJDXG>ZYFK. While soil thickness is 0.40-1.00 m, except SOCD of the karst plateau rises rapidly, SOCD of other types all increases slowly. The characteristics of the SOCD profile in different landforms show that: PDGY>YJCG>LBFC>HJDXG>ZYFK. Through multiple comparisons, Duncan, the inspection results show that SOCD in different karst landforms is obviously different, and SOCD in non-karst regions has no significant difference ($P<0.05$). Thus, it shows that karst landforms have significant influences on soil organic carbon content.

Composite Influence Factors of SOCD in Different Landforms

Influence Factors of SOCD in Non-Karst Region

The redundant analysis (RDA) method is used to analyze soil properties and their interrelation with environmental factors. Using Canoco 5.0 software, first organic carbon content was analyzed by DCA analysis (Detrended Canonical Correspondence Analysis) to generate gradient and length, then linear models (RDA or PCA) were selected using factors to carry out analysis and comparison. Land use was valued according to types of land use, and types of land use were divided into five classes according to the degree of human disturbance: Class 1 mainly includes arbor woodland, arbor and shrub woodland, and shrubby forest, which have the least human disturbance and the assignment value is 1; Class 2 is mainly wasteland with an assignment value of 2; Class 3 is mainly abandoned cultivated land with occasional animal husbandry phenomenon and an assignment value of 3; Class 4 is mainly grassland and




Fig. 2. SOC density of different landscapes under different statistical depths.

(YJCG: karst trough valley; XYFC: karst peak cluster depression; HJXG: karst canyon; PDGY: karst plateau; ZYFK: Non-karst region).

shrub grassland, mainly grazing disturbance, with an assignment value of 4; Class 5 is all kinds of cultivated land, including paddy fields, dry land, garden land, sloping farmland, fruit forest land, where the intensity of human disturbance is the highest with the assignment value of 5. This research adopted RDA sorting analysis of the relation between organic carbon density and the factors. A sorting diagram of relations between soil organic carbon and comprehensive factors in the non-karst region is shown in Fig. 3.

RDA sorting can reflect the influence degree of each influencing factor on the variation of organic carbon density directly. The length of connecting lines of environmental factors represents the correlation between soil organic carbon contents and influence factors and longer lines and smaller included angles in the RDA plot indicate stronger positive correlations between soil organic carbon content and environmental factors. When the included angle is less than 90° , it means the influence factor is positively correlated with the organic carbon content, while it is larger than 90° , it means the influence factor is negatively correlated with the organic carbon content. The smaller the angle included, the higher the correlation between organic carbon and the influence factor. As shown in Fig. 3, SOCD has a significant positive correlation with slope gradient, slope aspect, and slope position ($P < 0.05$). SOCD has a negative correlation with land use and soil thickness. While the influence of rock outcrops on SOCD shows correlations of different degrees in different soil thicknesses. Among them, SOCD (0-30), SOCD (0-10), and SOCD (0-20) were negatively correlated with rock

outcrops, while SOCD (0-100) was positively correlated with rock outcrops. The angle between SOCD (0-100) and slope gradient is the smallest indicating, that slope gradient has the greatest influence on SOCD (0-100), and the angle between SOCD (0-20) and slope aspect is the smallest, indicating that SOCD (0-20) is highly-significant and positively correlated with slope aspect, and the correlation coefficient is the largest ($P < 0.05$). On the whole, rock outcrops, land use, and soil thickness were remarkably negatively correlated with SOCD, indicating that SOCD in the non-karst region decreased gradually with the increase of soil thickness and the human disturbance degree (land use).

Influence Factors of SOCD in Karst Regions of Different Landforms

Landform plays a significant role in spatial distribution and variation of karst SOCD (Fig. 4). The driving factors of SOCD in different landforms are different. In the karst canyon region, SOCD has significant positive correlations with slope aspect, slope gradient, and soil thickness, and the angle between SOCD and slope aspect is the least, indicating the most significant correlation ($P < 0.05$). SOCD has significant negative correlations with slope position, land use, slope position, and rock outcrops. And the influence degree of each influence factor on organic carbon density from high to low is: soil thickness > rock outcrops > slope aspect > slope position > slope gradient > land use ($P < 0.05$). In the karst peak cluster depression region, SOCD has significant positive correlations with soil thickness,




Fig. 3. The relationship between soil organic carbon and geographic factors in non-karst areas.

slope position, and slope aspect. The significant negative correlation with various influence factors on organic carbon density from high to low is: soil thickness>slope gradient>slope position>rock outcrops>land use>slope aspect. In the karst plateau region, SOCD has significant positive correlations with slope gradient, slope position, and soil thickness, with a significant negative correlation with land use, rock outcrops, and slope aspects. The influence degree of influence factor on organic carbon content from high to low is: soil thickness>land use>slope aspect>slope position>slope gradient>rock outcrops. In the karst trough valley region, SOCD has significant positive correlations with land use, and slope position, while significant negative correlations with soil thickness, rock outcrops, slope aspects, and slope gradient. The influence degree of each influence factor on organic carbon content from high to low is: slope aspect>rock outcrops>soil thickness>slope position>slope gradient>land use.

Discussion

Comparison of Organic Carbon Content of Karst Soil and Non-Karst Soil

Soil is a very complex synthesis formed by secular evolution in nature, also the continuous variant of space-time, and has complexity and spatial variability due to soil formation from parent material, terrain, climate, vegetation, and so on, and the interference of human activity [25, 26]. Under the influences of many natural and human factors, such as different topographical features, landform factors, and human disturbances, the physicochemical property of soil shows a patchy or gradient pattern in spatial distribution [27]. The karst ecological environment is complex. Outcrop heterogeneity and diversity of crack structure of rocks under the ground form multi-layer ecological spatial structure. Rock outcroppings and extreme development of cracks caused the spatial heterogeneity of karst soil by making the horizontal and vertical distribution of soil discrete, the soil mantle extremely fractured, and the thickness distribution of soil uneven [28].




Fig. 4. The relationship between SOCD and geographic factors in different karst landforms. Note: (a) is karst canyon; (b) is karst peak cluster; (c) is karst plateau; (d) is karst trough valley.

In particular, there is a large amount of sloping soil in karst regions, and the average depth is only 4 to 20 cm. Therefore, the study of organic carbon and organic carbon density in topsoil (0-20 cm) in the karst region is the basis for evaluating soil carbon sequestration in this area. In different karst landforms, the soil organic carbon content (0-20 cm) was between 10.62 g and 89.11 g·kg⁻¹ (Table 3), and the average soil organic carbon content was 23.07 g·kg⁻¹. The content of soil organic carbon is similar to that of Sanjiang Plain in China and Urasnabala, Tanzania, which is higher than that in the small watershed of the Loess Plateau and lower than that of the Mediterranean Basin [29-31]. The geological landscape of the Loess Plateau is broken, and the low mountains and hills are continuous [32, 33]. Due to the long-term serious soil and water loss, the soil is barren and the soil and water loss is aggravated, resulting in a large amount of soil organic carbon loss, so the organic carbon content in the Loess Plateau is low. The Mediterranean Sea is a typical Mediterranean climate region, with dry heat and less rain in summer and warm and humid winter, which is beneficial to the growth of vegetation and increases the input of carbon sources to the soil, while the climate is dry with little rainfall in karst areas and the soil carbon input is smaller than that in the Mediterranean region. Organic carbon in the soil is the equilibrium result of the amount of plant residues entering the soil and the decomposition loss under the action of soil microorganisms. Its content is controlled by many physical, biological, and human factors, such as the climate, vegetation, soil properties, agricultural management practices, and so on, and there is an interaction between various factors [34].

Soil organic carbon reserve content in southwest China is relatively low. In this study, the organic carbon density of karst soil and non-karst soil is much lower than the average level of organic carbon density in China (9.6 kg·m⁻²). It is also much lower than the average SOC_D in the Loess Plateau region of northwest China and Sanjiang Plain in North China, which reflects that the karst geological environment in southwest China has a profound influence on SOC_D. The main reason is that the soil is dispersed and the soil thickness has obvious differences due to the karst geological environment in southwest China. The average depth of topsoil in

most areas in this study was 5 to 11 cm. Therefore, the topsoil in the karst area (0-20 cm) was studied. The soil carbon density in the study regions is also lower than that of karst in Guangxi, China (9.80 kg·m⁻²), which indicates that the high spatial heterogeneity of soil in karst regions results in large spatial variations of organic carbon density [35]. At the same time, due to the special binary hydrological structure and complex topographic forms such as complex peak cluster depression, hoodoo, funnel, and so on, the soil formation rate is slow, the soil layer is shallow and discrete, the exposed area of bedrock is large, the terrain is undulant and changeable, and the microrelief is very complex. Outcrop heterogeneity and diversity of crack structure of rocks under the ground form multi-layer spatial structures and many different types of small ecosystems, resulting in a large loss of organic carbon in karst soil [36, 37].

Driving Factors of SOC_D under Different Topographical Features

A large number of studies have shown that karst landforms in Guizhou can be divided into three genetic types due to the big differences in causes and combination forms of geological and topographic features: dissolution landform, dissolution-erosion landform, and dissolved structural landform [38]. Among them, the dissolution landform can be divided into four types: karst canyon, karst peak cluster depression, karst plateau, and karst trough valley. The karst area of the Guizhou Province accounts for 73.60% of the total land area of the province, and 95% of the counties (cities) in the province have karst distribution. Therefore, the study of the driving factors of SOC_D under different topographical features in the Guizhou area has an obvious promoting effect on the mechanism of soil organic carbon sequestration. The dynamics of soil carbon mainly depend on the dynamic balance between carbon input and output, and all the factors that can affect the accumulation and decomposition of soil organic carbon may affect the distribution of SOC_D [39, 40]. On the whole, the value of SOC_D is affected by many natural and human factors, such as environmental factors, soil properties, the change in land use, etc. A karst ecosystem has the characteristics of high

Table 3. Comparison of Carbon Contents in Karst and Non-Karst Soils.

Different research areas	SOC/(g·kg ⁻¹)		SOC _D /(kg·m ⁻²)	
	Content range	Mean	Content range	Mean
The Loess Plateau of China	1.28-24.55	16.98	1.39-33.41	10.92
San Jiang Plain of China	1.36-63.21	24.20	1.26-6.76	9.72
Mediterranean natural areas	24.99-49.43	37.21	0.63-10.60	-
Eastern Usambara Mountains	9.20-49.80	24.9	16.90-22.40	7.4
Guizhou Karst Region (this Study)	10.62-89.11	23.28	0.33-19.54	5.23
Guizhou non-karst region (this study)	11.31-60.11	22.87	0.33-19.54	3.98

calcium, carbonate, alkalinity, unbalanced nutrients, rapid hydrological processes, etc. The composition, structure, stability, and microbial transformation process of soil organic carbon may be different from other ecosystems. In addition, once the karst natural ecosystem is degraded or reclaimed into farmland (fertilizing) by human disturbance, soil organic carbon is easy to lose quickly [41, 42].

Comprehensive analysis shows that the distribution characteristics of organic carbon density in karst soil are the result of the interaction of environmental factors and human activities [43]. However, these environmental factors have both direct and indirect components under different topographical features. In karst canyon, and karst peak cluster depression regions, soil thickness has the greatest influence on organic carbon density, which is the direct influencing factor, and the influence degree of land use is the least, which is the indirect influencing factor [44]. So soil thickness is the key factor of SOCD in karst canyon and karst peak cluster depression regions. Because human activities in canyon and peak cluster depression regions are small and the land use is single. It has few influences on the input and output of SOCD. Land use is the largest, and the rock outcrops are the smallest in the karst plateau regions. People in the karst plateau regions tend to plan for all kinds of agricultural production land, resulting in a large amount of loss of topsoil due to frequent human disturbance. Therefore, as a result, rock outcrops and soil thickness directly determine the total amount of soil and then affect the organic carbon density in karst plateau regions. In karst trough valley regions, SOCD is most affected by rock slope aspects. The big difference in sunlight and temperature in different slope aspects results in different microclimates, vegetation types, and growth, and degradation rates of soil organic matter in the regions. The natural organic carbon density shows different characteristics. Land use in karst trough valley regions has the least influence, mainly because people in these regions tend to leave the land idle and succeed in grassland, various forest land, or degraded to wasteland through natural ecosystems. Human activities are less, and then its influence on SOCD is the smallest. Due to the complexity of SOCD and the diversity of karst environments, the main factors are different in regions of different landforms. Therefore, different ways and methods should be adopted according to the differences of the karst environment for reasonable and effective utilization of SOCD.

Conclusions

This study provided novel insights into how topography and landform features affected the content and stability of soil organic carbon. That emphasized significant variations in soil organic carbon density (SOCD) across different landscapes. The average SOCD (0-100 cm) was $5.61 \text{ kg}\cdot\text{m}^{-2}$ in non-karst regions

and $5.01 \text{ kg}\cdot\text{m}^{-2}$ in karst regions. The vertical variation characteristics of SOCD under different topography and landform characters are different. Generally, SOCD increased with soil depth. The SOCD order across different landforms was karst plateau>karst trough valley>karst peak cluster depression>karst canyon>non-karst area. The driving factors of soil organic carbon were different under different landform conditions. This study examined the factors affecting local karst soil organic carbon density (SOCD). Soil thickness primarily influenced SOCD in the karst canyon, karst peak cluster depression, and karst plateau regions. The slope aspect was the key determinant in karst trough valley regions. It reflected the global context from a local perspective. The study emphasized the unique landscapes of karst areas and highlighted the importance of maintaining and enhancing soil organic carbon for soil health, ecological balance, and erosion control. These fragile landscapes required concentrated efforts to increase soil carbon levels. The study underscored the importance of understanding organic carbon distribution for effective land-use planning. Implementing optimal planting and cultivation methods could enhance land-use efficiency, reduce dependence on fertilizers, and promote sustainable agriculture. This approach not only improved soil health but also contributed to mitigating climate change by sequestering CO_2 and reducing the greenhouse effect.

Author Contributions

Resources, conceptualisation writing-reviewing, and editing: XFH. Conceptualisation, data curation, formal analysis, methodology, visualisation, writing-original draft preparation, writing-reviewing, and editing: JH, WH. Visualisation, writing-reviewing and editing: HFW.

Funding

This work was financially supported by the Guizhou Provincial Science and Technology Projects (QKHZC [2023]-141, QKHZC [2023]-204)

Data Availability

The datasets generated during and/or analyzed during the current study are available from the corresponding author on request.

Conflict of Interest

The authors declare no financial and non-financial conflicts of interest. The funding sponsors had no role in the design of the study; in the collection, analyses,

or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

References

- ZHANG M., LIU Y., WEI Q., GU X., LIU L., GOU J. Biochar application ameliorated the nutrient content and fungal community structure in different yellow soil depths in the karst area of Southwest China. *Front Plant Science*. **13**, 1020832, **2022**.
- ZUMPARO V., PISANO L., PARISE M. An integrated framework to identify and analyze karst sinkholes. *Geomorphology*. **332**, 213, **2019**.
- JIANG Z.C., LUO W.Q., DENG Y., CAO J.H., QIN X.M., LI Y.Q. The Leakage of Water and Soil in the Karst Peak Cluster Depression and Its Prevention and Treatment. *Acta Geoscientica Sinica*. **3**, 535, **2014**.
- ROCK H., SUDARSANA M., ALMIATI R. Rock Physics Modeling and Seismic Interpretation to Estimate Shally Cemented Zone in Carbonate Reservoir Rock. *Journal of Geoscience, Engineering, Environment, and Technology*. **1**, 23, **2017**.
- GUTIÉRREZ F., PARISE M., DEWAELE J., JOURDE H. A review on natural and human-induced geohazards and impacts in karst. *Earth Science Review*. **138**, 61, **2014**.
- BARTOLOMÉ M., SANCHO C., BENITO G., MEDIALDEA A., CALLE M., MORENO A. Effects of glaciation on karst hydrology and sedimentology during the Last Glacial Cycle: The case of Granito cave, Central Pyrenees (Spain). *Catena*. **206**, 105252, **2021**.
- ALKHOURY I., BOITHIAS L., BAILEY R.T., OLLIVIER C., SIVELLE V., LABAT D. Impact of land-use change on karst spring response by integration of surface processes in karst hydrology: The ISPEEKH model. *Journal of Hydrology*. **626**, 130300, **2023**.
- YING B., XIONG K., WANG Q., WU Q. Can agricultural biomass energy provide an alternative energy source for karst rocky desertification areas in Southwestern China? investigating Guizhou Province as example. *Environmental Science and Pollution Research*. **28** (32), 44315, **2021**.
- TANG Q., XU Y., BENNETT S.J., LI Y. Assessment of soil erosion using RUSLE and GIS: a case study of the Yangou watershed in the Loess Plateau, China. *Environmental Earth Sciences*. **73** (4), 1715, **2015**.
- YE X., KUANG H. Evaluation of ecological quality in southeast Chongqing based on modified remote sensing ecological index. *Scientific Reports*. **12** (1), 15694, **2022**.
- QIU S., PENG J., DONG J., WANG X., DING Z., ZHANG H.Q. Understanding the relationships between ecosystem services and associated social-ecological drivers in a karst region: A case study of Guizhou Province, China. *Progress in Physical Geography. Earth and Environment*. **45**, 98, **2020**.
- ZHANG T., ZUO S., YU B., ZHENG K., CHEN S., HUANG L. Spatial patterns and controlling factors of the evolution process of karst depressions in Guizhou province, China. *Journal of Geographical Sciences*. **33** (10), 2052, **2023**.
- LU Q., ZHAO C., HUANG H. Comparative Study on the Temporal and Spatial Evolution of the Ecosystem Service Value of Different Karst Landform Types: A Case Study in Guizhou Province, China. *Applied Sciences*. **12**, 12801, **2022**.
- WEN H., LUO T., WANG Y., WANG S., LIU T., XIAO N. Molecular phylogeny and historical biogeography of the cave fish genus *Sinocyclocheilus* (Cypriniformes: Cyprinidae) in southwest China. *Integrative Zoology*. **17** (2), 311, **2022**.
- BARNA J.M., FRYAR A.E., CAO L., CURRENS B.J., PENG T., ZHU C. Variability in Groundwater Flow and Chemistry in the Houzhai Karst Basin, Guizhou Province, China. *Environmental & Engineering Geoscience*. **26** (3), 273, **2020**.
- ZHUO Z., CHEN Q., ZHANG X., CHEN S., GOU Y., SUN Z. Soil organic carbon storage, distribution, and influencing factors at different depths in the dryland farming regions of Northeast and North China. *Catena*. **210**, 105934, **2022**.
- BALKOVIČ J., SKALSKÝ R., FOLBERTH C., KHABAROV N., SCHMID E., MADARAS M. Impacts and Uncertainties of +2°C of Climate Change and Soil Degradation on European Crop Calorie Supply. *Earths Future*. **6** (3), 373, **2018**.
- TERRER C., PHILLIPS R.P., HUNGATE B.A., ROSENDE J., PETT-RIDGE J., CRAIG M.E. A trade-off between plant and soil carbon storage under elevated CO₂. *Nature*. **591** (7851), 599, **2021**.
- PENG X., DAI Q. Drivers of soil erosion and subsurface loss by soil leakage during karst rocky desertification in SW China. *International Soil and Water Conservation Research*. **10**, **2021**.
- ZHAO L., HOU R. Human causes of soil loss in rural karst environments: a case study of Guizhou, China. *Scientific Reports*. **9** (1), 3225, **2019**.
- ZHANG Z., ZHOU Y., HUANG X. Applicability of GIS-based spatial interpolation and simulation for estimating the soil organic carbon storage in karst regions. *Global Ecology and Conservation*. **21**, e00849, **2019**.
- WU Y., TIAN X., WANG R., ZHANG M., WANG S. Effects of vegetation restoration on distribution characteristics of heavy metals in soil in Karst plateau area of Guizhou. *PeerJ*. **11**, e15044, **2023**.
- LI Y., XIONG K., LIU Z., LI K., LUO D. Distribution and influencing factors of soil organic carbon in a typical karst catchment undergoing natural restoration. *Catena*. **212**, 106078, **2022**.
- ZHANG X.B., BAI X.Y., HE X.B. Soil creeping in the weathering crust of carbonate rocks and underground soil losses in the karst mountain areas of southwest China. *Carbonates and Evaporites*. **26** (2), 149, **2011**.
- BAI Y., ZHOU Y. The main factors controlling spatial variability of soil organic carbon in a small karst watershed, Guizhou Province, China. *Geoderma*. **357**, 113938, **2020**.
- LI C., WANG X., QIN M. Spatial variability of soil nutrients in seasonal rivers: A case study from the Guo River Basin, China. *PLOS ONE*. **16**, e0248655, **2021**.
- LUOBIN Y. Spatial variability in soil pH and land use as the main influential factor in the red beds of the Nanxiong Basin, China. *PeerJ*. **15**, 22, **2019**.
- ZHAO Z., SHEN Y.X., JIANG R., WANG Q. Rock outcrops change infiltrability and water flow behavior in a karst soil. *Vadose Zone Journal*. **19**, 344, **2020**.
- YU P., LI Y., LIU S., LIU J., DING Z., MA M. Afforestation influences soil organic carbon and its fractions associated with aggregates in a karst region of Southwest China. *Science of the Total Environment*. **814**, 152710, **2022**.
- HU P., LIU S., YE Y., ZHANG W., HE X., SU Y. Soil carbon and nitrogen accumulation following agricultural

- abandonment in a subtropical karst region. *Applied Soil Ecology*. **132**, 169-78, **2018**.
31. LAN J., WANG S., WANG J., QI X., LONG Q., HUANG M. The Shift of Soil Bacterial Community After Afforestation Influence Soil Organic Carbon and Aggregate Stability in Karst Region. *Frontiers in Microbiology*. **13**, 901126, **2022**.
 32. ZHANG Z., HUANG X., YUN C.Z. Spatial heterogeneity of soil organic carbon in a karst region under different land use patterns. *Ecosphere*. **11**, 554, **2020**.
 33. CHEN J.Q., JIA Y.N., HE Q.F., JIANG K., CHEN C., YE K. Effect of Land Use on the Stability of Soil Organic Carbon in a Karst Region. *Environmental Science*. **45** (1), 335, **2024**.
 34. ZHANG Y., XU X., LI Z., XU C., LUO W. Improvements in soil quality with vegetation succession in subtropical China karst. *Science of the Total Environment*. **775**, 145876, **2021**.
 35. WU M., LIU S., YE Y., ZHANG W., WANG K., CHEN H. Spatial heterogeneity and storage assessment method of surface soil organic carbon in high bulk-rock ratio slopes of Karst Regions. *Chinese Journal of Eco-Agriculture*. **23** (6), 676, **2015**.
 36. XIAO S., ZHANG W., YE Y., ZHAO J., WANG K. Soil aggregate mediates the impacts of land uses on organic carbon, total nitrogen, and microbial activity in a Karst ecosystem. *Scientific Reports*. **7** (1), 41402, **2017**.
 37. ZHANG W., ZHAO J., PAN F., LI D., CHEN H., WANG K. Changes in nitrogen and phosphorus limitation during secondary succession in a karst region in southwest China. *Plant Soil*. **391** (1), 77, **2015**.
 38. SCHWARZ K., GOCHT T., GRATHWOHL P. Transport of polycyclic aromatic hydrocarbons in highly vulnerable karst systems. *Environmental Pollution*. **159** (1), 133, **2011**.
 39. WANG M., XU S., ZHAO Y., SHI X. Climatic effect on soil organic carbon variability as a function of spatial scale. *Archives of Agronomy and Soil Science*. **63** (3), 375, **2017**.
 40. WEISSERT L.F., SALMOND J.A., SCHWENDENMANN L. Variability of soil organic carbon stocks and soil CO₂ efflux across urban land use and soil cover types. *Geoderma*. **271**, 80, **2016**.
 41. XIANG N., PENG G., YAN X.L., XIAO L. Impact of Different Afforestation Systems on Soil Organic Carbon Distribution Characteristics of Limestone Mountains. *Polish Journal of Environmental Studies*. **24**, 11436, **2015**.
 42. SEGONI S., MARTELLONI G., CATANI F. Different Methods to Produce Distributed Soil Thickness Maps and Their Impact on the Reliability of Shallow Landslide Modeling at Catchment Scale. In: Margottini C, Canuti P, Sassa K, editors. *Landslide Science and Practice: Volume 3: Spatial Analysis and Modelling*. Berlin, Heidelberg. Springer Berlin Heidelberg. **13**, 127, **2013**.
 43. YANG L., LUO P., WEN L., LI D. Soil organic carbon accumulation during post-agricultural succession in a karst area, southwest China. *Scientific Reports*. **6** (1), 37118, **2016**.
 44. PARRAS A.L., LOZANO G.B., BREVIK E.C., CERDÁ A. Soil organic carbon stocks assessment in Mediterranean natural areas: A comparison of entire soil profiles and soil control sections. *Journal of Environmental Management*. **155**, 219, **2015**.