

Original Research

Ameliorative Effect of Exogenously Applied Zinc on Cadmium-Stressed Sunflower (*Helianthus annuus* L.) by Modulating Growth, Photosynthetic Activity, Polyphenolic Compounds, and Yield Indices

Attiqa Rahman¹, Hafeez Ur Rehman¹, Hussam F. Najeeb Alawadi², Masood Iqbal Awan³, Athar Mahmood^{3*}, Mahmoud F. Seleiman⁴, Bilal Ahmad Khan^{5,6}, Abdul Wahid⁶, FNU Abdullah^{6,7}, Daniel O. Wasonga⁸

¹Department of Botany, University of Agriculture Faisalabad, 38040 Faisalabad, Pakistan

²College of Agriculture, Al-Qadisiyah University, Iraq

³Department of Agronomy, University of Agriculture Faisalabad, 38040 Faisalabad, Pakistan

⁴Department of Plant Production, College of Food and Agriculture Sciences, King Saud University, P.O. Box 2460, Riyadh 11451, Saudi Arabia

⁵Department of Agronomy, College of Agriculture, University of Sargodha, Sargodha, Pakistan

⁶Department of Soil and Crop Sciences, Texas A&M University, TX, USA

⁷Wheat Research Institute, Ayub Agriculture Faisalabad, Pakistan

⁸Department of Crop Sciences, University of Illinois Urbana-Champaign, Urbana, IL 61801, USA

Received: 9 May 2024

Accepted: 13 September 2024

Abstract

Sunflower (*Helianthus annuus* L.) is a type of annual plant that belongs to the Asteraceae family. However, various kinds of abiotic stresses, especially metal toxicity (Cd), substantially reduce its growth. The frequent release of mine and industrial wastewater as well as the overuse of chemical fertilizers are the main causes of this hazardous metal's rising proliferation. At the same time, even though plants have a robust defense against Cd toxicity, they are unable to combat greater levels of the metal's toxicity. Meanwhile, it has been demonstrated that applying zinc can significantly reduce the hazardous effects of cadmium. Therefore, the present experiment aimed to investigate the positive impacts of Zn application (0, 50, and 100 ppm) on the sunflower variety "FH-516" to combat Cd stress as an eco-friendly approach. Heavy metal (Cd) toxicity curtailed the root length, shoot length, leaf area, chlorophyll a/b ratio, 100-seed weight, and biological yield up to 32.03%, 27.03%, 26.1%, 31.33%, 34.46%, and 22.37%, respectively. Exogenous application of zinc significantly improved

*e-mail: athar.mahmood@uaf.edu.pk

the Cd-induced losses and increased the root length (37.9%), shoot length (48.4%), leaf area (56.2%), chlorophyll a/b ratio (65.64%), 100-seed weight (66.98%), and biological yield (74.04%). Mineral ions, e.g., Na⁺, of root and shoot (61.02% and 29.42%) increased under Cd stress, whereas K⁺ and Ca²⁺ decreased due to Cd toxicity in both root and shoot. Generally, the application of zinc improved the polyphenolic compounds and ion contents of plants under stressed conditions. At the same time, the application of Zn (100 ppm level as compared to 50) substantially increased the flavonoid contents up to 92.92% as compared to anthocyanin contents of 52.11% in Cd-stressed plants. Current findings reveal that using zinc could be an effective strategy against abiotic stresses and could be suggested for the decontamination of mild to moderately contaminated soils containing Cd.

Keywords: zinc application, cadmium stress, plant growth, polyphenolic compounds, photosynthetic potential

Introduction

The unusual shift in the natural environmental conditions because of growing population, elevated air and soil contamination, ecosystem toxicity from heavy metals, exhaustion of soil attributes, and worldwide climate variability has substantially influenced the potential of plants to respond to evolving climatic circumstances [1]. These alterations make agro-productivity systems vulnerable to fluctuating ecological conditions. Such hostile climatic events can cause harm to the physiological processes that plants undergo, known as stresses [2]. Environmental perturbations that limit plant development and agricultural productivity beneath the threshold point are known as abiotic stresses. These include extreme temperatures, metallic toxicities, water scarcity, soil salinity, and nutrient imbalance.

Cadmium is recognized as one of the most hazardous metal components in polluted agricultural lands due to its extensive contamination, considerable mobility, and extreme toxicity to human beings, livestock, and economic crops [3]. Both acute and long-term disclosure to Cd can lead to serious consequences, such as serious impairment and suppression of plant physiological activities, excessive production of nitrogen species and reactive oxygen species within cells that could also serve as harmful substances regulating an oxidative/nitrosative burst that can lead to an intensified process of senescence in plants, stunted growth, and ultimately plant decease [4]. Cd causes oxidative damage at the cellular level by serving to supply electrons to molecular oxygen, which in turn stimulates the formation of detrimental reactive oxygen species (ROS) [5]. Plants use an array of defense mechanisms to avert Cd toxicity, such as metal phytochelatin and retention, as well as the activation of ROS neutralization and combating systems involving both antioxidants (enzymatic and non-enzymatic) [6].

Sunflower (*Helianthus annuus* L.), an annual plant species, can accrue significant levels of U and Cd within its tissues [7]. One of its main contributions as an industrial product is its dry biomass. It has important agronomic traits like soil variability and resilience to intense climate variability [8]. Its sensory, nourishing,

and functional qualities make it a highly desirable protein source for the ingestion of humans [9]. The sunflower is recognized as a crucial plant species for use in phytotechnology due to the fact that it has a significant capacity for tolerance among heavy metals [10]. Previous research has shown that arsenic (As) and cadmium (Cd) drastically decrease the seed germination rate. Its capability to transport heavy metals from beneath the soil into the consuming parts of plants through the roots is quite promising [11]. Zinc (Zn) is an essential component of several kinds of enzymes, and it plays an integral part in the functioning role of enzymes to control a variety of cellular functions [12]. In addition to having more similarities with cadmium, zinc, which is a divalent ion, may compete with cadmium for binding sites in the soil and on the root surface, which may affect the translocation of cadmium in plants [13]. Zn application may reduce the Cd absorption from soil and perform a vital function in mitigating the noxious effects of Cd in plants by modulating a number of potential mechanisms, like minimizing the impact of restricted plant growth and regulating the process of Cd absorption and its relocation. However, these are only the basic effects of zinc application under cadmium stress. Furthermore, zinc application resulted in improved plant nutrient status, enhanced photosynthetic rate, activated antioxidant defense mechanisms, and lowered production of ROS, and at the same time also proved to be an effective strategy to reduce the harmful effects caused by cadmium on plants [14].

The objective of the present research was to find out how zinc application impacted the morpho-physiological and biological attributes of sunflowers under contexts of cadmium stress and contamination. The present study will assess the positive effects of zinc application on the overall botany of sunflower plants.

Experimental

The existing experiment was carried out in a pot at the University of Agriculture Faisalabad, postgraduate agriculture research station (Latitude: 31.383230648, Longitude: 72.9948305931) during the summer season

of 2021 in order to investigate the possible role of zinc in alleviating Cd stress in sunflower. A sunflower variety "FH-516" was obtained from the Ayub agriculture research station and sown in a pot (having dimensions 25 cm × 22 cm × 20 cm) containing an 8 kg clay soil mixture. A total of 27 pots were taken for an experiment that was organized in accordance with a two-factorial experimental layout and three replications under a completely randomized design. In each pot, 10 healthy seeds were sown up to 2 cm depth. One day before sowing, irrigation was done with tap water. These pots were 2-holed at the lower side (covered with muslin cloth) to prevent the accumulation of excess water. Regular watering was maintained, and one week after seeding, thinning was performed by maintaining four plants in every pot for further inspection. A very small quantity of urea was applied before the thinning of seedlings for better crop growth. Three levels of cadmium (0, 40, and 80 ppm) and three levels of zinc (0, 50, and 100 ppm) were used in this experiment, and the source of Cd and Zn was cadmium nitrate and zinc sulfate, respectively. These were taken from the laboratory of the Botany Department UAF-community college PARS in a required amount, and solutions were made in one-liter distilled water and applied exogenously to specific plants after 25 days of germination.

Morphological Parameters

After three weeks of supplying treatments, two plants were harvested from every pot. Plants were carefully uprooted and cleansed using ordinary tap water. Measurements were taken for morphological, biochemical, and yield attributes.

The fresh weight of the root and shoot samples was recorded immediately in a laboratory using a weighing scale. After that, the samples were stored in an oven at a temperature between 65 and 70°C for a period of two weeks in order to investigate the dry weight of the root and shoot samples. The root length, leaf length, and shoot length of each plant were recorded on a measuring scale, and the leaf area of each plant was measured using a leaf area meter.

Chlorophyll Contents

For the purpose of determining chlorophyll a, chlorophyll b, the a/b ratio, and carotenoids, the technique developed by Arnon was used [15]. Chlorophyll a, chlorophyll b, and carotenoids were assessed by measuring the absorbance of 0.1 grams of fresh leaf sample that was put in 80% acetone. The absorbance was measured at 663, 645, and 480 nm, respectively.

The following formulae were used to calculate the chlorophyll content quantitatively;

$$\text{Chl. } a \text{ (mg/g F. wt.)} = [12.7 \text{ (OD663)} - 2.69 \text{ (OD645)}] * V/1000 * W \quad (1)$$

$$\text{Chl. } b \text{ (mg/g F. wt.)} = [22.9 \text{ (OD645)} - 4.68 \text{ (OD663)}] * V/1000 * W \quad (2)$$

$$\text{Carotenoids (mg/g F.wt.)} = \text{Acar/Em-100} \quad (3)$$

$$\text{Acar} = \text{OD480} + 0.114 \text{ (OD 663)} - 0.638 \text{ (OD 645)}$$

$$\text{Em} = 2500$$

$$V = \text{Volume of the extract (ml)}$$

$$W = \text{Weight of fresh leaf tissue (g)}$$

Flavonoids

For determining flavonoids using the method proposed by Chang [16], 0.1 grams of fresh leaf material was taken and homogenized with 2 milliliters of a solution containing 80 percent acetone. Following this, 4 milliliters of distilled water were added to 1 milliliter of extracted material, and at 510 nanometers, the material absorbance was measured by using a spectrophotometer.

Anthocyanin

To determine the anthocyanin content of sunflower leaf material, using the Strack and Wray method [17], fresh leaf material weighing 0.1 grams was ground using 1 milliliter of acidic methanol. Afterward, the mixture was heated up to a temperature of 50 degrees Celsius for one hour, and then it was filtered. A spectrophotometer was used at a wavelength of 535 nm to determine the absorbance of the substance.

Ions (Na⁺, Ca²⁺, K⁺) Determination

For digestion of dry samples, freshly prepared nitric acid (HNO₃) and hydrochloric acid (HCl) together form a combination with a ratio of 3:1 (aqua regia) was used. For two hours, 1 gram of plant root and shoot samples were heated with 2 ml of aqua regia in a crucible at the flame, till all plant material was completely digested and black ash was obtained. After that, the ash was ground up into a high-quality powder and mixed with a 10% solution of nitric acid. After one hour, this sample solution was filtered and stored in airtight sampling bottles for the determination of Na⁺, K⁺, and Ca²⁺ of the root and shoot, according to the Vázquez method [18].

Seed Oil Contents (%)

Following the ginning, seed samples of sunflower plants were treated with acid and then dried for 24 hours at 40°C in the oven. The seed husk was

gently crushed using a nutcracker. The oil content of the embryo was then determined by Soxhlet extraction [19]. It can be inaccurate to describe the composition of sunflower seeds using percentages, particularly when the weight of the seeds varies. As a result, the embryo oil index was calculated to express composition based on weight rather than percentages. The oil content of the embryo was quantified both as a percentage of its weight (referred to as embryo oil percentage, EO%) and as a weight in milligrams (known as embryo oil index, EOI).

$$\begin{aligned} \text{Embryo oil index (weight of oil in 100 embryos)} \\ = \text{embryo oil \%} \times \text{acid-delinted, seed index}/100 \end{aligned}$$

Protein Contents

Proteins were estimated by using the Brad Ford method [20]. After grinding the sample, 0.2 grams of phosphate buffer with a pH of 7 were added to 10 milliliters of the solution. Afterward, the extract was filtered. Test tubes were filled with 1 milliliter of the plant sample and 1 milliliter of the extract. After allowing the test tubes to be kept at room temperature for half an hour, 0.5 milliliters of the Folin reagent were added. Again, they were kept at room temperature, and 5 mL of distilled water was added. Measurement of absorbance was performed at a wavelength of 620 nm. A blank was prepared by taking 1 mL of Buffer and 2 mL of Bradford.

Yield Attribute

The number of seeds of each flower was calculated through a simple counting method, and mean values were calculated, respectively. The weight of 100 completely grown seeds was measured using a weighing scale, and the average results were determined. The harvest index was calculated by computing the ratio of grain yield divided by total biomass at the time of final harvest.

The calculation was performed using the formula shown below.

$$\text{Harvest index (\%)} = \frac{\text{grain yield}}{\text{biological yield}} \times 100 \quad (4)$$

The sum of grain and straw yields was used to calculate biological yield.

Biological yield (g) = grain yield + straw yield was calculated using the formula biological

$$\text{yield (g)} = \text{grain yield} + \text{straw yield} \quad (5)$$

Determination of Leaf Cd Content

Following the washing process with deionized water, the leaf samples were subjected to drying in an oven at a temperature of 105°C for a duration of 30 minutes. Subsequently, the samples were incubated at a temperature of 75°C until a stable weight was achieved and maintained. The ground sample weighing 0.250 g was subjected to microwave digestion using a CEM MARS5 instrument (United States). The digestion process lasted for 30 minutes at a temperature of 180°C, using 8 mL of HNO₃. Afterward, the concentration of Cd in the leaves was quantified by ICP-MS (Agilent ICP-MS 7700ce, Agilent Technologies, Santa Clara, CA, United States).

Determination of Leaf Zn Content

Parts of the plant that were above ground, including the leaves, were taken from every treated plant and dried for three days in the sun and air. After that, they were dried in an oven at 60°C for seventy-two hours and then crushed to pass via a 1 mm mesh. A cylindrical cup, holding 1 gram of the sample that had been dried in the oven, was put inside a muffle burner made of carbolite (AAF1100). The cup was heated to a temperature of 300°C for a duration of one hour, resulting in the production of ash. The sample was cooled, and then the temperature was increased to 550°C for 8 to 9 hours. After the addition of 2 milliliters of concentrated hydrochloric acid and an adequate amount of water, the sample was left to evaporate in the combustion chamber for a duration of 15 to 20 minutes on a heated surface. After cooling down again, ten milliliters of 20% hydrogen nitrate were added to the sample, which was then left in the water bath with the closed furnace lid for one hour. The sample was first thoroughly sanitized and filled up to a 100 mL capacity before it was filtered by using Whatman filter paper No. 2 into a volumetric flask. Subsequently, an ordinary solution was formed. In order to determine the amount of zinc present in the combination, an ICP-Optical Emission Spectrometer (Perkin Elmer, Optima 8300, produced by PerkinElmer Corporation in Norwalk, United States of America) was used.

Statistical Data Processing and Analysis

The present experiment was designed with three replications using the CRD statistical design. Statistix 8.1 software was used to perform a two-factor analysis of variance, and Tukey's test was employed for comparing the means at the P≤0.05 level of significance. The Pearson correlation coefficient was determined using the statistical software Origin Pro 2022, and Microsoft Excel (Version Office 365) was used to construct the graphics of the obtained data.

Results

Morphological Parameters

Statistical analysis revealed that a significant effect of Zn under cadmium stress for morphological attributes, i.e., shoot fresh weight, root fresh weight, shoot dry weight, root dry weight, root length, shoot length, leaf length, and leaf area. Cadmium stress causes significant reductions in all morphological attributes of sunflowers. The morphological attributes of sunflower were considerably affected by cadmium stress at 80 ppm level as compared to 40 ppm (Fig. 1). The most significant reduction of morphological features was observed at non-zinc treatments. Shoot fresh weight, shoot dry weight, and root fresh weight showed more reduction under Cd stress (47.6%, 45.68%, and 39.14%), respectively, in comparison with control (Fig. 1. C, D, A). Whereas root length, root dry weight, shoot length, leaf length, and leaf area showed less reductions (Fig. 1. H, B, G, F, and E) under Cd stress (32.03%, 25.9%, 27.03%, 26.5%, and 26.1%), respectively, in comparison with treatments. However, the application of Zn (100 ppm) played a crucial role in mitigating Cd stress, which significantly increases the shoot fresh weight, shoot dry weight, root fresh weight, root dry weight, shoot length, leaf length, root length, and leaf area up to (77.2%, 77.3%, 70.7%, 62.8%, 48.4%, 36.2%, 37.9%, and 56.2%), respectively (Fig. 1 and Table 1). However, the overall interaction between all factors Zinc×Cd was statistically non-significant.

Photosynthetic Pigments and Phenolic Compounds

The acquired data for photosynthetic pigments showed a significant effect of Cd stress, which reduces the overall efficiency of the photosynthesis process. The higher level of Cd toxicity (80 ppm) reduced the Chl. a content (32.72%) (Fig. 2. A). Other photosynthetic pigments, e.g., Chl. b, Chl. ratio a to b, and carotenoids also showed maximum reduction under Cd stress up to 30.71%, 31.33%, and 23.15% as compared to non-stressed plants (Fig. 2. B, C, D). A significant effect of Zn application was recorded in the present study, which increased the Chl. a and carotenoids contents greatly by 85.28% and 84.88%, respectively. However, the application of Zn (100 ppm level as compared to 50) minimized the Cd-induced reduction. A significant increase of 53.82% and 65.64% was noticed for Chl. b and Chl. ratio a to b subsequently (Fig. 2). In short, there was no significant interaction seen between the stress caused by Cd and the administration of Zn. As the level of Cd toxicity increased from 40 ppm to 80 ppm, there was a reduction in photosynthetic pigments. Similarly, the higher level of Zn (100 ppm) application also enhanced the photosynthetic efficiency of sunflowers at the same rate (Table 1).

The data collected for flavonoids and anthocyanin showed that Cd stress significantly played a role in the overall reduction of plant phenolic contents. In the present study, flavonoids and anthocyanin contents were reduced by 16.37% and 18.37%, respectively, under the influence of Cd stress at an 80 ppm level (Fig. 2). However, the current investigation demonstrated considerable impacts of Cd toxicity on plants, which ultimately becomes the major constraint towards the self-defensive system of sunflowers against abiotic stresses. At the same time, application of Zn (100 ppm level as compared to 50) substantially increased the flavonoid contents up to 92.92% as compared to anthocyanin contents 52.11% in Cd-stressed plants (Fig. 2. E, F). The maximum increase in phenolic contents was recorded in non-stressed plants (Table 1). The interactive effect of Zn and Cd was noted to be non-significant.

Mineral Ions Content

The concentrations of various ions changed considerably under Cd toxicity. The Na⁺ ion concentration increased greatly in roots (61.02%) at 80 ppm Cd stress (Fig. 3. A), but the Na⁺ ion concentration in shoots was increased up to 29.42% under Cd stress (Fig. 3. B). However, the concentrations of root and shoot K⁺ and Ca²⁺ showed more reduction due to Cd stress. The root K⁺ revealed less reduction (35.72%) as compared to shoot K⁺ (43.87%) at 80 ppm Cd stress. The concentration of Ca²⁺ ions in roots showed more reduction (36.01%) as compared to shoot (26.98%) due to Cd stress at 80 ppm. Additionally, the exogenous application of Zn (100 ppm) showed significant results in terms of increased concentrations of root and shoot Na⁺, K⁺, and Ca²⁺ up to 22.25%, 91.12%, 103.92%, 45.72%, 68.07%, and 55.86%, respectively (Fig. 3. and Table 1). The combined effects of Zn and Cd were recorded as non-significant in the present study.

Yield Attributes

Cd stress and its interaction with plant yield attributes showed a significant trend at 40 and 80 ppm levels (Fig. 4). Recorded data for biological yield, harvesting index, number of seeds, 100-seed weight, oil contents, and protein contents revealed a decrease of 22.37%, 25.55%, 28.97%, 34.46%, 39.11%, and 42.84%, respectively, due to Cd stress at 80 ppm (Fig. 4. A-F). In the present study, the application of Zn as a growth promoter micronutrient for sunflowers showed maximum effects to mitigate Cd toxicity. With the application of Zn (100 ppm level), there was a substantial increase in yield parameters such as biological yield, harvesting index, number of seeds, 100-seed weight, oil contents, and protein contents up to 74.04%, 65.76%, 51.46%, 66.98%, 86.12%, and 104.37%, respectively (Fig. 4. A-F and Table 1). In conclusion, no significant interaction impact was seen between the stress caused by Cd and the administration of Zn.

Cd Contents in Leaf

The content of Cd in sunflower leaves exhibited considerable variations across several treatments of Cd-contaminated soil, e.g., 0 ppm, 40 ppm, and 80 ppm. At 0 ppm soil Cd control treatment, the Cd²⁺ concentration in leaves was maximum recorded up to 7 $\mu\text{g g}^{-1}$ due to already present some Cd concentration in soil. With an increase in the concentration of soil Cd treatment, there was a significant rise in the Cd content in the leaves (Fig. 5. A).

Zn Contents in Leaf

Application of Zn at different levels, e.g., 50 ppm and 100 ppm, increased the Zn²⁺ concentration in leaves of sunflower plants under Cd toxicity, which improves the overall growth attributes of Cd-stressed plants. The maximum recorded concentration of Zn²⁺ in leaves of sunflower at 50 ppm and 100 ppm Zn application ranged from 21 to 32 mg/kg. At Zn control application, 5 mg/kg Zn²⁺ concentration was also recorded in leaves due to already present Zn in soil (Fig. 5. B).




Fig. 1. Root fresh weight (A), Root dry weight (B), Shoot fresh weight (C), Shoot dry weight (D), Leaf area (E), Leaf length (F), Shoot length (G), Root length (H) of sunflower plant exogenously applied Zn with different levels as a growth promoter under Cd stress conditions. Significant differences among row spacing were measured by the least significant difference (LSD) at $p > 0.05$ and indicated by different letters.

Correlation and Polar Heatmap Analysis

Pearson's correlation study revealed a strong association between the morphological, photosynthetic, phenolic contents, mineral ions, and biological yield attributes of sunflowers. Various parameters, including root fresh weight, root dry weight, shoot fresh weight, shoot dry weight, shoot length, root length, leaf length, leaf area, anthocyanin, Chl. a, Chl. b, Chl. ratio a to b, and carotenoids, showed a strong positive correlation with the majority of indicators (Fig. 6). On the other hand, root Na^+ , shoot Na^+ , leaf Cd^{+2} uptake, and leaf Zn^{+2}

uptake revealed a negative correlation with most indices. Shoot K^+ , root K^+ , shoot Ca^{+2} , root Ca^{+2} , flavonoids, biological yield, harvesting index, number of seeds, 100-seed weight, oil contents, and protein contents did not show considerable correlations with most indexes (Fig. 6).

The results for all assessed characteristics of sunflowers in the nine treatments were shown in a polar heat map along with a dendrogram, facilitating basic comparisons of the effects of the treatments on the desired plant attributes. The findings revealed that nine therapies may be categorized into four main categories.




Fig. 2. Chlo. a (A), Chlo. b (B), Chlo. a/b (C), carotenoids (D), flavonoids (E), anthocyanin (F) of sunflower plant exogenously applied Zn with different levels as a growth promoter under Cd stress conditions. Significant differences among row spacing were measured by the least significant difference (LSD) at $p > 0.05$ and indicated by different letters.

Zn 50/Cd 80 ppm and Zn 0/Cd 80 ppm are clustered together; Zn 100/Cd 80 ppm, Zn 50/Cd 40 ppm, and Zn 0/Cd 40 ppm are clustered together; Zn 100/Cd 40 ppm, Zn 100/Cd 0 ppm are clustered together; and Zn 50/Cd 0 ppm and Zn 0/Cd 0 ppm are clustered together. Polar heatmap with dendrogram overall indicating high measured values for leaf area, medium measured values for number of seeds, and low measured values for all other growth, photosynthetic, phenolic contents, mineral ions uptake, and biological yield attributes of sunflower (Fig. 6).

Discussion

Plant growth is restricted, and plant production is negatively impacted by abiotic factors such as heat, salt, drought, and heavy metal stress, particularly cadmium [21]. The results of this experiment confirm our hypothesis that the use of Zn substantially ameliorates the morphological, photosynthetic, and yield attributes along with the phenolic and mineral ion contents of sunflowers under cadmium stress. Due to the negative effects of Cd strain, SFW, SDW, SL, RFW, RDW, RL,




Fig. 3. Root Na⁺ (A), Shoot Na⁺ (B), Root K⁺ (C), Shoot K⁺ (D), Root Ca²⁺ (E), Shoot Ca²⁺ (F) of sunflower plant exogenously applied Zn with different levels as a growth promoter under Cd stress conditions. Significant differences among row spacing were measured by the least significant difference (LSD) at $p > 0.05$ and indicated by different letters.

LA, and LL significantly decreased in contrast to the control treatments (Fig. 1). These drops in morphological attributes of sunflower were due to the harmful and toxic effects of Cd stress because Cd causes oxidative and ultrastructures damages, nutrient imbalance, and localization inside the subcellular space of plants [22]. In the current investigation, the application of Zn (100 ppm as compared to 50 ppm) shows a substantial function in mitigating Cd stress. The positive role of exogenous application of Zn was also previously studied by Rizwan [23]. The ameliorated proliferation of plants in terms of morphological attributes like SFW (77.2%),

SDW (77.3%), SL (48.4%), RFW (70.7%), RDW (62.8%), RL (37.9%), leaf area (56.2%), and leaf length (36.2%), respectively, were due to the application of Zn because it promotes cell division and elongation [24]. Auxin biosynthesis is improved with the usage of zinc, which is one of the insights given for the increases in plant dry and fresh weights [25]. In terms of the degradation of physiological and biochemical processes, excessive levels of cadmium stress (80 ppm as compared to 40 ppm) have a detrimental impact on the reduction of plant development, which ultimately causes damage to the defensive system of the plant. Plants can produce




Fig. 4. Biological yield (A), Harvesting index (B), Number of seeds (C), 100-seeds weight (D), Oil contents (E), Protein contents (F) of sunflower plant exogenously applied Zn with different levels as a growth promoter under Cd stress conditions. Significant differences among row spacing were measured by the least significant difference (LSD) at $p > 0.05$ and indicated by different letters.

Table 1. Mean values of morpho-physiological, biochemical, and biological yield attributes of sunflower under cadmium stress.

Source	DF	LA	LL	RDW	RFW	RL	SDW	SFW	SL
Cd	2	52982***	9.68***	0.32***	2.80***	16.82***	31.73***	100.09***	491.06***
Zn	2	124111***	9.10***	0.86***	2.98***	10.02***	29.20***	80.87***	802.50***
Cd*Zn	4	548 ^{ns}	0.04446 ^{ns}	0.00435 ^{ns}	0.06926 ^{ns}	0.1003 ^{ns}	0.1918 ^{ns}	0.361 ^{ns}	10.658 ^{ns}
Source	DF	Chl. a	Chl. b.	Chl. ratio a to b	Carotenoids	Anthocyanin	Flavonoids	-	-
Cd	2	0.51***	0.21***	0.68***	0.01***	0.02***	0.004***	-	-
Zn	2	1.64***	0.31***	1.89***	0.06***	0.10***	0.09***	-	-
Cd*Zn	4	0.01 ^{ns}	0.00083 ^{ns}	0.02 ^{ns}	0.0003 ^{ns}	0.0005 ^{ns}	0.00003 ^{ns}	-	-
Source	DF	Root Ca ⁺²	Root Na ⁺	Shoot Ca ⁺²	Shoot Na ⁺	Root K ⁺	Shoot K ⁺	-	-
Cd	2	0.70***	20.75***	10.41***	117.76***	8.26***	78.36***	-	-
Zn	2	3.11***	3.33***	20.49***	202.77***	19.64***	60.43***	-	-
Cd*Zn	4	0.02 ^{ns}	0.06 ^{ns}	0.12 ^{ns}	3.17 ^{ns}	0.20**	0.66 ^{ns}	-	-
Source	DF	BY	HI	Oil C	Pro. C	No. of seeds	100-seeds weight	-	-
Cd	2	239.37***	492.80***	672.44***	148.73***	9508.26***	7.033***	-	-
Zn	2	33.49***	102.13***	378.40***	74.99***	5880.15***	3.85***	-	-
Cd*Zn	4	0.51 ^{ns}	3.64 ^{ns}	4.26 ^{ns}	0.42 ^{ns}	13.81 ^{ns}	0.03 ^{ns}	-	-

*, **, ***, and ns represents low significant, moderately significant, highly significant, and non-significant respectively.

phenolic substances as a defensive strategy in order to mitigate the detrimental effects that metal stress puts on the surface of the cell which can lead to oxidative damages such as membrane damage and cell death [26]. In the present trial, Cd stress triggered a significant reduction in polyphenolic compounds of plants such as flavonoids and anthocyanin up to 16.37% and 18.37%. As a consequence of this, plants become more vulnerable to the harm that is brought about by abiotic stressors. However, the application of zinc played an antagonistic role against Cd stress and improved the plant polyphenolic compounds, e.g., flavonoids (92.92%) and anthocyanin (52.11%). Plants are protected from oxidative stress by the synthesis of polyphenolic chemicals, which have the ability to reduce the generation of free radicals and may function as antioxidants for metal ions [27].

The outcomes of the experiment that was being conducted revealed that yield attributes of sunflowers, such as biological yield, harvesting index, no. of seeds, 100-seed weight, oil contents, and protein contents, were decreased up to 22.37%, 25.55%, 28.97%, 34.46%, 39.11%, and 42.84%, respectively, due to the application of 80 ppm Cd stress. Marques [28] also illustrated that cadmium contamination has greatly reduced the accumulation of biomass in sunflower plants. Exposure to cadmium leads to the suppression of plant development due to its harmful effects, which also affect the regulatory functions of important metabolic enzymes and stimulate leaf necrosis [29]. As a result, the yield of overall plants is reduced considerably. However, the

application of zinc in plants has been proven effective in enhanced accumulation of their dry matter and protein contents as well as their total agricultural yield since it is essential for the formation of proteins and the breakdown of carbohydrates [30]. In our research trial, results revealed that exogenous applied Zn (100 ppm) significantly enhanced the yield attributes, e.g., biological yield (74.04%), harvesting index (65.76%), no. of seeds (51.46%), 100-seed weight (66.98%), oil contents (86.12%), and protein contents (104.37%) of sunflowers. The chlorophyll concentration, e.g., Chl. a (32.72%), Chl. b (30.71%), Chl. ratio a to b (31.33%), and carotenoids (23.15%), significantly reduced due to the toxic effects of Cd in contrast to control treatments. A decrease in the content of photosynthetic pigments has been observed as a frequent indication of toxicity to metals in several plant species [31]. Haseeb and Maqbool [32] studied that reduction in light-absorbing components of photosynthesis machinery (Chlorophyll a, b, ratio a to b, and carotenoids) was observed in sunflower plants due to noxious effects of abiotic stresses. According to studies by Vijayaragavan [33] and Touiserkani and Haddad [34], cadmium decreased the concentration of chlorophyll by either causing the Mg ion to be removed from its binding site in chlorophyll (which led to the deterioration of the chlorophyll molecule) or by hindering the activity of enzymes involved in the production of chlorophyll, such as protochlorophyllide reductase, porphobilinogen deaminase, and aminolevulinic dehydratase, which led to a shortage of Fe²⁺ and Mg²⁺, which in turn caused carbonic anhydrase activity to be inhibited due




Fig. 5. Cd²⁺ uptake in leaf (A), Zn²⁺ uptake in leaf (B) of sunflower plant exogenously applied Zn with different levels as a growth promoter and Cd²⁺ as an abiotic stress also applied with different levels. Zn²⁺ and Cd²⁺ uptake increases as their level of application increases. Significant differences among row spacing were measured by the least significant difference (LSD) at $p > 0.05$ and indicated by different letters.




Fig. 6. Pearson's correlation matrixes constructed (On Left Side) for morphological, photosynthetic, phenolic contents, mineral ions uptake, and biological yield attributes of sunflower under Cd stress. *Significant at $p \leq 0.05\%$; **Significant at $p \leq 0.01\%$; ***Significant at $p \leq 0.001\%$. Polar heatmap analysis between various attributes of sunflower (On Right Side) showed the impact of Zn application under Cd stress on morphological, photosynthetic, phenolic contents, mineral ions uptake, and biological yield attributes of sunflower; Zn:0/ Cd:0, Zn:50 ppm/ Cd:0, Zn:100 ppm/ Cd:0; Zn:0/ Cd:40 ppm, Zn:50/ Cd:40 ppm, Zn:100/ Cd:40 ppm; Zn:0/ Cd:80 ppm, Zn:50/ Cd:80 ppm, Zn:100/ Cd:80 ppm.

to Zn²⁺ limited supply [35]. However, in the present trial application of zinc enhanced the photosynthetic capacity of sunflowers up to 85.28%, 53.82%, 65.64%, and 84.88% were noticed for Chl. a, Chl. b, Chl. ratio a to b, and carotenoids subsequently under Cd stress. Wu [36] demonstrated that the use of Zn's impact on photosynthesis's enzyme activity might be the cause of improving photosynthetic attributes, as foliar treatments may enhance the associated zinc enzyme [37]. Previous studies have shown that the combination of amino acids and microelements enhances the rate of

photosynthesis in plants when applied independently or in response to external environmental stressors [38, 39]. Research has also shown that the reduced toxicity of Cd in wheat plants or the enhanced presence of zinc as a plant microelement might be a significant factor contributing to the increase in photosynthetic indices. The concentration of Na⁺ increased in both shoot and root while the concentrations of Ca²⁺ and K⁺ decreased due to Cd stress. Kosakivska [40] explained that Cd inhibits nitrate reductase, reducing the transport and intake of nitrate and interfering with the absorption

of Ca, P, K, Mg, and water. Metal transporters enable the transportation of inorganic minerals to and from vascular bundles during the loading and unloading process. Increased levels of Cd will lead to detrimental effects due to the interactions between Cd ions and other elements. Hence, the translocation of Cd ions in plant cells would be diminished if they were conveyed by cation channels or other transporters [41]. Nevertheless, in the current study, the use of zinc substantially increased the levels of plant root and shoot Na^+ , K^+ , and Ca^{2+} up to 22.25%, 91.12%, 103.92%, 45.72%, 68.07%, and 55.86%, respectively [42]. The decrease in the accumulation of mineral ions has an adverse effect on the processes of metabolism and photosynthetic indices [43]. A key approach to detoxifying heavy metal-contaminated soils is called phytoextraction, which functions via the use of accumulators to draw toxins from the soil and transfer them into plants. A phytoextraction plant species with excellent metal resistance and biological accumulation index should be ideal for the reclamation of heavy metal-affected soils for sustainable agriculture [44]. In our current research trial, there was considerable Cd^{2+} accretion in the leaves of sunflower plants, and Cd concentration in the leaves showed a substantial elevation with the increasing levels of Cd soil treatments, e.g., 40 ppm and 80 ppm (Fig. 5. A). At higher levels of Cd soil treatment, various attributes (photosynthetic, growth, and yield attributes) of sunflowers were affected considerably in terms of lower photosynthetic efficiency and reduced biological yield due to Cd toxicity. Lux [45] studied that the majority of plants are hazardous to leaf Cd contents over 5-10 $\mu\text{g g}^{-1}$. In the present study, the application of Zn significantly reduced the Cd-induced losses in sunflower plants. Zinc (Zn) is a crucial micronutrient that serves as a constituent of several enzymes responsible for regulating multiple metabolic events in plants. It is also necessary for the creation of auxins and proteins. Additionally, it facilitates enhanced assimilation of nitrogen (N) and phosphorus (P) by plants [46]. As the concentration of Zn application increases from 0 ppm (control) to 100 ppm, the Zn^{2+} concentration in the leaves of sunflower also increases, ranging from 5 to 32 mg/kg. This indicates the considerable effect of Zn application to improve the growth, photosynthetic, mineral ion uptake, and yield attributes of sunflower plants under Cd-stressed conditions.

Conclusions

The present study indicates the alterations in sunflower plants exposed to Cd stress in combination with Zn usage. The usage of Zn in general agricultural practices overall ameliorates the harmful effects of Cd toxicity that pose harmful impacts on morphological, photosynthetic, phenolic contents, mineral ions, and biological yield attributes of plants. It was also studied that Cd toxicity causes oxidative and ultrastructure

damage, nutrient imbalance, and subcellular localization in plants that ultimately impair plant development and adversely affect plant productivity. Translocation and uptake of mineral ions by plants also become disturbed due to Cd stress. However, application of Zn distinctly revealed its beneficial contribution to recouping losses carried on by stress. Furthermore, experiments needed to be accomplished alongside other micronutrients to investigate hazardous metal phytoremediation procedures in field trials on a larger scale.

Funding

This research was funded by Researchers Supporting Project (RSPD2025R751), King Saud University, Riyadh, Saudi Arabia.

Acknowledgment

This research was funded by Researchers Supporting Project (RSPD2025R751), King Saud University, Riyadh, Saudi Arabia.

Data Availability

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

Conflict of Interest

The authors declare no conflict of interest.

References

- AKRAM M., ULLAH S., IKRAM K., ZAHEER M.S., MANSHA M.Z., ALI H.H., NIAZ Y., SELEIMAN M.F., KHAN N. Effect of cadmium toxicity on Zn, Mn, Cd, Fe concentration and translocation, bioaccumulation, uptake of Cd in root and shoot of different rice varieties. *Pakistan Journal of Botany*, **56** (1), 55, **2024**.
- RAZA A., ASHRAF F., ZOU X., ZHANG X., TOSIF H. Plant adaptation and tolerance to environmental stresses: mechanisms and perspectives. *Plant Ecophysiology and Adaptation under Climate Change: Mechanisms and Perspectives I: General Consequences and Plant Responses*, **117**, **2020**.
- KHAN I., SELEIMAN M.F., CHATTHA M.U., JALAL R.S., MAHMOOD F., HASSAN F.A.S., IZZET W., ALHAMMAD B.A., ALI E.F., ROY R., USLU O.S., HASSAN M.U. Enhancing antioxidant defense system of mung bean with a salicylic acid exogenous application to mitigate cadmium toxicity. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*, **49** (2), 12303, **2021**.
- SELEIMAN M.F., AL-SUHAIBANI N., EL-HENDAWY S., ABDELLA K., ALOTAIBI M., ALDERFASI A.

- Impacts of long- and short-term of irrigation with treated wastewater and synthetic fertilizers on the growth, biomass, heavy metal content, and energy traits of three potential bioenergy crops in arid regions. *Energies*, **14** (11), 3037, **2021**.
5. FARAZ A., FAIZAN M., SAMI F., SIDDIQUI H., HAYAT S. Supplementation of salicylic acid and citric acid for alleviation of cadmium toxicity to Brassica juncea. *Journal of plant growth regulation*, **39**, 641, **2020**.
 6. SAIDI I., GUESMI F., KHARBECH O., HFAIEDH N., DJEBALI W. Gallic acid improves the antioxidant ability against cadmium toxicity: Impact on leaf lipid composition of sunflower (*Helianthus annuus*) seedlings. *Ecotoxicology and Environmental Safety*, **210**, 111906, **2021**.
 7. ALSABBAGH A.H., ABUQUDAIRA T.M. Phytoremediation of Jordanian uranium-rich soil using sunflower. *Water, Air, & Soil Pollution*, **228**, 1, **2017**.
 8. RIZWAN M., ALI S., ADREES M., RIZVI H., ZIAUR-REHMAN M., HANNAN F., QAYYUM M.F., HAFEZ F., OK Y.S. Cadmium stress in rice: toxic effects, tolerance mechanisms, and management: a critical review. *Environmental Science and Pollution Research*, **23**, 17859, **2016**.
 9. ZORZI C.Z., GARSKE R.P., FLÔRES S.H., THYS R.C.S. Sunflower protein concentrate: A possible and beneficial ingredient for gluten-free bread. *Innovative Food Science & Emerging Technologies*, **66**, 102539, **2020**.
 10. YUCE M., TASPINAR M.S., AYDIN M., AGAR G. Response of NAC transcription factor genes against chromium stress in sunflower (*Helianthus annuus* L.). *Plant Cell, Tissue and organ culture*, **136**, 479, **2019**.
 11. BASHIR S., QAYYUM M.A., HUSAIN A., BAKHSH A., AHMED N., HUSSAIN M.B., ELSHIKH M.S., ALWAHIBI M.S., ALMUNQEDHI B.M., HUSSAIN R. Efficiency of different types of biochars to mitigate Cd stress and growth of sunflower (*Helianthus*; L.) in wastewater irrigated agricultural soil. *Saudi Journal of Biological Sciences*, **28** (4), 2453, **2021**.
 12. SELEIMAN M.F., AHMAD A., BATTAGLIA M.L., BILAL H.M., ALHAMMAD B.A., KHAN N. Zinc oxide nanoparticles: A unique saline stress mitigator with the potential to increase future crop production. *South African Journal of Botany*, **159**, 208, **2023**.
 13. ZARE A., KHOSHGOFTARMANESH A., MALAKOUTI M., BAHRAMI H., CHANEY R. Root uptake and shoot accumulation of cadmium by lettuce at various Cd: Zn ratios in nutrient solution. *Ecotoxicology and Environmental Safety*, **148**, 441, **2018**.
 14. RIZWAN M., ALI S., REHMAN M.Z.U., MAQBOOL A. A critical review on the effects of zinc at toxic levels of cadmium in plants. *Environmental Science and Pollution Research*, **26**, 6279, **2019**.
 15. ARNON D.I. Copper enzymes in isolated chloroplasts. Polyphenoloxidase in Beta vulgaris. *Plant physiology*, **24** (1), 1, **1949**.
 16. CHANG C.-C., YANG M.-H., WEN H.-M., CHERN J.-C. Estimation of total flavonoid content in propolis by two complementary colorimetric methods. *Journal of food and drug analysis*, **10** (3), **2002**.
 17. STRACK D., WRAY V. Anthocyanins. *Methods in Plant Biochemistry*. Elsevier, **1**, 326, **1989**.
 18. VÁZQUEZ M., MIKHelson K., PIEPPONEN S., RÄMÖ J., SILLANPÄÄ M., IVASKA A., LEWENSTAM A., BOBACKA J. Determination of Na⁺, K⁺, Ca²⁺, and Cl⁻ Ions in Wood Pulp Suspension Using Ion-Selective Electrodes. *Electroanalysis: An International Journal Devoted to Fundamental and Practical Aspects of Electroanalysis*, **13** (13), 1119, **2001**.
 19. DE CASTRO M.L., GARCIA-AYUSO L. Soxhlet extraction of solid materials: an outdated technique with a promising innovative future. *Analytica chimica acta*, **369** (1-2), 1, **1998**.
 20. BRADFORD M.M. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical biochemistry*, **72** (1-2), 248, **1976**.
 21. IMRAN Q.M., FALAK N., HUSSAIN A., MUN B.-G., YUN B.-W. Abiotic stress in plants; stress perception to molecular response and role of biotechnological tools in stress resistance. *Agronomy*, **11** (8), 1579, **2021**.
 22. LI L., LI J., HE Q., ZHU S., CHEN J. Nutrients, ultrastructures, and Cd subcellular localization in the cottonseeds of three upland cotton cultivars under Cd stress. *Journal of soil science and plant nutrition*, **14** (2), 278, **2014**.
 23. RIZWAN M., ALI S., ALI B., ADREES M., ARSHAD M., HUSSAIN A., UR REHMAN M.Z., WARIS A.A. Zinc and iron oxide nanoparticles improved the plant growth and reduced the oxidative stress and cadmium concentration in wheat. *Chemosphere*, **214**, 269, **2019**.
 24. FARD S.R., KHOURGAMI A., RAFEE M., NASROLLAHI H. Study the effect of zinc spraying and plant density on seed yield and morphological characteristics of Green gram. *Annals of Biological Research*, **3** (8), 4166, **2012**.
 25. KOBRAEE S. Effect of foliar fertilization with zinc and manganese sulfate on yield, dry matter accumulation, and zinc and manganese contents in leaf and seed of chickpea (*Cicer arietinum*). *Journal of Applied Biology and Biotechnology*, **7** (3), 20, **2019**.
 26. ZOUARI M., AHMED C.B., ZORRIG W., ELLOUMI N., RABHI M., DELMAIL D., ROUINA B.B., LABROUSSE P., ABDALLAH F.B. Exogenous proline mediates alleviation of cadmium stress by promoting photosynthetic activity, water status and antioxidative enzymes activities of young date palm (*Phoenix dactylifera* L.). *Ecotoxicology and environmental safety*, **128**, 100, **2016**.
 27. MOUSAVI A., POURAKBAR L., MOGHADDAM S.S., POPOVIĆ-DJORDJEVIĆ J. The effect of the exogenous application of EDTA and maleic acid on tolerance, phenolic compounds, and cadmium phytoremediation by okra (*Abelmoschus esculentus* L.) exposed to Cd stress. *Journal of Environmental Chemical Engineering*, **9** (4), 105456, **2021**.
 28. MARQUES A.P., MOREIRA H., FRANCO A.R., RANGEL A.O., CASTRO P.M. Inoculating *Helianthus annuus* (sunflower) grown in zinc and cadmium contaminated soils with plant growth promoting bacteria—Effects on phytoremediation strategies. *Chemosphere*, **92** (1), 74, **2013**.
 29. AMAYA-CARPIO L., DAVIES F., FOX T., HE C. Arbuscular mycorrhizal fungi and organic fertilizer influence photosynthesis, root phosphatase activity, nutrition, and growth of *Ipomoea carnea* ssp. *fistulosa*. *Photosynthetica*, **47**, 1, **2009**.
 30. DROSTKAR E., TALEBI R., KANOUNI H. Foliar application of Fe, Zn and NPK nano-fertilizers on seed yield and morphological traits in chickpea under rainfed condition. *Journal of Resources and Ecology*, **4** (1), 221, **2016**.
 31. GIANNAKOULA A., THERIOS I., CHATZISSAVVIDIS C. Effect of lead and copper on photosynthetic apparatus

- in citrus (*Citrus aurantium* L.) plants. The role of antioxidants in oxidative damage as a response to heavy metal stress. *Plants*, **10** (1), 155, **2021**.
32. HASEEB M., MAQBOOL N. Influence of foliar applied nitrogen on reproductive growth of sunflower (*Helianthus annuus* L.) under water stress. *Agricultural Sciences*, **6** (12), 1413, **2015**.
 33. VIJAYARAGAVAN M., PRABHAHAR C., SURESHKUMAR J., NATARAJAN A., VIJAYARENGAN P., SHARAVANAN S. Toxic effect of cadmium on seed germination, growth and biochemical contents of cowpea (*Vigna unguiculata* L.) plants. *International multidisciplinary research journal*, **1** (5), **2011**.
 34. TOUISERKANI T., HADDAD R. Cadmium-induced stress and antioxidative responses in different Brassica napus cultivars. *Journal of Agricultural Science and Technology*, **14** (4), 929, **2012**.
 35. SADIQ R., MAQBOOL N., HASEEB M. Ameliorative effect of chelating agents on photosynthetic attributes of Cd stressed sunflower. *Agricultural Sciences*, **8** (02), 149, **2017**.
 36. WU C., DUN Y., ZHANG Z., LI M., WU G. Foliar application of selenium and zinc to alleviate wheat (*Triticum aestivum* L.) cadmium toxicity and uptake from cadmium-contaminated soil. *Ecotoxicology and Environmental Safety*, **190**, 110091, **2020**.
 37. HAGHIGHI M., RAMEZANI M.R., RAJAI N. Improving oxidative damage, photosynthesis traits, growth and flower dropping of pepper under high temperature stress by selenium. *Molecular biology reports*, **46**, 497, **2019**.
 38. LEE C.W., KANG H. UV Photolysis of Glycine on Ice Films: Implication for Photosynthesis and Photodestruction of Amino Acids in Interstellar Medium#. *Bulletin of the Korean Chemical Society*, **36** (3), 784, **2015**.
 39. MOUSAA I., RADH H. Photosynthesis of anticorrosive protective coatings for steel substrate based on acrylated oil containing unsaturated amino acid compounds. *Progress in Organic Coatings*, **107**, 18, **2017**.
 40. KOSAKIVSKA I.V., BABENKO L.M., ROMANENKO K.O., KOROTKA I.Y., POTTERS G. Molecular mechanisms of plant adaptive responses to heavy metals stress. *Cell Biology International*, **45** (2), 258, **2021**.
 41. LIU S.-L., YANG R.-J., MA M.-D., DAN F., ZHAO Y., JIANG P., WANG M.-H. Effects of exogenous NO on the growth, mineral nutrient content, antioxidant system, and ATPase activities of *Trifolium repens* L. plants under cadmium stress. *Acta Physiologiae Plantarum*, **37**, 1, **2015**.
 42. MARIUM A., KAUSAR A., ASHRAF M.Y., KANWAL H., NAZLI Z.-I.-H. Enhancement in biomass and nutrients uptake in cucumber through methionine and phenylalanine under saline soils. *Pakistan Journal of Agricultural Sciences*, **58** (1), **2021**.
 43. SADAK M.S., ABDALLA A.M., ABD ELHAMID E.M., EZZO M. Role of melatonin in improving growth, yield quantity and quality of *Moringa oleifera* L. plant under drought stress. *Bulletin of the National Research Centre*, **44** (1), 1, **2020**.
 44. ZHAO F., LOMBI E., MCGRATH S. Assessing the potential for zinc and cadmium phytoremediation with the hyperaccumulator *Thlaspi caerulescens*. *Plant and soil*, **249**, 37, **2003**.
 45. LUX A., MARTINKA M., VACULÍK M., WHITE P.J. Root responses to cadmium in the rhizosphere: a review. *Journal of experimental botany*, **62** (1), 21, **2011**.
 46. NITIN TRIVEDI N.T., DEVI SINGH D.S., VIJAY BAHADUR V.B., PRASAD V., COLLIS J. Effect of foliar application of zinc and boron on yield and fruit quality of guava (*Psidium guajava* L.). *Agricultural and Food Sciences. HortFlora Research Spectrum*, **1** (3), 281, **2012**.