

Original Research

# Individual and Combined Effects of Sublethal Cadmium and Lead on Tissue Accumulation, Hemato-Biochemical Parameters and Neurotoxicity in *Channa marulius*

Javed Ahmed Ujan<sup>1,2</sup>, Saima Majeed<sup>3</sup>, Sawera Maqaddas<sup>4</sup>, Wajeeha Afzal<sup>5</sup>,  
Syed Sikandar Habib<sup>6\*</sup>, Alia Mushtaq<sup>7</sup>, Saira Naz<sup>8</sup>, Muhammad Zahid<sup>9</sup>, Khalid Khan<sup>9</sup>,  
Muhammad Ismail<sup>9</sup>, Mohamed Mohany<sup>10</sup>

<sup>1</sup>Department of Zoology, Shah Abdul Latif University, Khairpur, 66020, Khairpur, Sindh, Pakistan

<sup>2</sup>Department of Animal Sciences, University of Florida, Gainesville, FL, 32608, USA

<sup>3</sup>Department of Maritime Sciences, Bahria University Karachi Campus, 75000, Karachi Sindh, Pakistan

<sup>4</sup>Department of Zoology, Government College University Faisalabad, 38000, Faisalabad Punjab, Pakistan

<sup>5</sup>Department of Zoology, University of the Punjab Lahore 05422, Pakistan

<sup>6</sup>Department of Zoology, University of Sargodha, 40100, Punjab, Pakistan

<sup>7</sup>Plant and Environmental Protection, National Agricultural Research Centre, Islamabad, 45500, Pakistan

<sup>8</sup>Centre for Research on Fish Nutrition and Environmental Ecology of the Ministry of Agriculture, Shanghai Ocean University, Shanghai 201306, China

<sup>9</sup>Department of Zoology, Islamia College University, Peshawar 25120, Khyber Pakhtunkhwa, Pakistan

<sup>10</sup>Department of Pharmacology and Toxicology, College of Pharmacy, King Saud University, P.O. Box 55760, Riyadh 11451, Saudi Arabia

Received: 27 June 2024

Accepted: 7 October 2024

## Abstract

This study evaluated the physiological effects of sublethal concentrations of Cd<sup>2+</sup> and Pb<sup>2+</sup>, both individually and in combination, on *Channa marulius*, an important species in local fisheries. A total of 160 fish were divided into four groups (control, Cd<sup>2+</sup>-treated, Pb<sup>2+</sup>-treated, and Cd<sup>2+</sup> + Pb<sup>2+</sup> treated) and exposed to 30% of the 96-hour LC<sub>50</sub> values for CdCl<sub>2</sub> (22.71 mg/L), PbCl<sub>2</sub> (16.02 mg/L), and their combination for 40 days. Results showed significantly ( $P < 0.05$ ) higher Cd<sup>2+</sup> and Pb<sup>2+</sup> concentrations in the tissues of exposed groups compared to controls, with the highest accumulation in the kidney, followed by gills and intestines. Enzyme activities of catalase (CAT) and superoxide dismutase (SOD) were significantly reduced, especially in the combined exposure group. Additionally, red blood cells,

hemoglobin, and total protein levels decreased, while cortisol and glucose levels increased, indicating stress. Acetylcholinesterase (AChE) activity in the gill and liver was also significantly inhibited in the combined exposure group. These findings highlight the risks posed by combined heavy metal exposure to fish physiology, underscoring potential threats to fish health and local fisheries.

**Keywords:** oxidative stress, metal mixtures, hematology, enzymatic activities, neurotoxicity

## Introduction

Exposure to heavy metals in aquatic environments is a significant environmental concern due to the potential accumulation of these metals in humans through the consumption of aquatic products like fish, posing health risks [1]. Cadmium ( $\text{Cd}^{2+}$ ) and lead ( $\text{Pb}^{2+}$ ) are particularly toxic, even at low exposure levels, especially for aquatic organisms [2]. Fish, being at higher trophic levels, can accumulate substantial amounts of these metals, and their accumulation patterns depend on uptake and elimination rates [3]. Metal exposure in aquatic environments leads to the accumulation of harmful substances in specific tissues of aquatic animals, necessitating research on bioaccumulation in fish tissues in contrast to biotransformation and excretion [4]. The accumulation in fish tissues is influenced by factors such as the type of metal, exposure concentration and duration, water temperature, salinity, hardness, fish species, age, and metabolic activity [5, 6]. Monitoring metal accumulation patterns in fish tissues is a reliable indicator for assessing contamination in aquatic environments [7]. While some metals are essential for fish physiology, excessive accumulation can become toxic, impacting physiological functions and potentially leading to high mortality and loss of indigenous fish species [8].

Changes in blood parameters can provide insights into the overall health and well-being of aquatic organisms [9]. By analyzing these parameters, researchers can diagnose structural and functional alterations in fish exposed to toxicants, contributing to a holistic understanding of the impact of  $\text{Cd}^{2+}$  and  $\text{Pb}^{2+}$  on fish physiology [10]. Environmental contaminants are often encountered as mixtures, complicating assessments based on individual chemical toxicity [11]. Water quality criteria typically consider individual chemicals in acute and chronic bioassays, but in reality, metals coexist in ambient waters and interact, affecting uptake, bioaccumulation, and toxicity [12].

$\text{Cd}^{2+}$  and  $\text{Pb}^{2+}$  exposure leads to metal buildup in animal brains through the blood-brain barrier, impacting the central nervous system and leading to neurological diseases due to cellular dysfunction and cerebral edema [13]. Acetylcholine is a neurotransmitter essential for movement control, memory, and cerebral blood flow in both central and peripheral nervous systems [14]. It is broken down by the enzyme acetylcholinesterase (AChE), a serine protease [15]. Measuring AChE activity serves as a key biomarker for neurotoxicity induced by metals exposure [16], as  $\text{Cd}^{2+}$  and  $\text{Pb}^{2+}$  toxicity causes

neurobehavioral disorders that directly impact AChE activity [17, 18].

The *Channa marulius* has been selected for the current study due to its importance in local fisheries, often targeted for its economic value and consumed as a food source in many regions [19]. Despite its ecological and economic significance, *C. marulius*, like many freshwater species, faces threats from habitat degradation, overfishing, and pollution [20]. Consuming fish contaminated with heavy metals poses significant health risks.  $\text{Cd}^{2+}$  can lead to kidney damage and bone weakening, while  $\text{Pb}^{2+}$  exposure may cause neurological issues, particularly in children, affecting their cognitive development [4, 5]. By examining the individual effects of metals, researchers can discern the unique contribution of each metal to tissue accumulation, hemato-biochemical alterations, and neurotoxicity in fish. Furthermore, investigating the combined effects of  $\text{Cd}^{2+}$  and  $\text{Pb}^{2+}$  is essential due to the frequent occurrence of environmental contaminants as mixtures. Therefore, the current study evaluates and compares the individual and combined sublethal effects of  $\text{Cd}^{2+}$  and  $\text{Pb}^{2+}$  on *C. marulius* tissue accumulation, hemato-biochemical parameters, and neurotoxicity.

## Materials and Methods

### Test Chemicals and Fish Sampling

Cadmium, represented as  $\text{CdCl}_2 \cdot \text{H}_2\text{O}$  in its monohydrate form and a purity of 98%, and lead, in the form of  $\text{PbCl}_2$  and a purity of 98%, obtained from Merck, Germany. A total of 160 mature catfish (*C. marulius*) were obtained from Chashma Lake in Mianwali, Punjab, Pakistan. These fish exhibited a mean body length of  $26 \pm 2.71$  cm and a mean body weight of  $144 \pm 22$  g. The selected fish, considered to be in good health, were promptly transported to the laboratory in a 500 L tank equipped with aeration systems. The handling procedures adhered to the guidelines outlined in the animal welfare protocol. Upon arrival at the laboratory, the catfish were given a 14-day acclimatization period in a fiberglass tank of dechlorinated tap water, which was refreshed daily. Throughout this acclimatization phase, the fish were provided with optimal conditions to adjust to their new environment. The daily diet (morning and evening) for the catfish (2% of their body) consisted of a commercial basal fish feed sourced from Aqua Feed Company in Pakistan with a crude protein level of 35%.

## Experimental Design

After the acclimatization period, the collected fish were distributed in triplicate into four groups (10 fish/tank), each treatment with 40 fish. The capacity of each tank was 90 L. Group 1 served as the control and was maintained in dechlorinated water. Meanwhile, Groups 2, 3, and 4 were exposed to 30% of the 96-hour  $LC_{50}$  values for  $CdCl_2$  (22.71 mg/L),  $PbCl_2$  (16.02 mg/L), and a combination of  $CdCl_2 + PbCl_2$  (22.71 + 16.02), respectively. This exposure occurred under constant conditions and a 12-hour light/dark cycle, spanning a duration of 40 days. Previous research indicated mean average concentrations of  $Cd^{2+}$  and  $Pb^{2+}$  in Chashma Lake water as 0.32 and 0.73 mg/L, respectively [21]. For this study, a 30%  $LC_{50}$  concentration was considered a critical threshold, potentially leading to adverse effects in fish exposed to  $Cd^{2+}$  and  $Pb^{2+}$ . The 96-hour  $LC_{50}$  values for  $Cd^{2+}$  and  $Pb^{2+}$  in *C. marulius* were reported as 75.70 and 53.42 mg/L, respectively, by Batool et al. [22]. Throughout the study, water quality parameters were diligently monitored using the APHA [23] method. The environmental conditions remained stable for all groups, with a temperature of  $26.05 \pm 0.04^\circ C$ , dissolved oxygen (DO) at  $5.97 \pm 0.17$  mg/L, total hardness of  $138.9 \pm 3.07$ , pH of  $7.13 \pm 0.04$ , total ammonia at  $0.22 \pm 0.05$ , nitrite at  $0.08 \pm 0.001$ , and nitrate levels at  $1.83 \pm 0.03$  mg/L. To ensure consistent aeration, both the treated and control test media were aerated using a capillary system integrated into the air pump.

## Fish Tissue Sampling and Assessment of Biochemical and Metal Contents

After the experimental period, sacrificial procedures were performed on five fish from each group. The gills, liver, kidney, and intestinal tissues were carefully dissected using sterile equipment as outlined by Habib et al. [21]. Subsequently, the liver tissue underwent a wash with phosphate-buffered saline (PBS) at a pH of 7.4 to remove any erythrocytes or clots. To generate a 10% homogenate from the tissue, a precise amount was extracted, finely minced, and then homogenized using ice-cold 0.1 M phosphate buffer with a pH of 7.4. After homogenization, the samples underwent centrifugation ( $1,500 \times g$  at  $4^\circ C$ ). The resulting supernatants were collected and stored at  $-80^\circ C$  for subsequent determination of catalase (CAT) and superoxide dismutase (SOD) levels. Commercially available kits from Biodiagnostics Co. were used for these analyses. Concentrations of  $Cd^{2+}$  in the sampled tissues, including gills, intestine, and kidney, were assessed using a graphite furnace atomic absorption spectrophotometer (AAS, PerkinElmer 3300, USA).

## Analysis of Hematological and Biochemical Parameters

Blood samples were obtained from both the control and treated fish groups at the end of the experimental period. The sampling involved randomly selecting five fish specimens from each tank after a fasting period of at least 24 hours. To ensure the fish were unconscious and stress was minimized during handling, a buffered aqueous solution of MS-222 at a concentration of 40 mg/l was employed. Blood was drawn from the caudal vein using a sterile syringe shortly after rendering the fish unconscious. The collected blood samples were transferred into tubes containing the anticoagulant agent EDTA for subsequent analysis. Various blood parameters, such as the red blood cell (RBC) count, were assessed using a Neubauer counting chamber following the method outlined by Habib et al. [24]. The concentration of hemoglobin (Hb) was determined using the cyanomethemoglobin method [25], and the hematocrit (Hct) value was determined according to the procedure outlined by Amouri et al. [26].

Plasma was isolated for further analysis of biochemical parameters by centrifuging the blood at 3000 rpm for 12 minutes. The isolated plasma was stored in a deep freezer. Blood plasma glucose levels were analyzed using an assay kit (Human Diagnostics Worldwide) following the procedure described by Trinder [27]. Cortisol concentration was assessed using a commercially available enzyme-linked immunoassay (ELISA) kit obtained from Alpha Diagnostic International (San Antonio, TX, USA). The total protein content was evaluated calorimetrically, following the method outlined by Henry [28]. Additionally, alanine aminotransferase (ALT) and aspartate aminotransferase (AST) levels were determined using a calorimetric technique, following the procedure outlined by Reitman and Frankel [29].

## Neurotransmitter

Gill and liver tissues were collected and used to measure AChE activity. Each type of tissue was homogenized ten times in a 0.1 M PBS solution. The homogenized samples were then centrifuged at  $10,000 \times g$  for 30 minutes at  $4^\circ C$ . The resulting supernatant was collected for analysis. AChE activity was assessed using the method described by Pretto et al. [30], with the activity reported in terms of nanomoles (nmol) per minute per unit of tissue protein.

## Statistical Analysis

Statistical analysis of the data involved employing one-way ANOVA. The results are expressed as mean  $\pm$  standard error. The Duncan multiple-range test was utilized to assess the differences among the treatment means. Significance was considered at  $P < 0.05$ .




Fig. 1. Concentration of heavy metals in tissues of *Channa marulius* after exposure for 40 days.

All statistical analyses were conducted using Prism GraphPad (version 8.1.2).

## Results and Discussion

The health and performance of aquatic organisms, particularly fish, are closely intertwined with the quality of their surrounding environment. Heavy metals like  $\text{Cd}^{2+}$  and  $\text{Pb}^{2+}$  enter aquatic ecosystems through various sources, including human activities, industry, and agriculture. The resulting toxicity from these heavy metals forces fish to adapt by adjusting both their behavioral and physiological activities to cope with encountered oxidative stress [31].

The concentration of  $\text{Cd}^{2+}$  and  $\text{Pb}^{2+}$ , both individually and in combination, in fish tissues is depicted in Fig. 1 after a 40-day exposure period. The results reveal a remarkable trend, indicating a significantly ( $P < 0.05$ ) higher concentration of these heavy metals in the fish tissues, specifically in the order of kidney > gills > intestine compared to the control group. Comparatively, in both fish gills and kidneys, the group treated with  $\text{Pb}^{2+}$  demonstrated a considerably ( $P < 0.05$ ) higher concentration compared to the group treated with  $\text{Cd}^{2+}$  individually. Conversely, in the intestine, no significant difference was observed between the  $\text{Cd}^{2+}$ -treated group and the  $\text{Pb}^{2+}$ -treated group. However, the concentration of heavy metals was markedly more pronounced when administered in a mixed form ( $\text{Cd}^{2+} + \text{Pb}^{2+}$  treated group) as opposed to their presence in a single form. Similar results were reported in other studies. Tunçsoy and Erdem [32] exposed *Oreochromis niloticus* to metals (Cd, Zn, and Co) both individually and in a mixture, while Qu et al. [33] found higher concentrations of

heavy metals in *Carassius auratus* tissues when exposed to a combination of metals. Different studies conducted in sublethal concentrations of  $\text{Pb}^{2+}$  and  $\text{Cd}^{2+}$ , such as Al-Balawi et al. [34], investigated the sublethal effects of  $\text{Pb}^{2+}$  on *Clarias gariepinus*, noting higher accumulation in the gills, liver, and kidney, with a lower level found in the muscle. In another study, Malarvizhi et al. [35] reported a higher concentration of  $\text{Cd}^{2+}$  in the gills, liver, and kidney of *Cirrhinus mrigala* after 30 days of exposure. In this study, the increased concentration of  $\text{Cd}^{2+}$  and  $\text{Pb}^{2+}$  in fish kidneys following sublethal exposure indicates a significant accumulation of these heavy metals within the renal tissues. This elevated concentration highlights the susceptibility of fish kidneys to the absorption and retention of  $\text{Cd}^{2+}$  and  $\text{Pb}^{2+}$  during sublethal exposure scenarios [36]. The accumulation and harmful impact of metals vary based on the specific species, developmental stage, gender, type of metal, its concentration, duration of exposure, and the physical and chemical attributes of the water [37, 38, 8].

The catalase CAT and SOD activities in fish liver following exposure to heavy metals, both individually and in combination, are illustrated in Fig. 2. The findings indicate that there is no significant difference in fish exposed to  $\text{Cd}^{2+}$  and  $\text{Pb}^{2+}$  when exposed individually. However, a significantly lower concentration was observed in the combined form compared to both individual exposures and the control group. These results align with the studies of Atli et al. [39], who subjected *Oreochromis niloticus* to sublethal levels of  $\text{Cd}^{2+}$  and  $\text{Pb}^{2+}$ . Rajeshkumar et al. [40] similarly investigated *Cyprinus carpio* after exposure to multiple heavy metals ( $\text{Cr}^{2+}$ ,  $\text{Cd}^{2+}$ , and  $\text{Pb}^{2+}$ ). The reduction in the activities of CAT and SOD in *C. marulius* after exposure




Fig. 2. CAT and SOD activity of control and metals treated groups of *Channa marulius*.

Table 1. Hemato-biochemical parameters of *Channa marulius* after exposure to heavy metals individually and in combined form for 40 days.

Parameters	Control	Treated-Cd	Treated-Pb	Treated-Cd +Pb
RBC count ( $10^6 \text{ mm}^{-3}$ )	$1.89 \pm 0.76^a$	$1.25 \pm 0.21^b$	$1.22 \pm 0.19^b$	$1.07 \pm 0.22^c$
Hct (%)	$35.42 \pm 0.92^a$	$24.74 \pm 0.47^b$	$21.37 \pm 0.73^b$	$12.43 \pm 0.42^c$
Hb (g/100 mL)	$7.23 \pm 1.25^a$	$3.42 \pm 0.16^b$	$3.31 \pm 0.28^b$	$3.12 \pm 0.23^b$
Cortisol (ng/l)	$16.06 \pm 1.33^c$	$47.32 \pm 7.67^b$	$49.73 \pm 2.44^b$	$57.32 \pm 2.06^a$
TP (g/100 ml)	$4.35 \pm 0.08^a$	$1.93 \pm 0.07^b$	$1.88 \pm 0.06^b$	$1.64 \pm 0.03^c$
Glucose (mg/L)	$63.11 \pm 2.17^c$	$84.26 \pm 2.52^b$	$87.42 \pm 2.73^b$	$98.64 \pm 2.33^a$
AST (IU/L)	$82.74 \pm 2.03^c$	$116.52 \pm 2.32^b$	$119.43 \pm 2.56^b$	$130.28 \pm 2.77^a$
ALT (IU/L)	$29.65 \pm 1.05^c$	$45.64 \pm 1.88^b$	$48.32 \pm 1.85^b$	$58.74 \pm 2.21^a$
Survival rate, %	100.00	94.29	91.43	85.71

Note: Statistical significance is denoted by different superscripts assigned to means within a similar row.






Fig. 3. Gill and liver AChE inhibition of control and metals treated groups of *Channa marulius*.

to  $\text{Cd}^{2+}$  and  $\text{Pb}^{2+}$  can be attributed to the harmful effects of these metals on the antioxidant defense system.  $\text{Cd}^{2+}$  and  $\text{Pb}^{2+}$  are known to induce oxidative stress by generating reactive oxygen species (ROS) within the organism [41, 42].

Hematological and biochemical indicators serve as valuable tools in assessing the health status of fish. They offer early detection capabilities for potential issues, aiding in the effective management of fish populations in both aquaculture and natural environments [43, 44]. Table 1 presents the hemato-biochemical parameters of *C. marulius* after a 40-day exposure to heavy metals, both individually and in combination. The fish group exposed to  $\text{Cd}^{2+}$  and  $\text{Pb}^{2+}$  in combined form exhibited significantly ( $P < 0.05$ ) lower concentrations of RBC count, Hb, and TP compared to those exposed to each heavy metal individually. Hct levels, however, showed no significant differences among all exposed groups, whether individually or in combined form. Similarly, Fazio et al. [5] found reduced RBC, Hb, Hct, and TP levels in *Mystus seenghala* following sublethal exposure to  $\text{Cd}^{2+}$ . The toxic effects of heavy metals on fish hematopoietic tissues, responsible for red blood cell production, were evident in our findings. Prolonged exposure to heavy metals can impede the normal functioning of the hematopoietic system, leading to a decline in red blood cell production [45-47].

In terms of cortisol, glucose, AST, and ALT levels, the fish group exposed to heavy metals in combined form demonstrated significantly ( $P < 0.05$ ) elevated values. Conversely, no significant differences were observed in the  $\text{Cd}^{2+}$  and  $\text{Pb}^{2+}$  treated groups when exposed individually. Survival rates indicated a higher percentage in the  $\text{Cd}^{2+}$ -treated group (94.29), followed by the  $\text{Pb}^{2+}$ -treated group (91.43) when exposed individually. However, in the combined form, the survival rate decreased to 85.71%. The studies, including Fazio et al. [5], demonstrated a substantial increase in glucose, cortisol, AST, and ALT levels in *M. seenghala* exposed to sublethal levels of  $\text{Cd}^{2+}$ .

#### Acetylcholinesterase Inhibition

AChE is crucial for cholinergic neurotransmission at neuromuscular junctions and brain synapses. It is produced in the endoplasmic reticulum and then processed in the Golgi apparatus, where it becomes either a membrane-bound enzyme or a secreted molecule [48-50]. Fig. 3 shows AChE activity in *C. marulius* exposed to  $\text{Cd}^{2+}$  and  $\text{Pb}^{2+}$ . The results indicate that AChE activity in both gill and liver tissues was significantly affected by exposure to  $\text{Cd}^{2+}$  and  $\text{Pb}^{2+}$ , whether individually or combined. The most substantial decrease in AChE activity was observed in fish exposed to both metals together, followed by those exposed to  $\text{Pb}^{2+}$  alone.

Interestingly, in liver tissue, there was no significant difference in AChE activity between the control group and the Cd<sup>2+</sup>-exposed group. However, fish exposed to both Cd<sup>2+</sup> and Pb<sup>2+</sup> in combination exhibited significantly lower AChE activity. Liu et al. [48] studied the neurotoxicity in zebrafish (*Danio rerio*) exposed to Pb<sup>2+</sup> and As, observing that combined exposure to these metals can lead to abnormal swimming behavior and exacerbate neurotoxicity.

### Conclusions

The findings from this study indicate that exposing *C. marulius* to a combination of metal forms results in a significant accumulation of these metals in fish tissues, causing more pronounced alterations in hemato-biochemical parameters compared to exposure to individual metals and control. Remarkably, the kidney exhibited the highest concentration of heavy metals, given its crucial role in filtration, reabsorption, and excretion processes, followed by the gills. The observed changes in hemato-biochemical parameters suggest a clear manifestation of physiological stress, characterized by rapid and transient modifications in specific stress-related indicators. With respect to neurotransmitter activity, Cd<sup>2+</sup> and Pb<sup>2+</sup> exposure notably reduced AChE activity in *C. marulius*. These research outcomes offer valuable insights for assessing pollution stress in aquatic environments and their inhabitants. This information is essential for shaping policies and strategies aimed at reducing the release of chemical compounds and heavy metals into freshwater ecosystems. To comprehensively understand the dynamics of aquatic organisms exposed to chemical mixtures in real-world scenarios, further research should explore the depuration potential of these organisms. Investigating the environmental realism of these scenarios will contribute to a more thorough understanding of the long-term impacts of chemical exposure on aquatic ecosystems.

### Acknowledgments

The authors extend their appreciation to the Researchers Supporting Project (RSPD2025R758), King Saud University, Riyadh, Saudi Arabia.

### Funding

This work was funded by the Researchers Supporting Project (RSPD2025R758), King Saud University, Riyadh, Saudi Arabia.

### Conflict of Interest

The authors declare no conflict of interest.

### References

1. NAZ S., FAZIO F., HABIB S.S., NAWAZ G., ATTAULLAH S., ULLAH M., HAYAT A., AHMED I. Incidence of heavy metals in the application of fertilizers to crops (wheat and rice), a fish (common carp) pond and a human health risk assessment. *Sustainability*, **14**, 13441, **2022**.
2. RAHMAN Z., SINGH V.P. The relative impact of toxic heavy metals (THMs) (arsenic (As), cadmium (Cd), chromium (Cr) (VI), mercury (Hg), and lead (Pb)) on the total environment: an overview. *Environmental Monitoring and Assessment*, **191**, 1, **2019**.
3. MADGETT A.S., YATES K., WEBSTER L., MCKENZIE C., MOFFAT C.F. The concentration and biomagnification of trace metals and metalloids across four trophic levels in a marine food web. *Marine Pollution Bulletin*, **173**, 112929, **2021**.
4. HABIB S.S., BATOOL A.I., REHMAN M.F.U., NAZ S. Assessment and bioaccumulation of heavy metals in fish feeds, water, and some tissues of *Cyprinus carpio* cultured in different environments (Biofloc Technology and Earthen Pond System). *Biological Trace Element Research*, **201**, 3474, **2023**.
5. FAZIO F., HABIB S.S., NAZ S., HASHMI M.A.H., SAOCHA C., ULLAH M. Cadmium sub-lethal concentration effect on growth, haematological and biochemical parameters of *Mystus seenghala* (Sykes, 1839). *Biological Trace Element Research*, **200**, 2432, **2022**.
6. YANG G., TANG Y., LIU X., WANG L., QIN L., LI D., SHEN X., KONG C., ZHAI W., FODJO EK., FAN C. Determination of Free Glycidol and Total Free Monochloropropanediol in Fish and Krill Oil with Simple Aqueous Derivatization and High-Performance Liquid Chromatography–Tandem Mass Spectrometry. *Foods*, **13**, 2340, **2024**.
7. TASLEEM S., MASUD S., HABIB S.S., NAZ S., FAZIO F., ASLAM M., ULLAH M., ATTAULLAH S. Investigation of the incidence of heavy metals contamination in commonly used fertilizers applied to vegetables, fish ponds, and human health risk assessments. *Environmental Science and Pollution Research*, **30**, 100646, **2023**.
8. SHAHJAHAN M., TASLIMA K., RAHMAN M.S., AL-EMRAN M., ALAM S.I., FAGGIO C. Effects of heavy metals on fish physiology—a review. *Chemosphere*, **300**, 134519, **2022**.
9. RIND K.H., HABIB S.S., UJAN J.A., FAZIO F., NAZ S., BATOOL A.I., ULLAH M., ATTAULLAH S., KHAYYAM K., KHAN K. The effects of different carbon sources on water quality, growth performance, hematology, immune, and antioxidant status in cultured Nile Tilapia with biofloc technology. *Fishes*, **8**, 512, **2023**.
10. HABIB S.S., NAZ S., NAWAZ S., AMEER I., KHATOON A., REHMAN H.U., JAWAD S.M., ALI H. Comparative analysis of hematological parameters of some farmed and wild fish species. *Pakistan Journal of Zoology*, **54**, 591, **2022**.
11. HEYS K.A., SHORE R.F., PEREIRA M.G., JONES K.C., MARTIN F.L. Risk assessment of environmental mixture effects. *RSC Advances*, **6**, 47844, **2016**.
12. GONG B., QIU H., ROMERO-FREIRE A., VAN GESTEL C.A., HE E. Incorporation of chemical and toxicological availability into metal mixture toxicity modeling: State

- of the art and future perspectives. *Critical Reviews in Environmental Science and Technology*, **52**, 1730, **2022**.
13. KARRI V., SCHUEMACHER M., KUMAR V. Heavy metals (Pb, Cd, As and MeHg) as risk factors for cognitive dysfunction: A general review of metal mixture mechanism in brain. *Environmental Toxicology and Pharmacology*, **48**, 203, **2016**.
  14. BEISHON L.C., HOSFORD P., GURUNG D., BRASSARD P., MINHAS J.S., ROBINSON T.G., HAUNTON V., PANERAI R.B. The role of the autonomic nervous system in cerebral blood flow regulation in dementia: A review. *Autonomic Neuroscience*, **240**, 102985, **2022**.
  15. ABUBAKAR M.U., ABUBAKAR D. Characterization of acetylcholinesterase from various sources: a mini review. *Journal of Environmental Bioremediation and Toxicology*, **4**, 24, **2021**.
  16. ROTUNDO R.L. Acetylcholinesterase at the neuromuscular junction. *Neuroscience Letters*, **735**, 135157, **2020**.
  17. ORTEGA D.R., ESQUIVEL D.F.G., AYALA T.B., PINEDA B., MANZO S.G., QUINO J.M., MORA P.C., DE LA CRUZ V.P. Cognitive impairment induced by lead exposure during lifespan: Mechanisms of lead neurotoxicity. *Toxics*, **9**, 23, **2021**.
  18. LV Z., HU J., HUANG M., PAN G., XU G., YANG M. Molecular mechanisms of cadmium-induced cytotoxicity in human ovarian granulosa cells identified using integrated omics. *Ecotoxicology and Environmental Safety*, **272**, 116026, **2024**.
  19. NAZIR S., KHAN N., FATIMA M., AZMAT H., NAVEED S., RAMZAN M.M., ASGHAR M., BANO S., KHIZER A., WAN A.H., DAVIES S.J. The influence of dietary protein concentration on digestive enzyme activities, growth, and body composition in juvenile bullseye snakehead (*Channa marulius*). *Plos one*, **18**, e0281274, **2023**.
  20. AHMED S.F., KUMAR P.S., KABIR M., ZUHARA F.T., MEHJABIN A., TASANNUM N., HOANG A.T., KABIR Z., MOFIJUR M. Threats, challenges and sustainable conservation strategies for freshwater biodiversity. *Environmental Research*, **214**, 113808, **2022**.
  21. BATOOL M., ABDULLAH S., IJAZ M.U., KOUSAR S., FATIMA M., ILYAS R., MUGHAL K.T. Heavy metals (cadmium and lead) induced oxidative stress in *Channa marulius* and *Wallago attu* during acute toxicity experiments. *Pakistan Journal of Zoology Supplementary Series*, **13**, 74, **2018**.
  22. American Public Health Association (APHA), American Water Works Association (AWWA), Water Environment Federation (WEF) Standard methods for the examination of water and wastewater, 22nd edition, **2012**.
  23. HABIB S.S., NAZ S., FAZIO F., CRAVANA C., ULLAH M., RIND K.H., ATTAULLAH S., FILICIOTTO F., KHAYYAM K. Assessment and bioaccumulation of heavy metals in water, fish (wild and farmed) and associated human health risk. *Biological Trace Element Research*, **202**, 725, **2024**.
  24. HABIB S.S., SAEED M.Q., MASUD S., KESBİÇ O.S., UJAN J.A., CRAVANA C., AL-REJAIE S.S., MOHANY M., FAZIO F. Dietary Effect of *Withania somnifera* Root Powder on Growth, Hematobiochemical Parameters, Immunity, and Disease Resistance to *Aeromonas hydrophila* in *Cyprinus carpio*. *Aquaculture Nutrition*, **2024**, 7321287, **2024**.
  25. HABIB S.S., BATOOL A.I., REHMAN M.F.U., NAZ S. Comparative analysis of hemato-biochemical profile, growth performance, and body composition of Common Carp cultured under different feed and habitats (biofloc technology and earthen pond systems). *North American Journal of Aquaculture*, **85**, 277, **2023**.
  26. AMOURI R.E., UJAN J.A., RAZA A., MUSHTAQ A., SAEED M.Q., MASUD S., HABIB S.S., MILOŠEVIĆ M., AL-REJAIE S.S., MOHANY M. Sublethal Concentrations of Cadmium and Lead: Effects on Hemato-Biochemical Parameters and Tissue Accumulation in *Wallago attu*. *Biological Trace Element Research*, **1**, **2024**.
  27. TRINDER P. Determination of glucose in blood using glucose oxidase with an alternative oxygen acceptor. *Annals of Clinical Biochemistry*, **6**, 24, **1969**.
  28. HENRY R.J. Colorimetric determination of total protein: clinical chemistry. Harper and Row, New York. **1964**.
  29. REITMAN S., FRANKEL S. A colorimetric method for the determination of serum glutamic oxalacetic and glutamic pyruvic transaminases. *American Journal of Clinical Pathology*, **28**, 56, **1957**.
  30. PRETTO A., LORO V.L., MORSCH V.M., MORAES B.S., MENEZES C., CLASEN B., HOEHNE L., DRESSLER V. Acetylcholinesterase activity, lipid peroxidation, and bioaccumulation in silver catfish (*Rhamdia quelen*) exposed to cadmium. *Archive of Environmental Contamination and Toxicology*, **58**, 1008, **2010**.
  31. SABULLAH M.K., AHMAD S.A., SHUKOR M.Y., GANSAU A.J., SYED M.A., SULAIMAN M.R., SHAMAAN N.A. Heavy metal biomarker: Fish behavior, cellular alteration, enzymatic reaction and proteomics approaches. *International Food Research Journal*, **22**, 435, **2015**.
  32. TUNÇSOY M., ERDEM C. Accumulation of copper, zinc and cadmium in liver, gill and muscle tissues of *Oreochromis niloticus* exposed to these metals separately and in mixture. *Fresenius Environmental Bulletin*, **23**, 1143, **2014**.
  33. QU R., WANG X., WANG Z., WEI Z., WANG L. Metal accumulation and antioxidant defenses in the freshwater fish *Carassius auratus* in response to single and combined exposure to cadmium and hydroxylated multi-walled carbon nanotubes. *Journal of Hazardous Materials*, **275**, 89, **2014**.
  34. AL-BALAWI H.F.A., AL-AKEL A.S., AL-MISNED F., SULIMAN E.A.M., AL-GHANIM K.A., MAHBOOB S., AHMAD Z. Effects of sub-lethal exposure of lead acetate on histopathology of gills, liver, kidney and muscle and its accumulation in these organs of *Clarias gariepinus*. *Brazilian Archives of Biology and Technology*, **56**, 293, **2013**.
  35. MALARVIZHI A., SARAVANAN M., POOPAL R.K., HUR J.H., RAMESH M. Accumulation of cadmium and antioxidant and hormonal responses in the Indian Major Carp *Cirrhinus mrigala* during acute and sublethal exposure. *Water Air Soil Pollution*, **228**, 1, **2017**.
  36. ORR S.E., BRIDGES C.C. Chronic kidney disease and exposure to nephrotoxic metals. *International Journal of Molecular Sciences*, **18**, 1039, **2017**.
  37. UNGUREANU E.L., MUSTATEA G. Toxicity of heavy metals. In: *Environmental Impact and Remediation of Heavy Metals*. IntechOpen, **2022**.
  38. LIU X., DAI B., CHUAI Y., HU M., ZHANG H. Associations between vitamin D levels and periodontal attachment loss. *Clinical Oral Investigations*, **27**, 4727, **2023**.
  39. ATLI G., CANLI E.G., EROGLU A., DOGAN Z.,



- CANLI M. Cadmium and lead alter the antioxidant and osmoregulation systems in the erythrocyte of fish (*Oreochromis niloticus*). *Turkish Journal of Fisheries and Aquatic Sciences*, **16**, 361, **2022**.
40. RAJESHKUMAR S., LIU Y., MA J., DUAN H.Y., LI X. Effects of exposure to multiple heavy metals on biochemical and histopathological alterations in common carp, *Cyprinus carpio* L. *Fish Shellfish and Immunology*, **70**, 461, **2017**.
41. UNSAL V., DALKIRAN T., ÇIÇEK M., KÖLÜKÇÜ E. The role of natural antioxidants against reactive oxygen species produced by cadmium toxicity: a review. *Advanced Pharmaceutical Bulletin*, **10**, 184, **2020**.
42. CHUAI Y., DAI B., LIU X., HU M., WANG Y., ZHANG H. Association of vitamin K, fibre intake and progression of periodontal attachment loss in American adults. *BMC Oral Health*, **23**, 303, **2023**.
43. FAZIO F. Fish hematology analysis as an important tool of aquaculture: a review. *Aquaculture*, **500**, 237, **2019**.
44. HU R., FENG H. Lenticulostrate artery and lenticulostrate-artery neural complex: new concept for intracerebral hemorrhage. *Current Pharmaceutical Design*, **23**, 2206, **2017**.
45. SCHARF P., BROERING M.F., OLIVEIRA DA ROCHA G.H., FARSKY S.H.P. Cellular and molecular mechanisms of environmental pollutants on hematopoiesis. *International Journal of Molecular Sciences*, **21**, 6996, **2020**.
46. ZHEN W., WANG Z., WANG Q., SUN W., WANG R., ZHANG W., ZHANG Y., QIN W., LI B., WANG Q., HONG, B. Cardiovascular disease therapeutics via engineered oral microbiota: Applications and perspective. *iMeta*, **3**, e197, **2024**.
47. HUI M., JIA X., LI X., LAZCANO-SILVEIRA R., SHI M. Anti-Inflammatory and Antioxidant Effects of Liposoluble C60 at the Cellular, Molecular, and Whole-Animal Levels. *Journal of Inflammation Research*, **16**, 83, **2023**.
48. BUKHARAEVA E., KHUZAKHMETOVA V., DMITRIEVA S., TSENTSEVITSKY A. Adrenoceptors modulate cholinergic synaptic transmission at the neuromuscular junction. *International Journal of Molecular Sciences*, **22**, 4611, **2021**.
49. WANG Z., SUN W., HUA R., WANG Y., LI Y., ZHANG H. Promising dawn in tumor microenvironment therapy: engineering oral bacteria. *International Journal of Oral Science* **16**, 24, **2024**.
50. LIU M., DENG P., LI G., LIU H., ZUO J., CUI W., ZHANG H., CHEN X., YAO J., PENG X., PENG L. Neurotoxicity of combined exposure to the heavy metals (Pb and As) in zebrafish (*Danio rerio*). *Toxics*, **12**, 282, **2024**.